

A new rheophytic species of *Syzygium* Gaertn. (Myrtaceae) from Assam, North East India

Jatindra SARMA, Hussain Ahmed BARBHUIYA & Santanu DEY



DIRECTEUR DE LA PUBLICATION : Bruno David
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF* : Thierry Deroin

RÉDACTEURS / *EDITORS* : Porter P. Lowry II ; Zachary S. Rogers

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS* : Emmanuel Côté (adanson@mnhn.fr) ; Anne Mabilie

MISE EN PAGE / *PAGE LAYOUT* : Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD* :

P. Baas (Nationaal Herbarium Nederland, Wageningen)
F. Blasco (CNRS, Toulouse)
M. W. Callmender (Conservatoire et Jardin botaniques de la Ville de Genève)
J. A. Doyle (University of California, Davis)
P. K. Endress (Institute of Systematic Botany, Zürich)
P. Feldmann (Cirad, Montpellier)
L. Gautier (Conservatoire et Jardins botaniques de la Ville de Genève)
F. Ghahremaninejad (Kharazmi University, Téhéran)
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)
K. Kubitzki (Institut für Allgemeine Botanik, Hamburg)
J.-Y. Lesouef (Conservatoire botanique de Brest)
P. Morat (Muséum national d'Histoire naturelle, Paris)
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)
S. E. Rakotoarisoa (Millenium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)
É. A. Rakotobe (Centre d'Applications des Recherches pharmaceutiques, Antananarivo)
P. H. Raven (Missouri Botanical Garden, St. Louis)
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)
J. G. West (Australian National Herbarium, Canberra)
J. R. Wood (Oxford)

COUVERTURE / *COVER* :

Syzygium nivae Barbhuiya, J. Sarma & S. Dey, sp. nov., inflorescence and fruits (photos by J. Sarma).

Adansonia est indexé dans / *Adansonia is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Adansonia est distribué en version électronique par / *Adansonia is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Adansonia est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Adansonia is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:

Geodiversitas, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Cryptogamie* sous-sections Algologie, Bryologie, Mycologie.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél. : 33 (0)1 40 79 48 05 / Fax : 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2019
ISSN (imprimé / *print*) : 1280-8571/ ISSN (électronique / *electronic*) : 1639-4798

A new rheophytic species of *Syzygium* Gaertn. (Myrtaceae) from Assam, North East India

Jatindra SARMA

Development, Research, Education and Working Plans, Assam Forest Department,
Aranya Bhawan, Panjabari, Guwahati 781037, Assam (India)

Hussain Ahmed BARBHUIYA

Landscape and Cosmetic Maintenance Section,
Architectural and Structural Engineering Division,
Bhabha Atomic Research Centre, Trombay, Mumbai 400085, Maharashtra (India)
hahmed@barc.gov.in

Santanu DEY

Department of Botany, Nagaland University, Lumami 798627, Nagaland (India)

Submitted on 14 July 2018 | accepted on 14 November 2018 | published on 20 May 2019

Sarma J., Barbhuiya H. A. & Dey S. 2019. — A new rheophytic species of *Syzygium* Gaertn. (Myrtaceae) from Assam, North East India. *Adansonia*, sér. 3, 41 (6): 53-58. <https://doi.org/10.5252/adansonia2019v41a6>. <http://adansonia.com/41/6>

ABSTRACT

A new species of *Syzygium* Gaertn. (Myrtaceae) is described from North East India: *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov. The new species is a rheophyte, which is restricted to the Kopili riverbed of West Karbi Anglong district of Assam. The present species shows little affinity with *Syzygium khasianum* (Duthie) N.P.Balakr. in having lanceolate leaves, panicle inflorescence and calyptrate petals, but differs in various aspects, such as shrubby rheophytic plant habit, narrower leaves, non-caudate leaf apex, shorter peduncles, larger flowers etc.

RÉSUMÉ

Une nouvelle espèce rhéophytique de *Syzygium* Gaertn. (Myrtaceae) d'Assam, nord-est de l'Inde.

Une espèce nouvelle de *Syzygium* Gaertn. (Myrtaceae) est décrite du nord-est de l'Inde: *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov. Cette espèce est une rhéophyte, circonscrite aux rives de la rivière Kopili, dans le district Karbi Anglong Ouest (Assam). Elle montre une faible affinité avec *S. khasianum* (Duthie) N.P. Balakr. par ses feuilles lancéolées, son inflorescence paniculée et ses pétales calyptrés, mais en diffère par divers caractères, tels que son port arbustif rhéophytique, ses feuilles plus étroites à apex non caudés, ses pédoncules plus courts, ses fleurs plus grandes, etc.

KEY WORDS

Myrtaceae,
Syzygium,
rheophyte,
endemic,
West Karbi Anglong,
Assam.

MOTS CLÉS

Myrtaceae,
Syzygium,
rhéophyte,
endémique,
Karbi Anglong Ouest,
Assam.

INTRODUCTION

Syzygium Gaertn. is the largest genus in Myrtaceae (clove family), which comprises *c.* 1200 species (Soh & Parnell 2015). It is a paleotropical genus with a wide range of occurrence mainly in southern and southeastern Asia, Australia, Malesia, and New Caledonia. Some species occur in east Africa, Madagascar, the Mascarenes, southwestern Pacific Islands, Taiwan, and southern Japan (Chen & Craven 2007; Soh 2017). Duthie (1878-1879), in Hooker's *Flora of British India*, treated the genus under *Eugenia* L. *sensu lato*, with 131 species which has been split into genera *Eugenia* and *Syzygium* with the majority of Indian species now placed in the latter genus (Byng *et al.* 2015). The genus shows high diversity in North Eastern India and Western Ghats. Kanjilal *et al.* (1938) enumerated *c.* 35 species from North East India under *Eugenia sensu lato* and Sujanapal & Kunhikannan (2017) reported *c.* 48 species of *Syzygium* from the Western Ghats. In India, the genus has received very little attention as till date there is no comprehensive revisionary work available for the area, except the discovery of few new species in recent years (Viswanathan & Manikandan 2008; Shareef *et al.* 2012, 2013, 2014; Ratheesh *et al.* 2014; Sujanapal *et al.* 2014; Venkat Ramana *et al.* 2014) and it is hoped that many more novel species awaits their formal scientific discovery.

During field explorations (2016-2018) at Koka, Panimur, Amreng and Zirikindeng regions of West Karbi Anglong District of Assam the senior author found this rheophytic shrub growing on rock crevices at the edges of Kopili riverbed near Koka. Observations in the field and critical examination of the voucher specimen have revealed number of morphological differences from all other hitherto known species of genus *Syzygium*, which warrants the description of a new species.

SYSTEMATICS

Syzygium nivae Barbhuiya, J.Sarma & S.Dey, sp. nov.
(Figs 1; 2)

The new species differs from *S. khasianum* (Duthie) N.P.Balacr. for its shrubby rheophytic plant habit, narrower leaves, non-caudate leaf apex, shorter peduncles, larger flowers and oblong to subglobose fruits.

TYPUS. — **India.** Assam, West Karbi Anglong District, Panimur, Koka, on the edges of Kopili riverbed, 25°43'–56.6"N, 92°49'–19.31"E, *c.* 90 m, 17.VIII.2016, fl., J. Sarma *s.n.* (holo-, HBARC[HBARC00006399]).

ETYMOLOGY. — The species is named after Mrs Niva Deka, the wife of first author of the paper, who constantly encouraged her husband to continue his botanical explorations in remote area of the state of Assam. By profession, Mrs Deka is an Engineer and presently working under the government of Assam.

DISTRIBUTION. — Endemic to the type locality, Koka, Panimur, West Karbi Anglong District, Assam, India (Fig. 3).

PHENOLOGY. — The plant flowers during the months of August to October and its fruit matures in October and November.

CONSERVATION STATUS. — *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov. is provisionally categorized as 'Data Deficient' (DD), as only 50 mature individuals were located during the survey at Koka locality. To ascertain its actual status, the entire course of Kopili River is required to be surveyed.

HABITAT AND ECOLOGY. — *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov. grows in rock crevices along the Kopili riverbed, at altitudes 86-102 m above MSL. The riverbed is characterized by the alternating bands of shale and sandstones. The sandstone are mainly composed of quartz, lithic fragments with meager amount of feldspar and cemented by calcareous and ferruginous cements (Bhuyan 2016). The rate of water flow of the river is very high during June-July and gradually slowed down from August, when the plant starts flowering. The other associated rheophytes occurring in the type locality were *Syzygium cyanophyllum*, *S. polypetalum*, *Ixora yunnanensis*, *Pavetta puffii*, *Carissa* sp., *Tarenna pumila*, *Eriobotrya angustissima*, *Millettia* sp., *Melastoma malabathricum* etc.

TABLE 1. — Morphological differences between *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov. and *S. khasianum* (Duthie) N.P.Balacr.

Characters	<i>Syzygium nivae</i> Barbhuiya, J.Sarma & S.Dey, sp. nov.	<i>Syzygium khasianum</i> (Duthie) N.P.Balacr.
Habit	A shrub, to 2.5 m high	A middle-size spreading tree
Leaf blade	narrowly lanceolate, 3.1-6.6 × 0.8-1.1 cm	ovate lanceolate, 5.0-7.5 × 1.1-2.7 cm
Leaf apex	acuminate, not caudate	abruptly acuminate, shortly caudate
Leaf base	attenuate	cuneate or attenuate
Lateral nerves	numerous, 0.5-1.6 mm apart	numerous, 0.4-1.4 mm apart
Petiole	8.3-11 mm long	8.0-17 mm long
Inflorescence	paniculate-cyme, terminal and axillary, to 12 cm long and 6.8 cm in diam.	paniculate-cyme, terminal and axillary, 10-14 cm long and 6-10 cm in diam.
Peduncle	1.8-2.5 cm long	5.0-7.5 cm long
Pedicel	absent or to 0.6 mm long	absent or to 1 mm long
Flower bud	pyriform, 5-6 mm long	pyriform, <i>c.</i> 3.8 mm long
Flowers	10-12 mm in across	<i>c.</i> 6.3 mm in across
Hypanthium	obconic, 4.9-7.4 mm long and 4.2-5.2 mm wide at mouth	obconic, 3.2-4.7 mm long and 2.5-3.5 mm wide at mouth
Petals	4, calyptrate, sub-orbicular, bowl-shaped, 2.4-3.0 mm in diam.	4, calyptrate, rarely free, bowl-shaped, 1.5-2.0 mm in diam.
Style	thickened at base slightly narrowed upwards, <i>c.</i> 8.3 mm long	uniform, <i>c.</i> 6.5 mm long
Fruit	slightly oblong or subglobose, 10-16 mm long	cylindric <i>c.</i> 6.3 mm long
Distribution	India (Assam: West Karbi Anglong District) (Fig. 3)	India (Meghalaya: Garo & Khasi Hills) (Fig. 3)




FIG. 1. — *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov.: **A**, type locality showing plant habit and habitat; **B**, inflorescence; **C**, flowers; **D**, fruits (photos by J. Sarma).



FIG. 2. — *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov.: **A**, apex of a branch showing inflorescence and leaves; **B**, a portion of the stem showing arrangement of leaves; **C**, leaves; **D**, leaf base showing petiole; **E**, middle portion of a leaf showing reticulate venation; **F**, leaf apex; **G**, a portion of the inflorescence showing flower buds; **H**, a portion of the inflorescence showing flowers; **I**, an opened flower bud; **J**, calyptra; **K**, L.S. of the flower; **L**, L.S. of hypanthium and gynoecium; **M**, petals; **N**, stamens; **O**, T.S. of calyx tube and style; **P**, T.S. of ovary (photos by H. A. Barbhuiya, based on holotype). Scale bars: A, C, 2 cm; B, G, H, 5 mm; D-F, 1 cm; I-N, 3 mm; O, 2 mm; P, 1 mm.




FIG. 3. — Distribution map of *Syzygium nivae* Barbhuiya, J.Sarma & S.Dey, sp. nov., *S. khasianum* (Duthie) N.P.Balakr. and *S. cyanophyllum* (P.C.Kanjilal & Das) Raizada. Map Created with SimpleMappr, <http://www.simplemappr.net> (Shorthouse 2010).

DESCRIPTION

A rheophytic shrub, to 2.5 m high; branchlets terete. Leaves opposite decussate, coriaceous; blade narrowly lanceolate, 3.1–6.6 × 0.8–1.1 cm, margins revolute, finely reticulate, both surfaces glabrous, deep green above, pale-green beneath, apex acuminate or obtuse, rarely retuse, base attenuate; midvein subcanaliculate above, raised beneath; lateral veins numerous, obscure, brochidodromous, 0.5–1.6 mm apart; intramarginal vein looped, 0.5–0.8 mm away from the leaf margin; petiole yellowish, 8.3–11 mm long, glabrous. Inflorescence a panicle-cyme, terminal and axillary at upper leaf axils, to 12 cm long and *c.* 6.8 cm in diam.; branches quadrangular, green, opposite spreading, 1.4–2.5 mm in diam.; peduncle subquadrangular, 1.8–2.5 cm long. Flowers sessile or shortly pedicellate; pedicel, to 0.6 mm long. Flower bud pyriform, 5–6 mm long, flowers 10–12 mm in across during anthesis; hypanthium obconic, 4.9–7.4 mm long and 4.2–5.2 mm wide at apex. Calyx tube *c.* 3.1 mm long and up to 5.2 mm in diam.; lobes inconspicuous. Petals 4, falling as a calyptra, suborbicular, bowl-shaped 2.4–3.0 mm in diam. Stamens many, of different lengths; filaments cylindric, narrowed at apex, 4.0–8.6 mm long and 0.1–0.3 mm in diam.; anthers minute, versatile, 0.4–0.6 × 0.2–0.4 mm. Ovary fleshy *c.* 3.1 × 1.7 mm, 2-locular, glabrous; style *c.* 8.3 mm long, 0.2–0.5 mm in diam., cylindric, thickened at base, narrowed upwards; stigma inconspicuous. Fruits slightly oblong or subglobose, 1.0–1.6 × 0.8–1.3 cm, apex with cup-shaped limb, reddish when young, blackish on ripening and slightly sweeter in taste.

REMARKS

Syzygium nivae Barbhuiya, J.Sarma & S.Dey, sp. nov. shows superficial similarity with *S. khasianum* (Duthie) N.P.Balakr.

But the latter is quite distinct from the former by its arboreal habit, abruptly acuminate and caudate leaf apex, longer peduncles, much smaller flowers, etc. The detailed comparison between the two species is summarized in the Table 1. The other closely related species occurring in the same locality was *S. cyanophyllum* (P.C.Kanjilal & Das) Raizada, which can be easily distinguished from the newly described species by its smaller leaves, shortly corymbose inflorescence bearing flowers with long hypanthium.

Acknowledgements

We are thankful to the Assam Forest Department for allowing us to use their resources in extensive field exploration works especially in remote areas. We extend our sincere thanks to Karbi Anglong Autonomous Council Authority and Hamren Territorial Division for providing logistic support during field tours. We acknowledge our heartiest thanks to officers and staffs of Southern Range Kheroni, Hamren Territorial Division for their continuous assistance in frequent field explorations. Finally yet importantly, we are indebted to Mr. Rowelson Engti, Forester I, Southern Range Kheroni, for his valuable help.

REFERENCES

- BHUYAN D. 2016. — Late Eocene unstable basin margin Kopili sedimentation and its reservoir characteristics in upper Assam petroliferous basin. *International Journal of Research and Scientific Innovation* 3 (2): 10–15.
- BYNG J. W., WILSON P. G. & SNOW N. 2015. — Typifications and nomenclatural notes on Indian Myrtaceae. *Phytotaxa* 217 (2): 101–116. <https://doi.org/10.11646/phytotaxa.217.2.1>

- CHEN J. & CRAVEN L. A. 2007. — Myrtaceae, in WU Z. Y., RAVEN P. H. & HONG D. Y. (eds), *Flora of China*. Vol. 13. *Clusiaceae through Araliaceae*. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis: 321-359.
- DUTHIE J. F. 1878-1879. — Order LIX. Myrtaceae, in HOOKER J. D. (ed.), *Flora of British India*. Vol. 2. L. Reeve & Co., London: 462-506.
- KANJILAL U. N., KANJILAL P. C. & DAS A. 1938. — *Flora of Assam*. Vol. II. Government of Assam, Shillong, v + 409 p.
- RATHEESH N., SHAREEF S. M., SHAJU T., SIVU A. R., SUJANA K. A., NANDAKUMAR M. K. & SATHEESH K. T. 2014. — A new species of *Syzygium* (Myrtaceae) from the southern Western Ghats of Kerala, India. *International Journal of Advanced Research* 2 (3): 1055-1058.
- SOH W. K. 2017. — Taxonomy of *Syzygium*, in NAIR K. N. (ed), *The Genus Syzygium; Syzygium cumini and Other Underutilized Species*. CRC Press, New York: 1-5.
- SOH W. K. & PARNELL J. 2015. — A revision of *Syzygium* Gaertn. (Myrtaceae) in Indochina (Cambodia, Laos and Vietnam). *Adansonia*, sér. 3, 37 (2): 179-275. <https://doi.org/10.5252/a2015n2a1>
- SHAREEF S. M., ROY P. E. & KRISHNARAJ M. V. 2014. — *Syzygium munnarensis* sp. nov. (Myrtaceae): an overlooked endemic species from southern Western Ghats of Kerala, India. *Webbia* 69: 53-57. <https://doi.org/10.1080/00837792.2014.899727>
- SHAREEF S. M., SANTHOSH KUMAR E. S. & ROY P. E. 2013. — A new species of *Syzygium* (Myrtaceae) from Kerala, India. *Phytotaxa* 129 (1): 34-38. <https://doi.org/10.11646/phytotaxa.129.1.2>
- SHAREEF S. M., SANTHOSH KUMAR E. S. & SHAJU T. 2012. — A new species of *Syzygium* (Myrtaceae) from the southern Western Ghats of Kerala, India. *Phytotaxa* 71: 28-33. <https://doi.org/10.11646/phytotaxa.71.1.5>
- SHORTHOUSE D. P. 2010. — SimpleMappr, an online tool to produce publication-quality point maps. Retrieved from <http://www.simplemappr.net> (accessed 14 Jul 2018).
- SUJANAPAL P. & KUNHIKANNAN C. 2017. — The Genus *Syzygium* in Western Ghats, in NAIR K. N. (ed), *The Genus Syzygium; Syzygium cumini and Other Underutilized Species*. CRC Press, New York: 15-56.
- SUJANAPAL P., ROBI A. J. & SASIDHARAN N. 2014. — *Syzygium sahyadricum* (Myrtaceae), a new tree species from India, and notes on the distribution of *S. spathulatum* Thwaites. *Phytotaxa* 174 (5): 285-290. <https://doi.org/10.11646/phytotaxa.174.5.5>
- VENKAT RAMANA M., CHORGHE A. & VENU P. 2014. — Two new species of *Syzygium* (Myrtaceae) from Saddle Peak National Park, Andaman and Nicobar Islands, India. *Blumea* 59: 42-48. <https://doi.org/10.3767/000651914X683593>
- VISWANATHAN M. B. & MANIKANDAN U. 2008. — A new species of *Syzygium* (Myrtaceae) from the Kalakkad-Mundanthurai Tiger Reserve in Peninsular India. *Adansonia*, sér. 3, 30 (1): 113-118.

*Submitted on 14 July 2018;
accepted on 14 November 2018;
published on 20 May 2019.*