


Lectotypification of two Turkish endemic taxa of *Klasea* Cass. (Asteraceae, Cardueae)

Kazem NEGARESH



DIRECTEUR DE LA PUBLICATION / *PUBLICATION DIRECTOR*: Gilles Bloch
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Thierry Deroin

RÉDACTEURS / *EDITORS*: Porter P. Lowry II; Zachary S. Rogers; Mathieu Gardère

ÉDITEUR TECHNIQUE (SUIVI ÉDITORIAL) / *DESK EDITOR (EDITORIAL PROCESS)*: Emmanuel Côté (adanson@mnhn.fr)

ÉDITEUR TECHNIQUE (PRODUCTION) / *DESK EDITOR (PRODUCTION)*: Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

F. Blasco (CNRS, Toulouse)
M. W. Callmender (Conservatoire et Jardin botaniques de Genève)
J. A. Doyle (University of California, Davis)
P. K. Endress (Institute of Systematic Botany, Zürich)
P. Feldmann (Cirad, Montpellier)
L. Gautier (Conservatoire et Jardin botaniques de Genève)
F. Ghahremaninejad (Kharazmi University, Téhéran)
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)
A. A. Khapugin (Tyumen State University, Russia)
J.-Y. Lesouef (Conservatoire botanique de Brest)
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)
S. E. Rakotoarisoa (Millenium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)
P. H. Raven (Missouri Botanical Garden, St. Louis)
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)
J. G. West (Australian National Herbarium, Canberra)
J. R. Wood (Oxford)

COUVERTURE / *COVER*:

Réalisée à partir des Figures de l'article/*Made from the Figures of the article.*

Adansonia est indexé dans / *Adansonia is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Adansonia est distribué en version électronique par / *Adansonia is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Adansonia est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Adansonia is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*: *Geodiversitas*, *Zoosystema*, *Anthropozoologica*,
European Journal of Taxonomy, *Naturae*, *Cryptogamie* sous-sections *Algologie*, *Bryologie*, *Mycologie*, *Comptes Rendus Palevol*

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)
Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

Les articles publiés dans *Adansonia* sont distribués sous licence CC-BY 4.0/*Articles published in Adansonia are distributed under a CC-BY 4.0 license.*
ISSN (imprimé / *print*): 1280-8571/ ISSN (électronique / *electronic*): 1639-4798

Lectotypification of two Turkish endemic taxa of *Klasea* Cass. (Asteraceae, Cardueae)

Kazem NEGARESH

Department of Horticultural Science, Faculty of Agriculture,
Agricultural Sciences and Natural Resources University of Khuzestan, Mollasani (Iran)
negaresh@asnrukh.ac.ir

Submitted on 8 April 2025 | accepted on 26 May 2025 | published on 27 January 2026

KEY WORDS

Asteraceae,
Klasea,
Serratula,
Schumeria,
Türkiye,
Turkey,
endemic,
lectotypifications.

MOTS CLÉS

Asteraceae,
Klasea,
Serratula,
Schumeria,
Turquie,
endémisme,
lectotypifications.

Negaresh K. 2026. — Lectotypification of two Turkish endemic taxa of *Klasea* Cass. (Asteraceae, Cardueae). *Adansonia*, sér. 3, 48 (2): 7-11. <https://doi.org/10.5252/adansonia2026v48a2>. <http://adansonia.com/48/2>

ABSTRACT

Two Turkish endemic taxa of *Klasea* Cass. (Asteraceae) are lectotypified, namely *Klasea aznavouriana* (Bornm.) Greuter & Wagenitz and *K. bornmuelleri* (Azn.) Greuter & Wagenitz. Lectotypes are deposited at B and G respectively. In addition, the relevant synonyms, type details and images of the type specimens are provided.

RÉSUMÉ

Lectotypification de deux taxons endémiques turcs du genre Klasea Cass. (Asteraceae, Cardueae).
Deux taxons endémiques turcs du genre *Klasea* Cass. (Asteraceae) sont lectotypifiés, à savoir *Klasea aznavouriana* (Bornm.) Greuter & Wagenitz et *K. bornmuelleri* (Azn.) Greuter & Wagenitz. Les lectotypes correspondants sont déposés à B et G. En outre, les synonymes pertinents, les détails du type et les images des spécimens types sont fournis.

INTRODUCTION

Klasea Cass., a genus in the subtribe Centaureinae of the tribe Cardueae in Asteraceae Bercht. & J.Presl (Hellwig 2004; Martins & Hellwig 2005; Martins 2006; Susanna & Garcia-Jacas 2007, 2009), comprises over 65 species that chiefly occur in Europe, the Mediterranean region, south-western Asia and northern Africa (Susanna & Garcia-Jacas 2007). According to modern taxonomy, the genus includes eight sections in Turkey: *Klasea* sect. *Coriaceae* L.Martins, *K.* sect. *Demetria* (Boriss.) L.Martins, *K.* sect. *Lasiocephalae* L.Martins, *K.* sect. *Klasea*, *K.* sect. *Melanolepis* L.Martins, *K.* sect. *Leuzeopsis* (Boriss.) L.Martins, *K.* sect. *Quinquefoliae* L.Martins and *K.* sect. *Schumeria* (Iljin) L.Martins (Martins 2006). It has been reported that *Klasea* contains 15 species in Turkey (Davis & Kupicha 1975; Martins 2006; Dogan *et al.* 2015). Five of these species are endemic to Turkey, resulting in an endemism ratio of 33.3%. The genus includes unarmed perennial herbs with involucre bracts mostly lacking apical appendages, tubular florets, and glabrous achenes, and a simple, seriate pappus composed of free, long bristles (Martins 2006; Susanna & Garcia-Jacas 2007).

Continuing my taxonomic study of the names in *Klasea* (Ranjbar *et al.* 2011, 2012, 2015; Ranjbar & Negaresh 2014; Negaresh 2023), I realized the necessity to lectotypify two species endemic to Turkey, i.e., *K. aznavouriana* (Bornm.) Greuter & Wagenitz and *K. bornmuelleri* (Azn.) Greuter & Wagenitz, because no specific herbarium sheet was cited as holotype in their protologues and the taxa were also not lectotypified earlier (Davis & Kupicha 1975). Following a thorough examination of the literature and of the type material, lectotypes are designated here for these taxa.

LECTOTYPIFICATIONS

Family ASTERACEAE Bercht. & J.Presl
Genus *Klasea* Cass.

Klasea aznavouriana (Bornm.) Greuter & Wagenitz
(Fig. 1)

Willdenowia 33: 58 (Greuter 2003). — Basionym: *Serratula aznavouriana* Bornm., *Bulletin de l'Herbier Boissier, sér. 2*, 6: 233 (Bornmüller 1906).

LECTOTYPE. — Türkiye • Anatolia bor. Prov. Pontus, in monte Tavschan-dagh [Tavşan Dağı] prope Mersivan [Merzifon]; 1904/1905; *J.J.Manissadjian s.n.*; lectotype (**designated here**): B[B100277137]! (Fig. 1) • same data; isolecotype: B[B100277136, B100277138]!, E[E00083602]!, G[G00012010, G00012011, G00012012, G00012014]!).

REMARKS

Bornmüller (1906) indicated the type of *Serratula aznavouriana* (*Klasea aznavouriana*) as 'Anatolia bor. Prov. Pontus, in monte Tavschan-dagh prope Mersivan [Merzifon], 1904/1905, *Manissadjian*'. Pertaining to the type specification provided in the protologue, eight duplicates

specimens of *Manissadjian s.n.* were traced (B100277136, B100277137, B100277138, E00083602, G00012010, G00012011, G00012012, G00012014). According to McNeill (2014), the existence of a holotype cannot be considered before 1990 unless one particular herbarium was indicated in the protologue (and only one specimen of the gathering was deposited there) or if it was made clear that only a single specimen of the gathering existed. Because there was more than one specimen for this species housed in more than one institution, they are duplicates. Based on Art. 40.2 Note 1 (Turland *et al.* 2018) these must all be treated as syntypes. The sheet B100277137 is selected as the lectotype here because it contains leaves, flowers, and capitula that are better developed than in the other duplicates, and the characters of the specimen agree with the protologue.

Klasea bornmuelleri (Azn.) Greuter & Wagenitz
(Fig. 2)

Willdenowia 33: 58 (Greuter 2003). — Basionym: *Serratula bornmuelleri* Azn., *Repert. Spec. Nov. Regni Veg.* 11: 397 (Aznavour 1912). *Schumeria bornmuelleri* (Azn.) Iljin, *Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk S.S.S.R.* 20: 364 (Iljin 1960), *comb. inval.*

LECTOTYPE. — Türkiye • [Malatya], collines au-dessus de Dérindeh [Darende]; 14.VII.1906; *G.Post & B.Post 30*; lectotype (**designated here**): G[G00012017]! (Fig. 2) • same data; isolecotype: B[B100277131]!, G[G00012016]!).

REMARKS

Aznavour (1912) described *Serratula bornmuelleri* (*Klasea bornmuelleri*) based on collections of *G.Post & B.Post 30* from Dérindeh, but did not designate a holotype nor did mention the name of the herbarium where the specimens were housed. Three specimens of *K. bornmuelleri* collected by *G.Post & B.Post* from Dérindeh were found. Based on Art. 40.2, Note 1 (Turland *et al.* 2018), these must all be treated as syntypes. Of these, the sheet G00012017 is better preserved and more complete than the other specimens, and best represents the species description given in the protologue; hence, it is designated here as the lectotype.

Acknowledgements

I am indebted to the personnel of the following herbaria: B, E, G, and JE, for their contribution during the revision of material and for providing the images of types. I would also like to thank the Agricultural Sciences and Natural Resources University of Khuzestan (project no. 1400/19) for financial support. The referees of the article, Prof. Ergin Hamzaoglu (Gazi University, Ankara, Türkiye), Dr. Zohreh Yousefi (Ataturk Universitesi, Erzurum Türkiye), Dr. Mina Khorasani (Farhangian University of Tehran, Tehran, Iran) and Thierry Deroin (Muséum national d'Histoire naturelle, Paris) are also thanked for their remarks on a previous version of the article.



FIG. 1. — Lectotype of *Klasea aznavouriana* (Bornm.) Greuter & Wagenitz, *Manissadjian s.n.* (B100277137).




Fig. 2. — Lectotype of *Klasea bornmuelleri* (Azn.) Greuter & Wagenitz, G.Post & B.Post 30 (G00012017).

REFERENCES

- AZNAVOUR G. V. 1912. — Une plante nouvelle d'Anatolie. *Repertorium Specierum Novarum Regni Vegetabilis* 11: 397-398. <https://www.biodiversitylibrary.org/page/237564>
- BORNMÜLLER J. F. N. 1906. — Ueber eine neue *Serratula*-Art der Anatolischen Flora (*Serratula aznavouriana* Bornm.). *Bulletin de l'Herbier Boissier, sér. 2*, 6: 233-234. <https://www.biodiversitylibrary.org/page/33915251>
- DAVIS P. H. & KUPICHA F. K. 1975. — *Serratula*, in DAVIS P. H. (ed.), *Flora of Turkey and the East Aegean Islands*. Vol. 5. Edinburgh University Press, Edinburgh: 452-460.
- DOGAN B., KOYUNCU O., DURAN A. & OCAK A. 2015. — *Klasea yunus-emrei* (Asteraceae), a new species from central Anatolia Turkey. *Plant Biosystems* 149 (6): 1010-1014. <http://doi.org/10.1080/11263504.2014.969356>
- GREUTER W. 2003. — The Euro Med treatment of Cardueae (Compositae): generic concepts and required new names. *Willdenowia* 33: 49-61. <http://doi.org/10.3372/wi.33.33104>
- HELLWIG F. H. 2004. — Centaureinae (Asteraceae) in the Mediterranean – history of ecogeographical radiation. *Plant Systematics and Evolution* 246: 137-162. <https://doi.org/10.1007/s00606-004-0150-2>
- ILJIN M. M. 1960. — Shumeriya: Novyi rod slozhnotsvetnykh, *Schumeria* Iljin. Genus novum Compositarum. *Botanicheskie Materialy Gerbariia Botanicheskogo Instituta imeni V. L. Komarova Akademii Nauk SSSR* 20: 363-369.
- MCNEILL J. 2014. — Holotype specimens and type citations: general issues. *Taxon* 63 (5): 1112-1113. <https://doi.org/10.12705/635.7>
- MARTINS L. & HELLWIG F. H. 2005. — Systematic position of the genera *Serratula* and *Klasea* within Centaureinae (Cardueae, Asteraceae) inferred from ETS and ITS sequence data and new combinations in *Klasea*. *Taxon* 54: 632-638. <https://doi.org/10.2307/25065420>
- MARTINS L. 2006. — Systematics and biogeography of *Klasea* (Asteraceae-Cardueae) and a synopsis of the genus. *Botanical Journal of the Linnean Society* 152: 435-464. <https://doi.org/10.1111/j.1095-8339.2006.00583.x>
- NEGARESH K. 2023. — Lectotypification of *Klasea coriacea* (Fisch. & C.A. Mey. ex DC.) Holub and *K. leptoclada* (Bornm. & Sint.) L.Martins (Asteraceae). *Annales Botanici Fennici* 6: 105-108. <https://doi.org/10.5735/085.060.0117>
- RANJBAR M. & NEGARESH K. 2014. — On the identity of *Centaurea paradoxa* (Asteraceae, Cardueae, Centaureinae) from Iran. *Willdenowia* 44: 13-20. <https://doi.org/10.3372/wi.44.44102>
- RANJBAR M., NEGARESH K. & KARAMIAN R. 2011. — Taxonomic notes on the *Klasea calcarea* group (Asteraceae) from Iran. *Feddes Repertorium* 122: 465-471. <https://doi.org/10.1002/fedr.201100031>
- RANJBAR M., NEGARESH K., KARAMIAN R. & JOHARCHI M. R. 2012. — *Klasea nana* (Asteraceae), a new species from NE Iran. *Annales Botanici Fennici* 49: 402-406. <https://doi.org/10.5735/085.049.0615>
- RANJBAR M., NEGARESH K., KARAMIAN R. & JOHARCHI M. R. 2015. — A synopsis of *Klasea* sect. *Schumeria* (Asteraceae, Cardueae) in Iran. *Novon* 24: 186-198. <https://doi.org/10.3417/2012065>
- SUSANNA A. & GARCIA-JACAS N. 2007. — Tribe Cardueae, in KADEREIT J. W. & JEFFREY C. (eds), *The Families and Genera of Vascular Plants*. Vol. 8. Springer Verlag, Berlin, Heidelberg & New York: 123-146.
- SUSANNA A. & GARCIA-JACAS N. 2009. — Cardueae (Carduoideae), in FUNK V. A., SUSANNA A., STUESSY T. F. & BAYER R. J. (eds), *Systematics, Evolution, and Biogeography of Compositae*. International Association for Plant Taxonomy, Vienna: 293-313.
- TURLAND N. J., WIERSEMA J. H., BARRIE F. R., GREUTER W., HAWKSWORTH D. L., HERENDEN P. S., KNAPP S., KUSBER W.-H., LI D.-Z., MARHOLD K., MAY T. W., MCNEILL J., MONRO A. M., PRADO J., PRICE M. J. & SMITH G. F. (eds) 2018. — International Code of Nomenclature for Algae, Fungi, and Plants (Shenzhen Code): Adopted by the Nineteenth International Botanical Congress Shenzhen, China. *Regnum Vegetabile* 159: 1-254. <https://doi.org/10.12705/Code.2018>

Submitted on 8 April 2025;
accepted on 26 May 2025;
published on 27 January 2026.