

Typification of *Salvia smyrnaea* Boiss.  
and *Salvia eriophora* Boiss. &  
Kotschy ex Boiss. (Lamiaceae)  
described by Pierre Edmond Boissier from Türkiye

Gianmarco TAVILLA & Ferhat CELEP

DIRECTEUR DE LA PUBLICATION / *PUBLICATION DIRECTOR*: Gilles Bloch  
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Thierry Deroin

RÉDACTEURS / *EDITORS*: Porter P. Lowry II; Zachary S. Rogers; Mathieu Gardère

ÉDITEUR TECHNIQUE (SUIVI ÉDITORIAL) / *DESK EDITOR (EDITORIAL PROCESS)*: Emmanuel Côté (adanson@mnhn.fr)

ÉDITEUR TECHNIQUE (PRODUCTION) / *DESK EDITOR (PRODUCTION)*: Fanny Herman

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

F. Blasco (CNRS, Toulouse)  
M. W. Callmänder (Conservatoire et Jardin botaniques de Genève)  
J. A. Doyle (University of California, Davis)  
P. K. Endress (Institute of Systematic Botany, Zürich)  
P. Feldmann (Cirad, Montpellier)  
L. Gautier (Conservatoire et Jardin botaniques de Genève)  
F. Ghahremaninejad (Kharazmi University, Téhéran)  
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)  
A. A. Khapugin (Tyumen State University, Russia)  
J.-Y. Lesouef (Conservatoire botanique de Brest)  
P. Morat (Muséum national d'Histoire naturelle, Paris)  
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)  
S. E. Rakotoarisoa (Millenium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)  
P. H. Raven (Missouri Botanical Garden, St. Louis)  
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)  
J. G. West (Australian National Herbarium, Canberra)  
J. R. Wood (Oxford)

COUVERTURE / *COVER*:

Réalisée à partir des Figures de l'article/*Made from the Figures of the article.*

*Adansonia* est indexé dans / *Adansonia is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

*Adansonia* est distribué en version électronique par / *Adansonia is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

*Adansonia* est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris  
*Adansonia is a fast track journal published by the Museum Science Press, Paris*

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish: Geodiversitas, Zoosystema, Anthropozoologica, European Journal of Taxonomy, Naturae, Cryptogamie* sous-sections *Algologie, Bryologie, Mycologie, Comptes Rendus Palevol*

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle  
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)  
Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40  
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

Les articles publiés dans *Adansonia* sont distribués sous licence CC-BY 4.0/*Articles published in Adansonia are distributed under a CC-BY 4.0 license.*  
ISSN (imprimé / *print*): 1280-8571/ ISSN (électronique / *electronic*): 1639-4798

**Typification of *Salvia smyrnaea* Boiss.  
and *Salvia eriophora* Boiss. & Kotschy ex Boiss.  
(Lamiaceae) described by Pierre Edmond Boissier  
from Türkiye**

**Gianmarco TAVILLA**

Competence Center for Economic, Ecological and Social Sustainability,  
Free University of Bozen-Bolzano, via Cassa di Risparmio 21, Bolzano I-39100 (Italy)  
gtavilla@outlook.com (corresponding author)

**Ferhat CELEP**

Department of Biology, Faculty of Engineering and Natural Sciences,  
Kırıkkale University, Kırıkkale (Türkiye)  
ferhat\_celep@hotmail.com

Submitted on 1 May 2025 | accepted on 5 June 2025 | published on 23 February 2026

Tavilla G. & Celep F. 2026. — Typification of *Salvia smyrnaea* Boiss. and *Salvia eriophora* Boiss. & Kotschy ex Boiss. (Lamiaceae) described by Pierre Edmond Boissier from Türkiye *Adansonia*, sér. 3, 48 (5): 29-34. <https://doi.org/10.5252/adansonia2026v48a5>. <http://adansonia.com/48/5>

**KEY WORDS**

Endemic species,  
second-step  
lectotypification,  
Türkiye,  
Ian Hedge.

**ABSTRACT**

A contribution to the typification of two endemic species of *Salvia* from Türkiye is presented. The names of *S. smyrnaea* Boiss. and *S. eriophora* Boiss. & Kotschy ex Boiss. are inadvertently lectotypified, and therefore, require second-step lectotypification. The designated lectotypes of *S. smyrnaea* and *S. eriophora* are housed at the Geneva herbarium from the Boissier collection.

**RÉSUMÉ**

*Typification de Salvia smyrnaea et Salvia eriophora (Lamiaceae) décrites par Pierre Edmond Boissier de Turquie.* Une contribution à la typification de deux espèces endémiques de *Salvia* de Türkiye est présentée. Les noms de *S. smyrnaea* Boiss. et *S. eriophora* Boiss. & Kotschy ex Boiss. ont été inopinément lectotypifiés et nécessitent donc une nouvelle lectotypification. Les lectotypes désignés pour *S. smyrnaea* et *S. eriophora* sont conservés à l'herbier de Genève dans la collection Boissier.

**MOTS CLÉS**

Espèces endémiques,  
second-step  
lectotypification,  
Türkiye,  
Ian Hedge.

## INTRODUCTION

The genus *Salvia* L. is the largest one in the family Lamiaceae having nearly 1000 species that are really diverse and worldwide distributed (Walker & Sytsma 2007; Rose *et al.* 2021; Celep & Doğan 2023; POWO 2025). The genus has its main center of species diversity in Central and South America, southwestern Asia and the Mediterranean region, and Eastern Asia (Walker & Sytsma 2007). Etlinger (1777) published the first botanical monograph on *Salvia* and recognized 49 species. The first infrageneric classification of the genus was made by Benthams (1833), based on the morphology of the calyx, corolla and stamen, and was later revised by Briquet (1897). Since then, many scientists have contributed to the evolution of these classifications (Pobedimova 1954; Wu 1977; Murata & Yamazaki 1993). On the other hand, some researchers have avoided subgeneric and sectional classifications by describing “species groups” or small sections (Epling 1938-1939; Hedge 1974; 1982). Currently, molecular phylogenetic studies have shown that *Salvia* is not monophyletic and five other genera, namely *Dorystaechas* Boiss. & Heldr. ex Benth., *Meriandra* Benth., *Perovskia* Kar., *Rosmarinus* L., and *Zhumeria* Rech.f. & Wendelbo, are embedded within it (Walker & Sytsma 2007; Walker *et al.* 2015; Drew *et al.* 2017; Fragoso-Martínez *et al.* 2018; Kriebel *et al.* 2019).

As part of ongoing studies on the taxonomy of the genus *Salvia* (Tavilla 2025), the species *S. smyrnaea* and *S. eriophora* are revised from a nomenclatural point of view. These two species are listed in the *Flora of Turkey and East Aegean Islands* (Hedge 1982), and Hedge indicated a holotype for each taxon. This statement should be interpreted as inadvertent lectotypification (Prado *et al.* 2015). Therefore, a second-step lectotypification for these two names is required in order to designate a single lectotype (Art. 9.17; Turland *et al.* 2018, hereafter ICN).

## TAXONOMY

Family LAMIACEAE Martinov  
Genus *Salvia* L.

*Salvia smyrnaea* Boiss.

*Diagnoses Plantarum Orientalium Novarum*, ser. 1, 5: 5 (Boissier 1844).

TYPE MATERIAL. — **Türkiye** • B1 Izmir, in cacumine montis Tartali (Tahtali Da.) ad orientem Smyrneam (Izmir); V.1842; *Boissier* (G); first-step lectotype designated by Hedge (1982: 429) • Cacumine montis ad orientem Smyrnae; V. 1842; *Boissier*; second-step lectotype (**here designated**): G-BOIS [G00156010]! (Fig. 1) • same data; isolectotypes: G[G00085141, G00085166, G00435332]!, GH[GH0001884]!, JE[JE00002705]!, K[K000509016, K000509017]!, LE[LE00016783]!, MW[MW0594584]!, NY[NY00000353]!, P[P00714794, P00714795, P00714797, P00714798]!, US[US00517381]! • same data; syntypes: HBG[HBG518237]!, M[M0099274]!

## DISTRIBUTION AND ECOLOGY

*Salvia smyrnaea* is a narrow endemic species found in the Kemalpaşa District of İzmir and the Söke District of Aydın provinces

in western Türkiye (Hedge 1982; Celep 2010; POWO 2025). It thrives on calcareous rocks in open *Pinus* sp. forests, around the summit of Nif mountain, in gravelly places at an altitude of 900-1510 m (Celep 2010; Celep *et al.* 2011). According to Celep & Doğan (2023), the conservation status of *S. smyrnaea* is assigned as Endangered (EN) based on IUCN (2022) criteria.

## NOTES

*Salvia smyrnaea* was validly described by Boissier (1844) in the *Diagnoses Plantarum Orientalium novarum*. In the prologue, Boissier indicated “Tartali ad orientem Smyrnae” as the habitat where the species was collected. Today, this area corresponds to Nif Mountain between the Torbalı and Kemalpaşa districts of İzmir. Based on herbarium searches, we found four herbarium sheets (G00156010, G00085141, G00085166, G00435332) in Geneva herbarium, which is typically the herbarium where Boissier type material is stored (Stafleu & Cowan 1976). Among them, one sheet (G00156010) belongs to the G-BOIS collection. Moreover, several duplicates were found in other herbaria, such as GH, HBG, JE, K, LE, M, MW, NY, P and US (herbarium acronyms follow Thiers 2025). The specimens housed in G that are labelled with a “Typus” red label typically indicate that the material has not been published as specifically designated. While other specimens in different herbaria are recognized as isotypes, these herbarium sheets often bear a yellow abbreviated label “Herb. E. Boissier, *Salvia smyrnaea* Boiss., Mons Tartali supra Smyrnam, Maio 1842”.

Hedge (1982) carried out an inadvertent lectotypification due to the statement “type” which can be considered a valid type designation and should be corrected under lectotype according to Art. 7.11 of ICN. However, there are several specimens in G which is the herbarium reported by Hedge (1982). Therefore, a second-step lectotypification is needed to clarify the name of *S. smyrnaea* and designate a single lectotype (Art. 9.17). According to the label on the G00085141 specimen, Hedge revised this one, which is part of the syntypes but is not found in Boissier’s herbarium. Therefore, we prefer to designate the G-BOIS specimen as a lectotype, which represents a well-preserved specimen of the species that displays all the diagnostic morphological features needed for the identification of the species, such as leaves mostly basal, ovate-oblong to oblong, flowering stems glandular below, calyx upper lip clearly 3-dentate, corolla with subequal lips. Therefore, it corresponds to the current application of the name (Hedge 1982).

*Salvia eriophora* Boiss. & Kotschy ex Boiss.

*Flora Orientalis* 4 (2): 611 (Boissier 1879).

TYPE MATERIAL. — **Türkiye** • C5 Adana/Nigde, ab Jool Baatch (Yol Bahçe) ad radices montis Allahdagh (Ala Da.) Ciliciae; 1525 m alt.: 3.VI.1859; *Kotschy* 229; G; first-step lectotype designated by Hedge (1982: 437) • ab Jool Baatch ad radices montis Allahdagh Ciliciae; 5000'; 3.VI.1859; *Kotschy* 229; second-step lectotype (**here designated**): G-BOIS [G00156025]!, (Fig. 2)



FIG. 1. — Lectotype of *Salvia smyrnaea* Boiss. (G00156010) © Conservatoire & Jardin botaniques de la Ville de Genève.




FIG. 2. — Lectotype of *Salvia eriophora* Boiss. & Kotschy ex Boiss. (G00156025) © Conservatoire & Jardin botaniques de la Ville de Genève.

• same data; isolectotypes: B[B100264866]!, BM[BM000950400]!, G[G00085150, G00446594]!, JE[JE00001470]!, K[K000929731]!, L[L.2793331]!, MW[MW0594577]!, P[P00714732, P00714733, P00714734]!, S[S08-10892]!

#### DISTRIBUTION AND ECOLOGY

The native range of this very rare endemic species is central and southern Türkiye, growing on rocky limestone slopes and high mountain steppe at an altitude of 1525-1880 m (Hedge 1982; POWO 2025). However, at present, this species could not be collected from the *locus classicus* (C5 Adana/Niğde, central southern Türkiye) due to uncertainty regarding the locality. Later, it was found at the second locality by P. H. Davis (Hedge 1982) that this species is naturally distributed only in the high mountain steppe in the pass (at an altitude of 1850-1880 m) between Pınarbaşı and Gürün in Kayseri/Sivas, with only a limited number of individuals. In this area, individuals have to survive under the pressure of overgrazing and road construction activities (Kahraman *et al.* 2012). According to IUCN (2022) criteria, this species can be classified as Critically Endangered (Kahraman *et al.* 2012; Celep & Doğan 2023).

#### NOTES

Boissier (1879) described this species in *Flora Orientalis*. He determined the material and described several new species to science, frequently relying solely on Kotschy's specimens (Lack 2020). Boissier indicated “Hab. ab Jool Baatch ad radices montis Allahdagh Ciliciae 5000' (Ky. 229!)”, which means he particularly observed the specimen numbered 229 from the Kotschy collection, and it can be considered original material. However, Hedge (1982) used the term holotype (“holo. G”) that must be corrected to lectotype. Due to the occurrence of three herbarium sheets with the same labels in G (G00085150, G00156025, G00446594), a second-step lectotype is required to designate a single specimen. Additionally, several duplicates of the original material were found in the B, BM, JE, K, L, MW, P, and S herbaria. The specimens in the G herbarium bear the typical red label with the word “Typus”, while the other syntypes, deposited in the herbaria BM, P, and S, refer to the type collection. Only the specimen deposited in the MW herbarium bears a modern label with the word “isolectotypus” but does not refer to any bibliographic source where a lectotypification could have been performed. Therefore, to the best of our knowledge, this name remains inadvertently typified according to the ICN articles. The sheet G00156025 from the Boissier collection is chosen here as second-step lectotype. It includes a label with the number 229; therefore, it is in accordance with the protologue reference. Furthermore, the selected specimen shows the main morphological features of *S. eriophora*: stems densely arachnoid-lanate above; leaves linear-oblong, calyx tubular-campanulate, teeth long-subulate, corolla lobes not widely diverging, tube straight, upper lip straight. Thus, it confirms the current application of the name (Hedge 1982).

#### Acknowledgements

We appreciate the reviewers Thierry Deroin (MNHN) and the anonymous referee for their valuable suggestions and comments.

#### REFERENCES

- BENTHAM G. 1833. — *Labiatarum Genera et Species, Vol. 3*. James Ridgway and Sons, London, 783 p. <https://www.biodiversitylibrary.org/item/144580> [accessed 17 February 2026].
- BOISSIER E. 1844. — *Diagnoses plantarum orientalium novarum ser. 1*, 5. Leipzig, Paris, 367 p. <https://www.biodiversitylibrary.org/item/142678> [accessed 17 February 2026].
- BOISSIER E. 1879. — *Flora Orientalis: sive, Enumeratio plantarum in Oriente a Graecia et Aegypto ad Indiae fines hucusque observatarum*, 4. Apud H. Georg, Geneva, 1276 p. <https://doi.org/10.5962/bhl.title.20323>
- BRIQUET J. 1897. — *Salvia*, in ENGLER A. & PRANTL K. (eds), *Die natürlichen Pflanzenfamilien nebst ihrer Gattungen und wichtigeren Arten, vol. IV*. Wilhelm Engelmann, Leipzig: 183-287.
- CELEP F. 2010. — Revision of the genus *Salvia* L. (Labiatae) in the Mediterranean and the Aegean geographic regions of Turkey. PhD dissertation, Middle East Technical University, Ankara, 287 p. <https://etd.lib.metu.edu.tr/upload/3/12612069/index.pdf> [accessed 12 February 2026].
- CELEP F. & DOĞAN M. 2023. — The Genus *Salvia* in Turkey: Morphology, Ecology, Phytogeography, Endemism and Threat Categories, in MATHÉ A. & TURGUT K. (eds), *Medicinal and Aromatic Plants of Turkey. Medicinal and Aromatic Plants of the World, vol. 10*. Springer, Cham: 107-120. [https://doi.org/10.1007/978-3-031-43312-2\\_5](https://doi.org/10.1007/978-3-031-43312-2_5)
- CELEP F., KAHRAMAN A. & DOĞAN M. 2011. — A new taxon of the genus *Salvia* (Lamiaceae) from Turkey. *Plant Ecology and Evolution* 144 (1): 111-114. <https://doi.org/10.5091/plecevo.2011.423>
- DREW B. T., GONZÁLEZ-GALLEGOS J. G., XIANG C. L., KRIEBEL R., DRUMMOND C. P., WALKED J. B. & SYTSMAN K. J. 2017. — *Salvia united: the greatest good for the greatest number*. *Taxon* 66: 133-145. <https://doi.org/10.12705/661.7>
- EPLING C. 1938-1939. — A revision of *Salvia*, subgenus *Calosphace*. *Feddes Repertorium Beihefte* 110: 1-383.
- ETLINGER A. E. 1777. — *De Salvia: Dissertatio inauguralis. Commentatio Botanico-Medica de Salvia*. Wolfgangi Waltheri, Erlangae, 112 p.
- FRAGOSO-MARTÍNEZ I., MARTÍNEZ-GORDILLO M., SALAZAR G. A., SAZATORNIL F., JENKS A. A., GARCÍA PEÑA M. R., BARRERA-AVELEIDA G., BENITEZ-VIEYRA S., MAGALLÓN S., CORNEJO-TENORIO G. & GRANADOS MENDOZA C. 2018. — Phylogeny of the Neotropical sages (*Salvia* subg. *Calosphace*; Lamiaceae) and insights into pollinator and area shifts. *Plant Systematics and Evolution* 304: 43-55. <https://doi.org/10.1007/s00606-017-1445-4>
- HEDGE I. C. 1974. — A Revision of *Salvia* in Africa including Madagascar and the Canary Islands. *Notes from the Royal Botanic Garden Edinburgh* 33: 1-121.
- HEDGE I. C. 1982. — *Salvia* L., in DAVIS P.-H. (ed.), *Flora of Turkey and East Aegean Islands, vol. 7*. University Press, Edinburgh: 400-461.
- IUCN 2022. — The IUCN Red List of Threatened Species vers. 2020-1. <http://www.iucnredlist.org> [accessed 28 April 2025].
- KAHRAMAN A., BAGHERPOUR S., KARABACAK E., DOĞAN M., DOĞAN H. M., UYSAL I. & CELEP F. 2012. — Reassessment of conservation status of the genus *Salvia* (Lamiaceae) in Turkey II. *Turkish Journal of Botany* 36 (2): 103-124. <https://doi.org/10.3906/bot-1007-2>
- KRIEBEL R., DREW B. T., DRUMMOND C. P., GONZÁLEZ-GALLEGOS J. G., CELEP F., MAHDJOUR M. M., ROSE J. P., XIANG C.-L., HU G.-X., WALKER J. B., LEMMON E. M., LEMMON A. R. & SYTSMAN K. J. 2019. — Tracking temporal shifts in area, biomes, and pollinators in the radiation of *Salvia* (sages) across continents: leveraging anchored hybrid enrichment and targeted sequence data. *American Journal of Botany* 106: 573-597. <https://doi.org/10.1002/ajb2.1268>

- LACK H. W. 2020. — Theodor Kotschy in Iran, 1841-1843. Botanical collections and an early printed vegetation profile. *Candollea* 75: 31-43. <https://doi.org/10.15553/c2020v751a3>
- MURATA G. & YAMAZAKI T. 1993. — *Salvia*, in IWATSUKI K., YAMAZAKI T., BOUFFORD D. E. & OHBA H. (eds), *Flora of Japan, vol. IIIa*. Kodansha, Tokyo: 302-307.
- POBEDIMOVA E. G. 1954. — *Salvia*, in SHISHKIN B. K. (ed.), *Flora of the U.S.S.R., vol. 21*. Izdatel'stvo Akademii Nauk SSR, Moscow: 154-255.
- PRADO J., HIRAI R. Y. & MORAN R. C. 2015. — Proposals concerning inadvertent lectotypifications (and neotypifications). *Taxon* 64: 651. <https://doi.org/10.12705/643.29>
- POWO 2025. — *Plants of the World Online*. The Royal Botanic Gardens, Kew. <https://powo.science.kew.org/> [accessed 4 April 2025].
- ROSE J. P., KRIEBEL R., KAHAN L., DI NICOLA A., GONZÁLEZ-GALLEGOS J. G., CELEP F., LEMMON E. M., LEMMON A. R., SYTSMA K. J. & DREW B. T. 2021. — Sage insights into the phylogeny of *Salvia*: Dealing with sources of discordance within and across genomes. *Frontiers in Plant Science* 12: 2606. <https://doi.org/10.3389/fpls.2021.767478>
- STAFLEU, F. A. & COWAN R. S. 1976. — *Taxonomic literature*. Ed. 2. Vol. 1. Bohn, Scheltema & Holkema, Utrecht, 1136 p. <http://dx.doi.org/10.5962/bhl.title.48631>
- TAVILLA G. 2025. — Typification of two names in the genus *Salvia* (*Lamiaceae*). *Kew Bulletin* 80: 209-213. <https://doi.org/10.1007/s12225-024-10241-0>
- THIERS B. 2025. — Index Herbariorum: a Global Directory of Public Herbaria and Associated Staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [accessed 4 April 2025]
- TURLAND N. J., WIERSEMA J. H., BARRIE F. R., GREUTER W., HAWKSWORTH D. L., HERENDEEN P. S., KNAPP S., KUSBER W.-H., LI D.-Z., MARHOLD K., MAY T. W., MCNEILL J., MONRO A. M., PRADO J., PRICE M. J. & SMITH G. F. (eds). 2018. — International Code of Nomenclature for algae, fungi and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Vegetabile* 1-254. Glashütten: Koeltz Botanical Books. <https://doi.org/10.12705/Code.2018>
- WALKER J. B. & SYTSMA K. J. 2007. — Staminal evolution in the genus *Salvia* (*Lamiaceae*): Molecular phylogenetic evidence for multiple origins of the staminal lever. *Annals of Botany* 100 (2): 375-391. <https://doi.org/10.1093/aob/mcl176>
- WALKER J. B., DREW B. T. & SYTSMA K. J. 2015. — Unravelling species relationships and diversification within the iconic California Floristic Province sages (*Salvia* subgenus *Audibertia*, *Lamiaceae*). *Systematic Botany* 40: 826-844. <https://doi.org/10.1600/036364415X689285>
- WU C. Y. 1977. — *Salvia*, in WU C. Y. (ed.), *Flora Reipublicae Popularis Sinicae, vol. 66*. Science Press, Beijing: 70-196.

Submitted on 1 May 2025;  
accepted on 5 June 2025;  
published on 23 February 2026.