

cryptogamie

Algologie

2020 • 41 • 3

Pseudoderbesia eckloniae, sp. nov.
(Bryopsidaceae, Ulvophyceae)
from Western Australia

John M. HUISMAN & Heroen VERBRUGGEN

DIRECTEUR DE LA PUBLICATION : Bruno DAVID,
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / EDITOR-IN-CHIEF : Line LE GALL

ASSISTANT DE RÉDACTION / ASSISTANT EDITOR : Audrina NEVEU (algo@cryptogamie.com)

MISE EN PAGE / PAGE LAYOUT : Audrina NEVEU

RÉDACTEURS ASSOCIÉS / ASSOCIATE EDITORS

Ecoevolutionary dynamics of algae in a changing world

Stacy KRUEGER-HADFIELD

Department of Biology, University of Alabama, 1300 University Blvd, Birmingham, AL 35294 (United States)

Jana KULICHOVA

Department of Botany, Charles University, Prague (Czech Republic)

Cecilia TOTTI

Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona (Italy)

Phylogenetic systematics, species delimitation & genetics of speciation

Sylvain FAUGERON

UMI3614 Evolutionary Biology and Ecology of Algae, Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Av. Bernardo O'Higgins 340, Santiago (Chile)

Marie-Laure GUILLEMIN

Instituto de Ciencias Ambientales y Evolutivas, Universidad Austral de Chile, Valdivia (Chile)

Diana SARNO

Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli (Italy)

Comparative evolutionary genomics of algae

Nicolas BLOUIN

Department of Molecular Biology, University of Wyoming, Dept. 3944, 1000 E University Ave, Laramie, WY 82071 (United States)

Heroen VERBRUGGEN

School of BioSciences, University of Melbourne, Victoria, 3010 (Australia)

Algal physiology & photosynthesis

Janet KÜBLER

California State University Northridge, Department of Biology, California State University, Northridge, CA 91330-8303 (United States)

Prokaryotic algae

Nico SALMASO

IASMA Research and Innovation Centre, Fondazione Mach-Istituto Agrario di S. Michele all'Adige, Limnology and River Ecology, Via E. Mach, 1, 38010 San Michele all'Adige, Trento (Italy)

Vitor VASCONCELOS

Faculdade de Ciências da Universidade do Porto and CIIMAR, Rua do Campo Alegre, s/n, 4169-007 Porto (Portugal)

COUVERTURE / COVER:

Extraits d'éléments de la Figure 1A / Extracts of the Figure 1A

Cryptogamie, Algologie est indexé dans / *Cryptogamie, Algologie* is indexed in:

- Aquatic Sciences & Fisheries Abstracts Part I.
- Biological Abstracts
- Chemical Abstracts
- Current Contents
- Marine Science Contents Tables (FAO)
- Science Citation Index
- Publications bibliographiques du CNRS (Pascal).

Cryptogamie, Algologie est distribué en version électronique par / *Cryptogamie, Algologie* is distributed electronically by:

- BioOne® (<http://www.bioone.org/loi/crya>)

Cryptogamie, Algologie est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Cryptogamie, Algologie is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / The Museum Science Press also publishes:

Adansonia, Geodiversitas, Zoosystema, Anthropozoologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Bryologie, Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2020

ISSN (imprimé / print): 0181-1568 / ISSN (électronique / electronic): 1776-0984

***Pseudoderbesia eckloniae*, sp. nov.**
(Bryopsidaceae, Ulvophyceae)
from Western Australia

John M. HUISMAN

Western Australian Herbarium, Biodiversity and Conservation Science,
Department of Biodiversity, Conservation and Attractions, Locked Bag
104, Bentley Delivery Centre, Western Australia 6983 (Australia)
john.huisman@dbca.wa.gov.au (corresponding author)

Heroen VERBRUGGEN

School of BioSciences, University of Melbourne,
Victoria 3010 (Australia)

Submitted on 10 September 2019 | Accepted on 20 November 2019 | Published on 4 March 2020

Huisman J. M. & Verbruggen H. 2020. — *Pseudoderbesia eckloniae*, sp. nov. (Bryopsidaceae, Ulvophyceae) from Western Australia. *Cryptogamie, Algologie* 41 (3): 19–23. <https://doi.org/10.5252/cryptogamie-algologie2020v41a3>. <http://cryptogamie.com/algologie/41/3>

ABSTRACT

Pseudoderbesia eckloniae Huisman & Verbruggen, sp. nov., is described for specimens collected from Cape Peron, Western Australia, growing epiphytically on the kelp *Ecklonia radiata* (C.Agardh) J.Agardh. The new species forms a velvety turf on the host and is distinguished from related species by a combination of molecular analyses (*rbcL*) and morphological characteristics.

RÉSUMÉ

Pseudoderbesia eckloniae, sp. nov. (Bryopsidaceae, Ulvophyceae) d'Australie occidentale.

Pseudoderbesia eckloniae Huisman & Verbruggen, sp. nov., est décrit pour des spécimens collectés au cap Péron, en Australie occidentale, poussant de manière épiphytique sur le kelp *Ecklonia radiata* (C.Agardh) J.Agardh. La nouvelle espèce forme un gazon dense sur l'hôte et se distingue des espèces apparentées par une combinaison d'analyses moléculaires (*rbcL*) et de caractéristiques morphologiques.

KEY WORDS

Australia,
Bryopsidaceae,
new species.

MOTS CLÉS

Australie,
Bryopsidaceae,
espèce nouvelle.

INTRODUCTION

The genus *Pseudoderbesia* Calderon & Schnetter (1991) was described for small, siphonous green algae with rhizoidally attached stoloniferous bases that give rise to erect axes with dichotomously branched, tapering siphons. Reproduction is via biflagellate zooids that arise in transformed branches and are released through papillae in the siphon wall. The genus is currently monospecific and was poorly known until the review of Leliaert *et al.* (2014), who described specimens from Greece that conformed to *Pseudoderbesia* but could not be confidently assigned to its only species, *P. arbuscula* Calderon & Schnetter (1991), and were left as *Pseudoderbesia* sp. (Greece).

A recent Western Australian collection of a tufted green alga, growing epiphytically on the kelp *Ecklonia radiata* (C.Agardh) J.Agardh, morphologically agreed with *Pseudoderbesia* but differed in subtle ways from *P. arbuscula* and other taxa described in Leliaert *et al.* (2014), and is described here as a new species.

SYSTEMATICS

Family BRYOPSIDACEAE Bory
Genus *Pseudoderbesia* Calderon & Schnetter

Pseudoderbesia eckloniae
Huisman & Verbruggen, sp. nov.
(Fig. 1)

HOLOTYPE. — PERTH 09134638. Collected at Cape Peron, Western Australia (32°16.18'S, 115°41.178'E), epiphytic on *Ecklonia radiata* at 2 m depth, 9.I.2019, J.M.Huisman 9.I.19.1A-C (three specimens mounted on sheet) (Fig. 1B).

PARATYPE. — PERTH 09134646. *Loc. id.*, 30.XII.2018, J.M.Huisman 30.12.18.

ETYMOLOGY. — The epithet “eckloniae” refers to the species’ only known host, *Ecklonia radiata*.

DESCRIPTION

Thallus grass green, forming a velvety layer on the surface of degrading branches of *Ecklonia radiata* (Fig. 1A), spreading laterally for up to 10 cm, siphonous, with dichotomously branched upright axes arising from a stoloniferous base (Fig. 1C). Stolons terete, 35–50 µm in diameter, attached by elongate, terete rhizoidal siphons. Upright axes dichotomously (rarely trichotomously) branched 4–7 times when mature, dichotomies mostly at similar levels but occasionally unequal, with slight constrictions commonly present at branch bases (Fig. 1D). Upright axes proximally 50 µm in diameter when young, 120–200 µm in diameter when mature; distal branches markedly narrowing, with ultimate branches 12–25 µm in diameter. Chloroplasts numerous, elongate to ovate, each with a single pyrenoid (Fig. 1E). Reproduction not observed.

MOLECULAR ANALYSIS

The *rbcL* gene of four samples of the new species was sequenced following the procedures of Verbruggen *et al.* (2009) and submitted to GenBank (MT108929). The four sequences of *Pseudoderbesia eckloniae*, sp. nov., were identical and only one was included for further analysis. The sequence was aligned with those of Bryopsidineae used in Leliaert *et al.* (2014) and *Lambia antarctica* (Skottsberg) Delépine (Cremen *et al.* 2019). A maximum likelihood phylogenetic tree was inferred with RAxML 8.2.10, using a GTR+G+I model and 100 standard bootstraps (Stamatakis 2014). The phylogenetic tree (Fig. 2) shows that the genus *Pseudoderbesia* is related to *Lambia* Delépine and *Bryopsis* J.V.Lamouroux, but the exact relationships among these genera is not recovered with sufficient bootstrap support. The divergence between *Pseudoderbesia eckloniae*, sp. nov., and *Pseudoderbesia* sp. from Greece is clear in the tree, and the *rbcL* interspecific divergence between these two *Pseudoderbesia* species is 3.87%.

AFFINITIES

The only currently named species of *Pseudoderbesia* is the generitype, *P. arbuscula* Calderon & Schnetter (1991). Based on the comparative table provided in Leliaert *et al.* (2014), upright axes in *P. eckloniae*, sp. nov., grow to approximately twice the dimensions of *P. arbuscula*, to 8 mm tall (as opposed to 4.3 mm), with lower siphons to 200 µm in diameter (54–84 µm). In addition, rhizoids in *P. arbuscula* are short, distally lobed and surrounded by mucilage, differing from the elongate rhizoids without mucilage of *P. eckloniae*, sp. nov. Two other undescribed entities (potentially species) were described from the Canary Islands and Greece (Calderon & Schnetter 1991; Leliaert *et al.* 2014). In both, the maximum siphon diameter near the base is smaller than that of *P. eckloniae*, sp. nov. The Greek entity also differs in having short rhizoids. While the morphological variation described above could be encompassed in a single, broadly defined species, our molecular analysis indicates species-level variation between *P. eckloniae*, sp. nov., and *Pseudoderbesia* sp. (Greece), the only other taxon currently represented in molecular libraries. Based on this, plus the geographical separation between the Australian and Columbian localities, we prefer to describe our material as a new species.

Leliaert *et al.* (2014) also noted three species currently placed in *Derbesia* Solier that, based on morphology, might be better placed in *Pseudoderbesia*. These species are *D. attenuata* Dawson (1954), *D. padinae* Trono (1972) and *D. fastigiata* Taylor (1928). The dichotomously branched upright axes of *P. eckloniae*, sp. nov., are somewhat similar to those of *D. attenuata*, described originally from Vietnam (Dawson 1954) but also recorded from north-western Australia (Huisman 2015) and elsewhere. As the original material of *D. attenuata* lacked reproductive structures, Dawson expressed some doubt regarding its generic placement, and the subsequent Australian collection added no new information. In contrast to *P. eckloniae*, sp. nov., Dawson’s species was described as “usually solitary” with unconstricted upright siphons. Moreover, siphon diameters of *D. attenuata*




FIG. 1. — *Pseudoderbesia eckloniae* Huisman & Verbruggen, sp. nov.: **A**, specimen from the type collection growing on *Ecklonia radiata* (C.Agardh) J.Agardh at Cape Peron, Western Australia; **B**, one of three specimens from the holotype sheet (PERTH 09134638); **C**, closer view, showing stolons (arrow) and dichotomously branched upright axes; **D**, detail of upper branches, with slight constrictions at dichotomies. A trichotomy is also present (arrow); **E**, siphon detail showing elongate chloroplasts, each with a single pyrenoid. Scale bars: A, B, 1 cm; C, 500 µm; D, 100 µm; E, 25 µm.




Fig. 2. — Maximum likelihood phylogenetic tree of the *rbcL* gene showing the new species *Pseudoderbesia eckloniae* Huisman & Verbruggen, sp. nov., clustering with the Greek *Pseudoderbesia* strain. The scale bar is in estimated substitutions per site. Branch labels indicate bootstrap support, shown only when >70.

are markedly less than those of *P. eckloniae*, sp. nov., as is also true for *D. padinae*, which grades from lower axes of 50–60 µm diameter to ultimate branches of 6–8 µm diameter. As described by Littler & Littler (2000), the tropical American *D. fastigiata* can grow to a larger size (to 2 cm), but again the siphons do not reach the dimensions of *P. eckloniae*, sp. nov. A further distinguishing feature is the siphons of *D. fastigiata* often being swollen below dichotomies (Littler & Littler 2000).

Acknowledgements

Heroen Verbruggen thanks Marisa Pasella for molecular lab support. Funding was provided by The Australian Biological Resources Study (RFL213-08) and the University of Melbourne Botany Foundation. John M. Huisman thanks the staff of the Western Australian Herbarium. The authors also thank the associate editor, Stacy Krueger, as well as Frederik Leliaert and an anonymous reviewer for their helpful comments.

REFERENCES

- CALDERON E. & SCHNETTER R. 1991. — *Pseudoderbesia*, nuevo género de algas marinas (Bryopsidaceae, Chlorophyta). *Caldasia* 16: 285–294.
- CREMEN M. C. M., LELIAERT F., WEST J., LAM D. W., SHIMADA S., LOPEZ-BAUTISTA J. M. & VERBRUGGEN H. 2019. — Reassessment of the classification of Bryopsidales (Chlorophyta) based on chloroplast phylogenomic analyses. *Molecular Phylogenetics and Evolution* 130: 397–405. <https://doi.org/10.1016/j.ympev.2018.09.009>
- DAWSON E. Y. 1954. — Marine Plants in the Vicinity of the Institute Océanographique de Nha Trang, Viêt Nam. *Pacific Science* 8: 372–469. <http://hdl.handle.net/10125/9121>
- HUISMAN J. M. 2015. — *Algae of Australia: Marine Benthic Algae of North-western Australia. 1. Green and Brown Algae*. ABRIS & CSIRO Publishing, Canberra & Melbourne, [i]–viii, 320 p.
- LELIAERT F., VERBRUGGEN H., D'HONDT S., LOPEZ-BAUTISTA J. M. & DE CLERCK O. 2014. — The forgotten genus *Pseudoderbesia* (Bryopsidales, Chlorophyta). *Cryptogamie, Algologie* 35: 207–219. <https://doi.org/10.7872/crya.v35.iss3.2014.207>
- LITTLER D. S. & LITTLER M. M. 2000. — *Caribbean Reef Plants. An Identification Guide to the Reef Plants of the Caribbean*.

- Bahamas, Florida and Gulf of Mexico*. Offshore Graphics, Washington, 542 p.
- STAMATAKIS A. 2014. — RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312-1313. <https://doi.org/10.1093/bioinformatics/btu033>
- TAYLOR W. R. 1928. — The marine algae of Florida with special reference to the Dry Tortugas. *Publications of the Carnegie Institution of Washington* 379: [i-v], [1]-219.
- TRONO G. C. JR. 1972 [1971]. — Some New Species of Marine Benthic Algae from the Caroline Islands, Western-Central Pacific. *Micronesica* 7: 45-77.
- VERBRUGGEN H., ASHWORTH M., LODUCA S. T., VLAEMINCK C., COCQUYT E., SAUVAGE T., ZECHMAN F. W., LITTLER D. S., LITTLER M. M., LELIAERT F. & DE CLERCK O. 2009. — A multi-locus time-calibrated phylogeny of the siphonous green algae. *Molecular Phylogenetics and Evolution* 50: 642-653. <https://doi.org/10.1016/j.ympev.2008.12.018>

*Submitted on 10 September 2019;
accepted on 20 November 2019;
published on 4 March 2020.*