

cryptogamie

Bryologie

2020 • 41 • 6

The two faces of Mt Gede, Java –
Species richness, composition and
zonation of epiphytic bryophytes

Eka Aditya Putri ISKANDAR
Michael STECH & Sylvia M. De OLIVEIRA

DIRECTEUR DE LA PUBLICATION / *PUBLICATION DIRECTOR*: Bruno David,
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Denis LAMY

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS*: Marianne SALAÜN (bryo@cryptogamie.com)

MISE EN PAGE / *PAGE LAYOUT*: Marianne SALAÜN

RÉDACTEURS ASSOCIÉS / *ASSOCIATE EDITORS*

Biologie moléculaire et phylogénie / *Molecular biology and phylogeny*

Bernard GOFFINET

Department of Ecology and Evolutionary Biology, University of Connecticut (United States)

Mousses d'Europe / *European mosses*

Isabel DRAPER

Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid (Spain)

Francisco LARA GARCÍA

Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid (Spain)

Mousses d'Afrique et d'Antarctique / *African and Antarctic mosses*

Rysiek OCHYRA

Laboratory of Bryology, Institute of Botany, Polish Academy of Sciences, Krakow (Pologne)

Bryophytes d'Asie / *Asian bryophytes*

Rui-Liang ZHU

School of Life Science, East China Normal University, Shanghai (China)

Bioindication / *Biomonitoring*

Franck-Olivier DENAYER

Faculté des Sciences Pharmaceutiques et Biologiques de Lille, Laboratoire de Botanique et de Cryptogamie, Lille (France)

Écologie des bryophytes / *Ecology of bryophyte*

Nagore GARCÍA MEDINA

Department of Biology (Botany), and Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid (Spain)

COUVERTURE / *COVER*:

Extraits d'éléments de la Figure 1 / Extracts of the Figure 1

Cryptogamie, Bryologie est indexé dans / *Cryptogamie, Bryologie is indexed in*:

- Biological Abstracts
- Current Contents
- Science Citation Index
- Publications bibliographiques du CNRS (Pascal).

Cryptogamie, Bryologie est distribué en version électronique par / *Cryptogamie, Bryologie is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Cryptogamie, Bryologie est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris

Cryptogamie, Bryologie is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:

Adansonia, Geodiversitas, Zoosystema, Anthropolzoologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Algologie, Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2020

ISSN (imprimé / *print*): 1290-0796 / ISSN (électronique / *electronic*): 1776-0992

The two faces of Mt Gede, Java – Species richness, composition and zonation of epiphytic bryophytes

Eka Aditya Putri ISKANDAR

Naturalis Biodiversity Center, PO Box 9517, 2300 RA Leiden
and Leiden University, Leiden (Netherlands)
and Cibodas Botanic Garden – Indonesian Institute of Sciences (LIPI),
Jl. Kebun Raya Cibodas, Cipanas, Cianjur 43253 (Indonesia)
eka.iskandar@naturalis.nl (corresponding author)

Michael STECH

Sylvia M. De OLIVEIRA

Naturalis Biodiversity Center, PO Box 9517, 2300 RA Leiden
and Leiden University, Leiden (Netherlands)

Submitted on 13 June 2019 | Accepted on 27 January 2020 | Published on 6 May 2020

Iskandar E. A. P., Stech M. & De Oliveira S. M. 2020. — The two faces of Mt Gede, Java – Species richness, composition and zonation of epiphytic bryophytes. *Cryptogamie, Bryologie* 41 (6): 69–81. <https://doi.org/10.5252/cryptogamie-bryologie2020v41a6>. <http://cryptogamie.com/bryologie/41/6>

ABSTRACT

Bryophytes have been included in rather few studies of elevational gradients and studies on epiphytic bryophytes on different slopes of a single mountain have not been conducted in Indonesia. This research aims to compare species richness pattern, species composition, and zonation of epiphytic bryophytes on tree trunks on two different slopes of Mt Gede, West Java, Indonesia. We established an elevational transect on the North-North-East (Cibodas) and the South-South-West (Selabintana) slopes of the mountain. The transects ranged from 1500 to 2700 m each and sampling was carried out at 200 m intervals. The Cibodas slope is richer than the Selabintana slope, with 160 species (71 mosses and 89 liverworts) compared to 149 species (57 mosses and 92 liverworts), respectively. Species richness pattern on both slopes showed quadratic responses for mosses and cubic responses for liverworts. Pairwise elevation comparison between slopes showed that species composition is significantly different except for the highest plot (2700 m). Pairwise elevation comparison within the two transects revealed significant species composition changes at different elevations, indicating the presence of an elevational zonation of epiphytic bryophytes along the slopes. Based on overall bryophyte species composition, five elevational zones were recognized on each slope.

KEY WORDS

Elevational gradient,
mountain slope,
mosses,
liverworts,
Indonesia.

RÉSUMÉ

Les deux visages du mont Gede, Java – Richesse en espèces, composition et zonation des bryophytes épiphytes. Les bryophytes n'ont été considérés que dans peu d'études de gradients altitudinaux. Les études sur les bryophytes épiphytes le long des pentes d'une même montagne n'ont pas été menées en Indonésie. La recherche, présentée ici, compare le patron de richesse en espèces, la composition spécifique et la zonation des bryophytes épiphytes sur des troncs d'arbres de deux pentes différentes du mont Gede, Java ouest, Indonésie. Un transect altitudinal a été réalisé sur les pentes nord-nord-est (Cibodas) et sud-sud-ouest (Selabintana) du mont Gede. Les transects organisés entre 1500 à 2700 m et l'échantillonnage ont été conduits tous les 200 m. La pente Cibodas avec 160 espèces (71 mousses et 89 hépatiques) est plus riche que la pente Selabintana, avec 149 espèces (57 mousses et 92 hépatiques). Le patron de richesse spécifique sur les deux pentes a montré des réponses quadratiques pour les mousses et cubiques pour les hépatiques. La comparaison par paires entre les pentes a montré que la composition des espèces est significativement différente sauf pour les points les plus hauts (2700 m). La comparaison par paires de l'altitude dans les deux transects a révélé des changements significatifs de la composition spécifique à différentes altitudes, montrant la présence d'une zonation altitudinale des bryophytes épiphytes le long des pentes. Sur la base de la composition spécifique globale des bryophytes, cinq zones altitudinales ont été reconnues sur chaque pente.

MOTS CLÉS

Gradient d'altitude,
aspect de la pente,
mousses,
hépatiques,
Indonésie.

INTRODUCTION

Biodiversity changes along elevational gradients have fascinated scientists around the world, starting in the early 1800s with Humboldt in the Ecuadorian Andes, Darwin in the Chilean Andes in 1839 and 1859, and Wallace in Indonesia in 1876 and 1878 (Lomolino 2001). Various aspects of species response to the environmental differences along these gradients have been addressed since then. For bryophytes, these aspects include: 1) species richness patterns, composition, distribution, biomass, cover, and life form changes; 2) bryozone formation based on those community descriptors and assessment of the significance of certain taxa for each zone; 3) comparison between mountains; and 4) understanding the factors influencing the subjects above.

Few studies specifically dealt with richness and composition of epiphytic bryophytes along elevational gradients (e.g. Van Reenen & Gradstein 1983; Wolf 1993; Mucina *et al.* 2000; Ah-Peng *et al.* 2012; Sanger & Kirkpatrick 2015; Song *et al.* 2015; Tabua *et al.* 2017), and even fewer focused on South-East Asian countries (Thailand, Chantanaorrapint 2010; Indonesia, Gradstein & Culmsee 2010; Permadi 2015; Malaysia, Suleiman & Jotan 2015). As in other groups of organisms (Rahbek 1995; 2004; Nogués-Bravo *et al.* 2008; Fischer *et al.* 2011), the most common pattern observed for epiphytic bryophytes is the hump-shaped (quadratic) pattern. While species richness was shown to follow a common pattern, species composition changes along the elevational gradient were not as clear. Kessler (2000) found distinct elevational boundaries in the floristic composition of epiphytic bryophytes in a Colombian transect analyzed by Wolf (1993), but those boundaries were not distinctly detected along a Peruvian transect studied by Gradstein & Frahm (1987). Variation within one transect was observed by Sanger & Kirkpatrick (2015), where moss species com-

position was similar between the two highest elevations and most significantly different between the lowest ones. Distinct epiphytic bryophyte communities were recorded on eastern Viti Levu, Fiji Islands (Tabua *et al.* 2017). Floristic composition of bryophytes, species numbers, the combination of those two, estimation of cover, the spectra of life form, and phytomass estimation were used to infer forest zonation (Frahm 1990; Frahm & Gradstein 1991; Frahm *et al.* 2003). In South-East Asia, studies of the bryophyte zonation and the species associated with them were conducted on Huon Peninsula of Papua New Guinea (Enroth 1990) and Mt Kinabalu (Frahm 1990; Kürschner 1990; Frey *et al.* 1990).

Very few studies of elevational gradients included different slopes of the same mountain. Van Reenen (1987) summarized a comparison of the three Colombian Cordilleras using total percentage cover of terrestrial and epiphytic bryophytes, showing differences in zone boundaries between east and west slopes. Subsequently, Gradstein *et al.* (1989) compared two slopes of the central of the three Cordilleras, where the wet western slope has more understory bryophyte species than the drier eastern slope. No studies have been conducted in South-East Asia comparing different slopes of a mountain.

In a study on the elevational distribution of lichens and mosses on Mt Gede, West Java, Indonesia, Seifriz (1924) recognized two moss-rich zones on the Cibodas slope of the mountain, each with a distinct species composition. The first zone was located at *c.* 1700–2100 m a.s.l. and the second at *c.* 2400–2700 m. According to Seifriz (1924), the zones below 1700 m and above 2700 m were moss-poor zones dominated by lichens. Enroth (1990) argued that Seifriz's collecting efforts were insufficient and that some species reported to be exclusive to Seifriz's zones were actually widespread. He then mentioned that there were separate moss



FIG. 1. — Mt Gede, showing the SSW (left side) and the NNE (right side) slopes.

zones, but they were not as clear-cut as Seifríz suggested. The existence of what seemed like two hump-shaped patterns of bryophyte species richness on this mountain, each with a distinct species composition, and the question whether the same zonation occurs on other slopes of the mountain, still needs to be investigated.

Here we deal with the elevational gradients of epiphytic bryophytes of Mt Gede by comparing the North-North-East (NNE) slope with the South-South-West (SSW) one in terms of species richness and composition. We hypothesize that a hump-shaped species richness pattern exists on each slope for epiphytic mosses, liverworts, and total bryophytes. Species composition is hypothesized to indicate the presence of an elevational zonation corresponding to vegetational and topographic differences within slope, and to differ between slopes at the same elevation due to climate. We further discuss the zonation formed by dissimilarities of species composition in the light of climate and vegetation differences.

MATERIAL AND METHODS

STUDY AREA

The study was conducted in Gede-Pangrango National Park, West Java, Indonesia. It is the oldest National Park in Indonesia and encompasses two volcanoes, Mt Gede (2958 m a.s.l.) and Mt Pangrango (3109 m). Both mountains are connected through a saddle at approximately 2400 m. Following Van

Steenis (1972), the vascular plant vegetation consists of three zones, the Submontane (<1500 m), Montane (1500–2400 m), and Subalpine (>2400 m) zone.

SAMPLING

Along an elevational gradient ranging from 1500 to 2700 m, bryophytes were sampled in two transects on Mt Gede: one transect on the NNE slope explored through the Cibodas track and the other transect on the SSW slope along the Selabintana track (Fig. 1). Along each transect, seven sample sites were chosen with 200 m intervals. At each site, ten trees with diameter at breast height (DBH) larger than 20 cm were sampled. Trees selected were separated from each other by at least 15 m, belonged to different species and differed in bark texture as much as possible. Tree species, DBH and elevation above sea level of each tree were recorded. At higher elevations, where tree diameters were smaller, we summed up smaller trunks of the same tree in order to balance sampling effort. Epiphytic bryophytes (mosses and liverworts) were collected from the first zone of tree zonation (Johansson 1974; 0–2 m above ground). All material associated with this study is vouchered and deposited at Cibodas Botanic Garden Herbarium (CHTJ). Voucher specimens were identified by standard practice of using identification keys, mainly Bartram (1939), Eddy (1988, 1990, 1996), and Gradstein (2011), and followed the nomenclature within “Tropicos” (n.d.) and Söderström *et al.* (2010, 2016). Only specimens identified to species level were used in the data analysis. The identification results were recorded as presence and absence

of species per tree per elevation site, representing species richness per elevation.

DATA ANALYSIS

All statistical analyses were performed using R software (R Core Team 2018) packages. Overall species richness similarity between Selabintana and Cibodas slope was tested using permutation test with function 'independence_test' in 'coin' package (Hothorn *et al.* 2008). Multiple pairwise comparisons of species richness among elevations from both slopes were tested using Kruskal-Wallis test. Dunn post hoc test with p-values adjusted with Benjamini-Hochberg method was performed following rejection of null hypothesis of Kruskal-Wallis test. Dunn tests were performed using 'FSA' package (Ogle *et al.* 2018) function 'dunnTest'. Correlations among variables were checked using correlation test with Spearman index. Correlation test function 'rcorr' was used from 'Hmisc' Package (Harrell Jr. *et al.* 2018). These tests were performed to match the data properties following data exploration procedure by Zuur *et al.* (2010).

Generalized Linear Models (GLM) with Poisson distribution and log link were used to relate species richness and elevation, with tree surface area added as covariate. Since overdispersion was detected, the standard errors were corrected using quasipoisson-GLM models. Generalized Linear Models were performed using 'glm' function. Comparison of full model (polynomials of third-degree order) with simpler models was used to choose for best model. Nonsignificant terms were dropped stepwise by eliminating the most complex term and re-running the selection until only the significant term and the lower order term were left. Best models were chosen after running diagnostic tests.

The relation between species composition and elevation was analyzed using Permutational Multivariate Analysis of Variance (PERMANOVA) with Bray-Curtis dissimilarity index. The homogeneity of dispersion on the species composition was checked using Permutational Analysis of Multivariate Dispersions (PERMDISP). Each elevation site was used as grouping. Permutation tests (9999 for PERMANOVA and 999 for PERMDISP, respectively) were applied to test the significance. Package 'vegan' (Oksanen *et al.* 2018) functions 'adonis' and 'betadisper' were used to compute PERMANOVA and PERMDISP, respectively. Species composition of a particular elevation was assigned to a different zone than the adjacent elevation when PERMANOVA results showed significant p-values ($p \leq 0.01$) and PERMDISP showed non-significant p-values.

Indicator Species Analysis was conducted to test whether zones distinguished by their bryophyte species composition were associated with elevation. The Indicator Species Analysis was performed using 'indicspecies' package (De Cáceres & Legendre 2009) functions 'multipatt' with 9999 permutations. Indicator species were selected when the Squared Indicator Value $> 30\%$ and Benjamini-Hochberg p-adjusted value = 0.05, the latter correcting for false positives in the original, lower p-values that resulted from the analysis (De Cáceres & Legendre 2009).

RESULTS

SPECIES RICHNESS

The Cibodas slope yielded a higher number of bryophyte species (160, of which 71 mosses and 89 liverworts) than the Selabintana slope (149, of which 57 mosses and 92 liverworts) ($Z = -3.1451$, $p\text{-value} = 0.0017$), as shown in Figure 2. Overall species richness of bryophytes in each elevational belt was higher on Selabintana slope, and species richness of liverworts was generally higher than that of mosses on both slopes, albeit not significantly (data not shown). Exceptions were observed at 1500 and 1900 m on Cibodas slope, where richness of mosses was higher.

SPECIES RICHNESS PATTERNS

The variable 'tree surface area' was eliminated during the process of selecting the best model, leaving elevation as the only variable in the model. Total bryophyte species richness showed a significant quadratic response pattern, resulting in a hump-shaped curve for Cibodas slope that peaked at 2000 m (Fig. 3A). For Selabintana slope, in contrast, a cubic (hump-shaped, then increasing) response was observed peaking at 1900 m (Fig. 3A). Both Cibodas and Selabintana species richness showed a quadratic response pattern for mosses and a cubic pattern for liverworts (Fig. 3B, C). Moss species richness peaked at 1900 m on the Cibodas slope and at 2100 m on Selabintana, whereas liverwort richness peaked at 1900 m on both slopes and increased again above 2400 m.

SPECIES COMPOSITION

The PERMANOVA analyses for pairwise elevations within each slope (Table 1A) revealed different species compositions of bryophytes on both slopes, indicating the presence of an elevational zonation of epiphytic bryophytes. The PERMDISP analyses revealed that bryophyte compositions were homogeneous ($p\text{-value} < 0.01$) except for Cibodas pairwise elevation at 2500-2700 m. At corresponding sampling elevations on opposite slopes, total bryophyte species composition was generally different (Table 1B).

The zones and respective indicator species are listed in Table 2 and their diagrammatic visualization is shown in Figure 4. Within each slope, five bryophyte zones were distinguished based on the compositional similarity between elevations (Table 2, Fig. 4). Selabintana slope had a wider zone 2, which was comprised of three sampling elevation instead of only two on Cibodas. On the other hand, zone 5 on Cibodas slope was one sampling elevation wider than the corresponding zone on Selabintana. Other than that, the zonation between those two were similar in width. Except for zone 4 on Selabintana, each bryophyte zone had its own indicator species (Table 2, Fig. 4). No indicator species were shared among zones of Cibodas and Selabintana, except for *Syrrhopodon tristichus* Nees ex Schwägr. in zone 2. On Cibodas slope, 10 species were resolved as indicator species with indicator values ranging from 32.7% to 77.1%, while Selabintana slope had 13 indicator species with indicator value ranging from 31.3% to 90.9% (Table 2).




FIG. 2. — Species richness of **A**, Cibodas and **B**, Selabintana slopes along the elevational gradient.

DISCUSSION

Our study shows that elevational gradients of epiphytic bryophytes on Mt Gede are expressed by changes in species richness as well as species composition, and differ among slopes. The differences between slopes seem to be associated with climatic and vegetational differences.

SPECIES RICHNESS: DIFFERENT PATTERNS FOR MOSSES AND LIVERWORTS

While the drier Cibodas slope is overall significantly richer in epiphytic bryophyte species, the wetter Selabintana slope

has more species than Cibodas at corresponding elevations although the differences were not significant. Dearborn & Danby (2017) observed similar patterns in alpine-subarctic plant communities. In that study, the drier, south-facing slope had a higher β -diversity than the wetter, north-facing slope, while the mean species richness per elevation between aspects was similar, which might have been caused by less pronounced vegetation changes on the wetter, north-facing slope. On Mt Gede a more variable topography and more pronounced forest stand changes are seen on the Cibodas slope. Climbing up this slope from lower elevation, one passes a river, small lake, swamp, hot water river, waterfall, and a steep

TABLE 1. — Non-parametric analysis of variance for pairwise elevation comparison within slope **A** and between slopes **B** for Cibodas and Selabintana bryophytes. Pairwise comparisons of group mean dispersions are shown as F ratio and p value (9999 permutations). Pairwise comparisons of permutational analysis of variance are shown as coefficient of determination R^2 and p value (999 permutations). ** = $p \leq 0.01$; *** = $p \leq 0.001$.

A	Permdisp		Permanova	
Pairwise elevation	F	p	R ²	p
Cibodas				
1500-1700	0.814	0.370	0.155	0.001 ***
1700-1900	0.096	0.760	0.120	0,027
1900-2100	0.255	0.618	0.139	0.005 **
2100-2300	0.000	0.998	0.202	0.001 ***
2300-2500	1.137	0.299	0.173	0.001 ***
2500-2700	21.700	0.0001 ***	0.298	0.001 ***
Selabintana				
1500-1700	0.162	0.697	0.160	0.004 **
1700-1900	0.570	0.458	0.115	0,018
1900-2100	0.580	0.452	0.078	0,138
2100-2300	4.154	0.055	0.258	0.001 ***
2300-2500	3.762	0.067	0.202	0.001 ***
2500-2700	0.114	0.741	0.281	0.001 ***

B	Permdisp		Permanova	
Pairwise elevation	F	p	R ²	p
Cibodas-Selabintana				
1500	0.557	0.465	0.156	0.001 ***
1700	0.281	0.602	0.168	0.003 **
1900	0.331	0.572	0.170	0.003 **
2100	0.241	0.630	0.241	0.001 ***
2300	2.111	0.163	0.326	0.001 ***
2500	0.014	0.906	0.308	0.001 ***
2700	17.781	0.0005 ***	0.336	0.001 ***

trail up to the summit, whereas the trail on the Selabintana slope mainly follows a ridge and then passes a river up to the summit, without much topographic variation. The more variable topography and more pronounced forest stand changes on the Cibodas slope probably caused the higher epiphytic bryophytes species richness seen on this slope.

The generally higher liverwort diversity observed on each elevational gradient in this study is in accordance with Frahm (2002), Gradstein *et al.* (1989), Santos *et al.* (2017), Song *et al.* (2015), Tabua *et al.* (2017) and Wolf (1993). In Neotropical moist forests, liverworts are the dominant component in the epiphytic layer (Cornelissen & Ter Steege 1989; Gradstein *et al.* 2001). This is also the case with Mt Gede, where mosses are outnumbered from the very beginning of the elevational gradients. According to Gradstein *et al.* (1989), the ratio of mosses and liverworts reflects the humidity of a location, where a higher proportion of liverworts signifies higher humidity. The higher liverwort richness in Selabintana should reflect the wetter climate on this slope as compared with Cibodas.

The quadratic response pattern for total bryophytes richness on the Cibodas slope, resulting in a hump-shaped curve, confirms previous observations (Wolf 1993; Ah-Peng *et al.* 2012; Sanger & Kirkpatrick 2015; Song *et al.* 2015). However, different patterns were observed for mosses (quadratic) and liverworts (cubic) on both slopes of Mt Gede. A similar cubic pattern was observed for terrestrial bryophytes along

an altitudinal gradient in China (Sun *et al.* 2013). In contrast, Grau *et al.* (2007), Henriques *et al.* (2016) and Tabua *et al.* (2017) did not find differences between mosses and liverworts, both showing quadratic patterns. Another pattern was found by Bruun *et al.* (2006), who observed increasing species richness with elevation for both mosses and liverworts in an alpine environment, the increase being steeper for liverworts. In the Canary Islands, on the other hand, Hernández-Hernández *et al.* (2017) found a significantly quadratic pattern for liverworts but not for mosses. In Cibodas, moss and total bryophyte richness exhibited a similar pattern, while in Selabintana liverworts matched the total bryophyte richness pattern. In the Colombian Andes, Wolf (1993) also observed that the overall bryophyte pattern was mainly determined by the liverwort component, while the moss pattern was similar to that of lichens.

The quadratic response pattern on Mt Gede for total bryophytes is attributed to the suitability of the microclimate at mid-elevation for bryophyte growth. The peak richness of total bryophytes at mid-elevation coincides with maximum humidity due to cloud belt formation at 2000 m (Van Steenis 1972). The decline towards higher elevation is most likely due to the combination of lesser cloud frequency (pers. obs.) and the decrease in tree size above 2000 m (Van Steenis 1972), providing less substrate for epiphytic bryophytes. Another possible explanation for the quadratic pattern would be the Mid Domain Effect (MDE), where the peak of the species richness at mid elevation is formed by the stochastic overlapping of species ranges along the gradient. In the latter case, the pattern observed should be the same for mosses and liverworts, which was not the case in the present study. Grau *et al.* (2007) found quadratic patterns for mosses and liverworts in the Himalayas that peaked at different elevations and argued that such patterns could be explained by the interaction between climate and species richness. They suggested that MDE may not be important for bryophytes. On Mt Gede, the vegetation zonation of the forest stands might explain the cubic pattern for liverworts. The last increase in liverwort richness with elevation occurs at approximately 2400 m and coincides with the upper limit of the montane vegetation zone. Above that elevation the subalpine zone starts, characterized by lower, denser, thinner-stemmed, more open forest with single canopy layer (Van Steenis 1972). This zone promotes liverworts better than mosses, presumably due to the higher light intensity. Increasing humidity along a transect causes an increase in liverwort richness (Frahm & Ohlemüller 2001). More open forest means higher overall light intensity and light transmission to the forest floor, allowing sun-tolerant species to grow at lower heights on the trunks (Acebey *et al.* 2003) and thus adding to liverwort species richness. Another explanation could be non-proportional sampling, with taxa sampled from the smaller trees at high elevation possibly including those of higher tree zones. Since more liverworts are known to inhabit higher vertical zones within a tree in tropical rainforests (Cornelissen & Ter Steege 1989; Wolf 1993; Holz 2003; Mota de Oliveira & Ter Steege 2013), the species richness of liverworts is thus further increased.




FIG. 3. — The relationship between species richness of **A**, bryophytes; **B**, mosses; and **C**, liverworts and elevation with GLM functions superimposed for the Cibodas (left) and Selabintana (right) slopes.

SPECIES COMPOSITION: SLIGHT DIFFERENCES IN ZONATION ACCORDING TO SLOPE

Comparing pairwise elevations across the Cibodas and Selabintana slopes, in general, species richness did not significantly differ, but species composition did. This might be due

to the different climate on each slope, with Cibodas (NNE) being drier than Selabintana (SSW). The moisture brought by western monsoon winds from the Pacific and Indian Ocean is released as rain on the windward Selabintana slope, leaving the leeward Cibodas slope relatively less wet. The difference

TABLE 2. — Indicator species of elevational zones on the Cibodas and Selabintana slopes. **A** and **B** represent the ‘specificity’ and the ‘fidelity’ components of indicator value. Only (IndVal)² > 30% and p-adj. ≤ 0.05 are shown.

Slope	Elevation (m a.s.l.)	zone	Indicator species	A	B	IndVal	(IndVal) ² (%)	p-adj.
Cibodas	1500	1	<i>Leucophanes angustifolium</i> Renauld & Cardot	1.000	0.400	0.632	40.0	0.007
	1700-1900	2	<i>Plagiochila spathulifolia</i> Mitt.	0.929	0.650	0.777	60.4	0.001
			<i>Syrrhopodon tristichus</i> Nees ex Schwägr.	1.000	0.400	0.632	40.0	0.005
			<i>Lejeunea micholitzii</i> Mizut.	1.000	0.400	0.632	40.0	0.006
	2100	3	<i>Hypnodendron dendroides</i> (Brid.) A. Touw	0.714	0.500	0.598	35.7	0.012
			<i>Trachyloma indicum</i> Mitt.	0.546	0.600	0.572	32.7	0.023
	2300	4	<i>Pogonatum cirratum</i> (Sw.) Brid.	0.875	0.700	0.783	61.3	0.001
			<i>Distichophyllum brevicuspis</i> M. Fleisch	0.750	0.600	0.671	45.0	0.002
	2500-2700	5	<i>Bazzania japonica</i> (Sande Lac.) Lindb.	0.857	0.900	0.878	77.1	0.001
			<i>Herbertus dicranus</i> (Gottsche, Lindenb. et Nees) Trevis.	1.000	0.650	0.806	65.0	0.001
Selabintana	1500	1	<i>Heteroscyphus argutus</i> (Reinw., Blume et Nees) Schiffn.	0.600	0.600	0.600	36.0	0.013
	1700-2100	2	<i>Syrrhopodon tristichus</i> Nees ex Schwägr.	0.944	0.567	0.732	53.5	0.002
			<i>Bazzania javanica</i> (Sande Lac.) Schiffn.	0.842	0.533	0.670	44.9	0.006
	2300	3	<i>Symphysodontella attenuatula</i> M. Fleisch.	0.909	1.000	0.953	90.9	0.001
			<i>Fissidens wichurae</i> Broth. & M. Fleisch	1.000	0.400	0.632	40.0	0.009
	2500	4	—	—	—	—	—	—
	2700	5	<i>Plagiochila arbuscula</i> (Brid. ex Lehm. et Lindenb.) Lindenb.	0.900	0.900	0.900	81.0	0.001
			<i>Lepidozia stahliae</i> Steph.	1.000	0.500	0.707	50.0	0.001
			<i>Cylindrocolea kiaeri</i> (Austin) Váňa	0.636	0.700	0.667	44.5	0.002
			<i>Cryptolophocolea ciliolata</i> (Nees) L.Söderstr.	0.833	0.500	0.645	41.7	0.005
			<i>Heteroscyphus aselliformis</i> (Reinw., Blume et Nees) Schiffn.	0.833	0.500	0.645	41.7	0.005
			<i>Drepanolejeunea ternatensis</i> (Gottsche) Schiffn.	0.500	0.800	0.632	40.0	0.011
			<i>Frullania junghuhniana</i> Gottsche	0.800	0.400	0.566	32.0	0.024
			<i>Bazzania praerupta</i> (Reinw., Blume et Nees) Trevis.	0.625	0.500	0.559	31.3	0.023

between wet and dry season in the West Javan mountains is not as pronounced as in East Java. Rainfall recordings on Mt Gede’s twin peak Mt Pangrango show that even during the driest months, rain still falls every other day (Docters van Leeuwen 1933; Van Steenis 1972). And since Java is positioned south of the equator, the NE side of Mt Gede receives more sunshine than its SW side (Van Steenis 1972). Similar climatic conditions are observed on Mt Kilimanjaro (Pepin *et al.* 2017) – also located in the southern hemisphere – where the SW slope is generally cooler and moister, and receives sunshine later during the day than the NE slope.

The climatic settings of different slopes influence the vegetation, as observed on Mt Kilimanjaro (Hemp 2006) and on Baima Snow Mountain (Yang *et al.* 2016) where changes in forest species composition along the elevational gradient differ between slopes. This, in turn, together with microclimatic differences, affects the species composition of the epiphytic vegetation component, including the bryophytes. In the case of Mt Gede, there are more bryophyte species in Selabintana at single sampling elevations than in Cibodas, but the differences were not significant.

The forest stand components create a microclimate suitable for particular bryophytes assemblages. The degree of vegetation density around the phorophyte might influence the effect of sunlight and wind at the lower trunk. Király & Ódor (2010) found that sapling density increased the epiphytic bryophyte species richness significantly and had a considerable effect on species composition. In addition, tree species contribute to epiphytic bryophyte composition (Barkman 1958, Sim-Sim

et al. 2011, Mežaka *et al.* 2012) through variation in bark texture (Szövényi *et al.* 2004), bark chemical properties (Mežaka *et al.* 2012, Batista & Santos 2016, Dislich *et al.* 2018), and shaded cover (Ariyanti *et al.* 2008). The more heterogeneous the phorophyte composition, the more variable the bryophyte species composition.

Another factor possibly causing differences in bryophyte composition between the two slopes is human disturbance. Visitors of Gede-Pangrango National Park usually climb up to the peak via the Cibodas track as it is more easily accessible, the main office of the National Park is located at the entrance, more water resources are available on the way up, and the visitor quota allowed to enter daily through this track is three times that of Selabintana (Ketentuan Umum Penda- kian Gunung Gede Pangrango (n.d.); 300 persons per day). Although this study was conducted in sites with minimal human disturbance, it is possible that human activities along the track had influenced the studied bryophyte communities.

BRYOPHYTE ZONATION ON MT GEDE IN A BROADER PERSPECTIVE

Applying the tropical rainforest zonation terminology using bryophytes proposed by Frahm (1990) and Frahm & Gradstein (1991), bryophyte composition changes on Mt Gede correspond with the vegetation zonation in other studies on the Mt Gede-Mt Pangrango twin mountain, as shown in Figure 5.

The numbers and width of zones varied between those studies. A minimum of three zones, Tropical Lower Mon-




FIG. 4. — Diagrammatic visualization of zonation along the elevational gradient of the Selabintana and Cibodas slopes with their indicator species. For taxa authors see Table 2.

tane, Tropical Upper Montane, and Tropical Subalpine, were recognized in all of them. On Mt Gede, the lowest elevation zone corresponds to the Tropical Submontane zone and the highest to the Subalpine forest of Van Steenis (1972). The Tropical Submontane zone was not part of the transect on Mt Pangrango (Ohsawa 1991) and thus unrecognized. The zones on Mt Gede (zone 2, 4, 5; this study) are mostly narrower than on neighbouring Mt Pangrango (except for zone 2, see below), which may be due to the 'Massenerhebung' effect where the width of a forest zone was narrower on the smaller mountain than on the bigger one (Grubb 1971). The wider zone 2 (Tropical Lower Montane zone) on Mt Gede should be due to the wetter climate on the Selabintana slope, as mentioned before. This finding is in accordance with Van Reenen (1987), Gradstein *et al.* (1989), and Frahm & Gradstein (1991) who found that the width of the Tropical Montane zone is narrower on drier slopes, and its boundaries different accordingly. The wider zone 5 (Tropical Subalpine) on the Cibodas slope as shown from the species composition dispersion (PERMDISP) analyses, was due to the nestedness of the species composition, with the species at higher elevation being a subset of the lower one. Phorophyte richness on the Cibodas slope was highly reduced in the summit site and dominated here by *Vaccinium varingaefolium* Miq., which was also part of the phorophyte composition at lower elevations. The reduced phorophyte richness at the summit might be due to the harsh environmental conditions here, including the effect of toxic substances emitted from the crater of Mt Gede.

Zone 1 (Tropical Submontane) corresponds to Seifriz's moss-poor *Altingia* zone. The next three zones comprised Van Steenis' Montane zone. In his elevational zonation scheme of the mountains of Java, Van Steenis (1972) did not divide the Montane zone into subzones. Ohsawa (1991) proposed to divide Van Steenis' Subalpine zone into an Upper Montane and Subalpine zone, as supported by his recognition of Lower and Upper Montane zones on Mt Pangrango, the twin mountain of Gede. Our study using bryophytes reflects the detailed forest stand zonation of Mt Gede and shows that the Montane zone has three divisions – Tropical Lower Montane, transition, Tropical Upper Montane –, which was also recorded by Arrijani (2008) based on an analysis of tree diversity on the Cibodas slope. These three zones correspond to Seifriz's *Podocarpus*, herbs, and *Vaccinium* zones, respectively. His observation of the moss-rich zone was also confirmed by the present study in which epiphytic bryophytes peaked in the Tropical Lower Montane zone and then declined in the transition zone. On the opposite slope (Selabintana), Permadi (2015) found a similar bryophyte zonation although he did not further subdivide the Montane zone.

Zone 3 can be regarded as a transition zone between the Tropical Lower Montane and Upper Montane zone, due to its significantly different species composition that consists of a mix of species from adjacent zones. Yamada (1976) mentioned the presence of a forest transition zone on Mt (Gede-) Pangrango, situated on the saddle shared by the two mountains with a maximum width at 2300 m.




Fig. 5. — Elevational zonation comparison on mountains of Java, especially Mt Gede-Pangrango. Zonations were based on bryophytes coverage (Seifriz 1924), species composition dissimilarity (this study) and higher plants composition (Ohsawa 1991, Seifriz 1923, Van Steenis 1972).

Since bryophyte species composition is influenced by forest stand, the presence of a transitional forest stand along the elevational gradient explains the existence of a bryophyte transition zone. Despite using Yamada's data, Ohsawa (1991) did not recognize the transition zone and included it in a wider Lower Montane Zone. The methodologies applied by Ohsawa (1991) to determine the zonation of Mt (Gede-) Pangrango probably did not allow for a transition zone to be detected, since such inference would require species turnover analysis (cf. Bach & Gradstein 2011). The Subalpine zone in the latter study coincided within Seifriz's *Vaccinium* zone (approximately 2400-2700 m). Although described by Seifriz

as the second 'moss-rich' zone, the present study found that it is actually a liverwort-rich zone.

The characterization of the zones on the Selabintana and Cibodas slope of Mt Gede by species indicators reinforces the differences in species composition. Different plant communities inhabiting the same zone on opposite slopes were also observed by Kappelle *et al.* (1995) in Costa Rica. Since epiphytic bryophytes are influenced by phorophyte communities, the occurrence of different bryophyte communities in the same zone on opposite slopes is to be expected. Selabintana has more liverworts as indicator species, reflecting the wetter condition and higher liverwort richness on this slope. The

results indicate that altitudinal zonation is more strongly driven by slope conditions than by elevation.

Acknowledgements

We thank the Research and Innovation in Science and Technology Project (RISET-Pro) of the Ministry of Research, Technology and Higher Education of the Republic of Indonesia (World Bank Loan No. 8245-ID) for the PhD scholarship awarded to the first author. The initial research was supported by the PPKP 2011 Incentive Program of the Indonesian Budget for the Ministry of Research, Technology and Higher Education in 2011. The first author is also grateful to the Mt Gede Pangrango National Park authority (BBTNGGP) of the Ministry of Environment and Forestry of the Republic of Indonesia for the fieldwork permit and the expedition team from Cibodas Botanic Garden for support during the field work. The authors thank Robbert Gradstein and an anonymous reviewer for their valuable contributions.

REFERENCES

- ACEBEY A., GRADSTEIN S. R. & KRÖMER T. 2003. — Species richness and habitat diversification of corticolous bryophytes in submontane rain forest and fallows of Bolivia. *Journal of Tropical Ecology* 19 (1): 9-18. <https://doi.org/10.1017/S026646740300302X>
- AH-PENG C., WILDING N., KLUGE J., DESCAMPS-JULIEN B., BARDAT J., CHUAH-PETIOT M., STRASBERG D. & HEDDERSON T. A. J. 2012. — Bryophyte diversity and range size distribution along two altitudinal gradients: Continent vs. island. *Acta Oecologica* 42: 58-65. <https://doi.org/10.1016/j.actao.2012.04.010>
- ARIYANTI N. S., BOS M. M., KARTAWINATA K., TJITROSOED-IRDJO S. S., GUHARDA E. & GRADSTEIN S. R. 2008. — Bryophytes on tree trunks in natural forests, selectively logged forests and cacao agroforests in Central Sulawesi, Indonesia. *Biological Conservation* 141 (10): 2516-2527. <https://doi.org/10.1016/j.biocon.2008.07.012>
- ARRIJANI A. 2008. — Struktur dan Komposisi Vegetasi Zona Montana Taman Nasional Gunung Gede Pangrango. *Biodiversitas* 9 (2): 134-141. <https://doi.org/10.13057/biodiv/d090212>
- BACH K. & GRADSTEIN S. R. 2011. — A comparison of six methods to detect altitudinal belts of vegetation on tropical mountains. *Ecotropica* 17 (1): 1-13.
- BARKMAN J. J. 1958. — Phytosociology and ecology of cryptogamic epiphytes, Van Gorcum, Assen. 628 p.
- BARTRAM E. B. 1939. — Mosses of the Philippines. *Philippine Journal of Science* 68: 437 p.
- BATISTA W. V. S. M. & SANTOS N. D. 2016. — Can regional and local filters explain epiphytic bryophyte distributions in the Atlantic Forest of southeastern Brazil? *Acta Botanica Brasiliica* 30 (3): 462-472. <https://doi.org/10.1590/0102-33062016abb0179>
- BRUUN H. H., MOEN J., VIRTANEN R., GRYNES J.-A., OKSANEN L. & ANGERBJÖRN A. 2006. — Effects of altitude and topography on species richness of vascular plants, bryophytes and lichens in alpine communities. *Journal of Vegetation Science* 17 (1): 37-46. [https://doi.org/10.1658/1100-9233\(2006\)017\[0037:EOAAT O\]2.0.CO;2](https://doi.org/10.1658/1100-9233(2006)017[0037:EOAAT O]2.0.CO;2)
- CHANTANAORRAPINT S. 2010. — Ecological Studies of Epiphytic Bryophytes Along Altitudinal Gradients in Southern Thailand. Dissertation, Universität Bonn. 107 p. Retrieved from <http://www.archive-for-bryology.com/Special%20volume%206.pdf>
- CORNELISSEN J. H. C. & TER STEEGE H. 1989. — Distribution and ecology of epiphytic bryophytes and lichens in dry evergreen forest of Guyana. *Journal of Tropical Ecology* 5 (2): 131-150. <https://doi.org/10.1017/S0266467400003400>
- DE CACERES M. & LEGENDRE P. 2009. — Associations between species and groups of sites: Indices and statistical inference. *Ecology* 90 (12): 3566-3574. <http://sites.google.com/site/miqueldecaceres/>
- DEARBORN K. D. & DANBY R. K. 2017. — Aspect and slope influence plant community composition more than elevation across forest-tundra ecotones in subarctic Canada. *Journal of Vegetation Science* 28 (3): 595-604. <https://doi.org/10.1111/jvs.12521>
- DISLICH R., PINHEIRO E. M. L. & GUIMARÃES M. 2018. — Corticolous liverworts and mosses in a gallery forest in Central Brazil: Effects of environmental variables and space on species richness and composition. *Nova Hedwigia* 107 (3): 385-406. https://doi.org/10.1127/nova_hedwigia/2018/0480
- DOCTERS VAN LEEUWEN W. M. 1933. — Biology of plants and animals occurring in the higher parts of Mount Pangrango-Gedeh in West-Java. Verhandelingen der Koninklijke Akademie van Wetenschappen te Amsterdam, Afdeling Natuurkunde (tweede sectie) Deel 31. Noord-Hollandsche Uitgevers-Maatschappij, Amsterdam.
- EDDY A. 1988. — Sphagnales to Dicranales, in *A Handbook of Malesian Mosses. Vol. 1*. British Museum of Natural History, London, 204 p.
- EDDY A. 1990. — Leucobryaceae to Buxbaumiaceae, in *A Handbook of Malesian Mosses. Vol. 2*. Natural History Museum Publications, London, 256 p.
- EDDY A. 1996. — Splachnobryaceae to Leptostomataceae, in *A Handbook of Malesian Mosses. Vol. 3*. Natural History Museum and HMSO, London, 276 p.
- ENROTH J. 1990. — Altitudinal zonation of bryophytes on the Huon Peninsula, Papua New Guinea. A floristic approach, with phytogeographic considerations. *Tropical Bryology* 2: 61-90. <https://doi.org/10.11646/bde.2.1.6>
- FISCHER A., BLASCHKE M. & BÄSSLER C. 2011. — Altitudinal gradients in biodiversity research: The state of the art and future perspectives under climate change aspects. *Waldökologie, Landschaftsforschung und Naturschutz* 11: 35-47.
- FRAHM J.-P. 1990. — The altitudinal zonation of bryophytes on Mt Kinabalu. *Nova Hedwigia*, 51: 133-149.
- FRAHM J.-P. 2002. — Ecology of bryophytes along altitudinal and latitudinal gradients in Chile. *Tropical Bryology* 21: 67-79. <https://doi.org/10.11646/bde.21.1.12>
- FRAHM J.-P. & GRADSTEIN S. R. 1991. — An altitudinal zonation of tropical rain forests using bryophytes. *Journal of Biogeography* 18: 669-678. <https://doi.org/10.2307/2845548>
- FRAHM J.-P. & OHLEMÜLLER R. 2001. — Ecology of bryophytes along altitudinal and latitudinal gradients in New Zealand. Studies in austral temperate rain forest bryophytes 15. *Tropical Bryology* 20: 117-137. <https://doi.org/10.11646/bde.20.1.20>
- FRAHM J.-P., PÓCS T., O'SHEA B., KOPONEN T., PIIPPO S., ENROTH J., RAO P. & FANG Y. 2003. — Manual of tropical bryology. *Tropical Bryology* 23: 1-196. Retrieved from <http://core.ac.uk/download/pdf/14530151.pdf>
- FREY W., GOSSOW R. & KÜRSCHNER H. 1990. — Verteilungsmuster von Lebensformen, wasserleitenden und wasserspeichernden Strukturen in epiphytischen Moosgesellschaften am Mt Kinabalu (Nord-Borneo). *Nova Hedwigia* 51: 87-119.
- GRADSTEIN S. R. 2011. — Guide to the Liverworts and Hornworts of Java. SEAMEO-BIOTROP, Bogor, 146 p.
- GRADSTEIN S. R. & CULMSEE H. 2010. — Bryophyte diversity on tree trunks in montane forests of Central Sulawesi, Indonesia. *Bryophyte Diversity and Evolution* 31 (1): 95-105. <https://doi.org/10.11646/bde.31.1.16>
- GRADSTEIN S. R., GRIFFIN III D., MORALES M. I. & NADKARNI N. M. 2001. — Diversity and habitat differentiation of mosses and liverworts in the cloud forest of Monteverde, Costa Rica. *Caldasia* 23 (1): 203-212. Retrieved from www.jstor.org/stable/23641822
- GRADSTEIN S. R., VAN REENEN G. B. A. & GRIFFIN III D. 1989. —

- Species richness and origin of the bryophyte flora of the Colombian Andes. *Acta Botanica Neerlandica* 38 (4): 439-448. <https://doi.org/10.1111/j.1438-8677.1989.tb01375.x>
- GRADSTEIN S. R. & FRAHM J.-P. 1987. — Die floristische Höhengliederung der Moose entlang des BRYOTROP-Transektes in NO-Peru. *Beihefte Nova Hedwigia* 88: 105-113.
- GRAU O., GRYNES J.-A. & BIRKS H. J. B. 2007. — A comparison of altitudinal species richness patterns of bryophytes with other plant groups in Nepal, Central Himalaya. *Journal of Biogeography* 34 (11): 1907-1915. <https://doi.org/10.1111/j.1365-2699.2007.01745.x>
- GRUBB P. J. 1971. — Interpretation of the 'Massenerhebung' effect on tropical mountains. *Nature* 229: 44-45. <https://doi.org/10.1038/229044a0>
- HARRELL JR F. E. 2018. — Hmisc: Harrell Miscellaneous. R package version 4.1-1 with contributions from Charles Dupont and many others. <https://CRAN.R-project.org/package=Hmisc>
- HEMP A. 2006. — Continuum or zonation? Altitudinal gradients in the forest vegetation of Mt Kilimanjaro. *Plant Ecology* 184 (1): 27-42. <https://doi.org/10.1007/s11258-005-9049-4>
- HENRIQUES D. S. G., BORGES P. A. V., AH-PENG C. & GABRIEL R. 2016. — Mosses and liverworts show contrasting elevational distribution patterns in an oceanic island (Terceira, Azores): The influence of climate and space. *Journal of Bryology* 38 (3): 183-194. <https://doi.org/10.1080/03736687.2016.1156360>
- HERNÁNDEZ-HERNÁNDEZ R., BORGES P. A., GABRIEL R., RIGAL F., AH-PENG C. & GONZÁLEZ-MANCEBO J. M. 2017. — Scaling a-and b-diversity: bryophytes along an elevational gradient on a subtropical oceanic Island (La Palma, Canary Islands). *Journal of Vegetation Science* 28: 1209-1219. <https://doi.org/10.1111/jvs.12573>
- HOLZ I. 2003. — Diversity and Ecology of Bryophytes in Montane Quercus Forests of Costa Rica. Dissertation, Universität zu Göttingen. 172 p. Retrieved from <https://d-nb.info/972640606/34>
- HOTHORN T., HORNIK K., VAN DE WIEL M. A. & ZEILEIS A. 2008. — Implementing a class of permutation tests: The coin package. *Journal of Statistical Software* 28 (8): 1-23. <http://www.jstatsoft.org/v28/i08/>
- JOHANSSON D. 1974. — Ecology of vascular epiphytes in West African rain forest. *Acta Phytogeographica Suecica* 59: 1-136.
- KAPPELLE M., VAN UFFELEN J. G. & CLEEF A. M. 1995. — Altitudinal zonation of montane *Quercus* forests along two transects in the Chirripó National Park, Costa Rica. *Vegetatio* 119: 119-153
- KESSLER M. 2000. — Altitudinal zonation of Andean cryptogam communities. *Journal of Biogeography* 27 (2): 275-282. <https://doi.org/10.1046/j.1365-2699.2000.00399.x>
- KETENTUAN UMUM PENDAKIAN GUNUNG GEDE PANGRANGO. N.D. — <https://gedepangrango.org/booking/> (accessed 09 April 2019).
- KIRÁLY I. & ÓDOR P. 2010. — The effect of stand structure and tree species composition on epiphytic bryophytes in mixed deciduous-coniferous forests of Western Hungary. *Biological Conservation* 143 (9): 2063-2069. <https://doi.org/10.1016/j.biocon.2010.05.014>
- KÜRSCHNER H. 1990. — Die epiphytischen Moosgesellschaften am Mt Kinabalu (Nord-Borneo, Sabah, Malaysia). *Nova Hedwigia*, 51: 1-75.
- LOMOLINO M. V. 2001. — Elevation gradients of species-density: Historical and prospective views. *Global Ecology and Biogeography* 10 (1): 3-13. <https://doi.org/10.1046/j.1466-822x.2001.00229.x>
- MEŽÁKA A., BRÜMELIS G. & PĪTERĀNS A. 2012. — Tree and stand-scale factors affecting richness and composition of epiphytic bryophytes and lichens in deciduous woodland key habitats. *Biodiversity and Conservation* 21 (12): 3221-3241. <https://doi.org/10.1007/s10531-012-0361-8>
- MOTA DE OLIVEIRA S. & TER STEEGE H. 2013. — Floristic overview of the epiphytic bryophytes of terra firme forests across the Amazon basin. *Acta Botanica Brasilica* 27 (2): 347-363. <https://doi.org/10.1590/S0102-33062013000200010>
- MUCINA L., VALACHOVIC M., DIMOPOULOS P., TRIBSCH A. & PISUT I. 2000. — Epiphytic lichen and moss vegetation along an altitude gradient on Mount Aenos (Kefallinia, Greece). *Biologia, Bratislava* 55 (1): 43-54
- NOGUÉS-BRAVO D., ARAÚJO M. B., ROMDAL T. & RAHBK C. 2008. — Scale effects and human impact on the elevational species richness gradients. *Nature* 453 (7192): 216-219. <https://doi.org/10.1038/nature06812>
- OGLE D. H., WHEELER P. & DINNO A. 2018. — FSA: Fisheries Stock Analysis. R package version 0.8.22. <https://github.com/droglenc/FSA>
- OHSAWA M. 1991. — Structural comparison of tropical montane rain forests along latitudinal and altitudinal gradients in south and east Asia. *Vegetatio* 97 (1): 1-10
- OKSANEN J., BLANCHET F. G., FRIENDLY M., KINDT R., LEGENDRE P., MCGLINN D., MINCHIN P. R., O'HARA R. B., SIMPSON G. L., SOLYMOS P., STEVENS M. H. H., SZOECS E. & WAGNER H. 2018. — Vegan: Community Ecology Package. R package version 2.5-2. <https://CRAN.R-project.org/package=vegan>
- PEPIN N. C., PIKE G., SCHAEFER M., BOSTON C. M. & LOVELL H. 2017. — A comparison of simultaneous temperature and humidity observations from the SW and NE slopes of Kilimanjaro: The role of slope aspect and differential land-cover in controlling mountain climate. *Global and Planetary Change* 157: 244-258. <https://doi.org/10.1016/j.gloplacha.2017.08.006>
- PERMADI D. 2015. — Lumut Epifit (Mosses, Divisio: Bryophyta) Sepanjang Gradien Ketinggian Di Taman Nasional Gunung Gede Pangrango, Jawa Barat. Bachelor Thesis. Bandung, Institut Teknologi Bandung, 56 p.
- R CORE TEAM. 2018. — R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- RAHBK C. 1995. — The elevational gradient of species richness: A uniform pattern? *Ecography* 18 (2): 200-205. <https://doi.org/10.1111/j.1600-0587.1995.tb00341.x>
- RAHBK C. 2004. — The role of spatial scale and the perception of large-scale species-richness patterns: Scale and species-richness patterns. *Ecology Letters* 8 (2): 224-239. <https://doi.org/10.1111/j.1461-0248.2004.00701.x>
- SANGER J. C. & KIRKPATRICK J. B. 2015. — Moss and vascular epiphyte distributions over host tree and elevation gradients in Australian subtropical rainforest. *Australian Journal of Botany* 63 (8): 696-704. <https://doi.org/10.1071/BT15169>
- SANTOS N. D., COSTA D. P., KINOSHITA L. S. & SHEPHERD G. J. 2017. — Variations in bryophyte communities in a short elevational gradient in Atlantic Forest of Southeastern Brazil. *Cryptogamie, Bryologie* 38 (2): 191-211. <https://doi.org/10.7872/cryb/v38.iss2.2017.191>
- SEIFRIZ W. 1923. — The altitudinal distributions of plants on Mt Gedeh, Java. *Bulletin of the Torrey Botanical Club* 50 (9): 283-306.
- SEIFRIZ W. 1924. — The altitudinal distribution of lichens and mosses on Mt Gedeh, Java. *The Journal of Ecology* 12 (2): 307-313. <https://doi.org/10.2307/2255252>
- SIM-SIM M., BERGAMINI A., LUÍS L., FONTINHA S., MARTINS S., LOBO C. & STECH M. 2011. — Epiphytic bryophyte diversity on Madeira Island: Effects of tree species on bryophyte species richness and composition. *The Bryologist* 114 (1): 142-154. <https://doi.org/10.1639/0007-2745-114.1.142>
- SÖDERSTRÖM L., GRADSTEIN S. R. & HAGBORG A. 2010. — Checklist of the hornworts and liverworts of Java. *Phytotaxa* 9 (1): 53-149. <https://doi.org/10.11646/phytotaxa.9.1.7>
- SÖDERSTRÖM L., HAGBORG A., VON KONRAT M., BARTHOLOMEW-BEGAN S., BELL D., BRISCOE L., BROWN E., CARGILL D. C., COSTA D. P., CRANDALL-STOTLER B. J., COOPER E. D., DAUPHIN G., ENGEL J. J., FELDBERG K., GLENNY D., GRADSTEIN S. R., HE X., HEINRICHS J., HENTSCHEL J., ILKIU-BORGES A. L., KATAGIRI T., KONSTANTINOVA N. A., LARRAÍN J., LONG D. G., NEBEL M., PÓCS T., PUCHE F., REINER-DREHWALD E., REN-

- NER M. A., SASS-GYARMATI A., SCHÄFER-VERWIMP A., MORAGUES J. G., STOTLER R. E., SUKKHARAK P., THIERS B. M., URIBE J., VÁÑA J., VILLARREAL J. C., WIGGINTON M., ZHANG L. & ZHU R. L. 2016. — World checklist of hornworts and liverworts. *PhytoKeys* 59: 1-828. <https://doi.org/10.3897/phytokeys.59.6261>
- SONG L., MA W.-Z., YAO Y.-L., LIU W.-Y., LI S., CHEN K., LU H.-Z., CAO M., SUN Z.-H. & TAN Z.-H. 2015. — Bole bryophyte diversity and distribution patterns along three altitudinal gradients in Yunnan, China. *Journal of Vegetation Science* 26 (3): 576-587. <https://doi.org/10.11646/phytotaxa.9.1.7>
- SULEIMAN M. & JOTAN P. 2015. — Diversity of epiphytic mosses along an altitudinal gradient at Minduk Sirung Trail in Crocker Range Park, Sabah, Malaysia. *Sepilok Bulletin* 21 (22): 49-58
- SUN S.-Q., WU Y.-H., WANG G.-X., ZHOU J., YU D., BING H.-J. & LUO J. 2013. — Bryophyte species richness and composition along an altitudinal gradient in Gongga Mountain, China. *PLoS ONE* 8 (3): e58131. <https://doi.org/10.1371/journal.pone.0058131>
- SZÖVÉNYI P., HOCK Z. S. & TÓTH Z. 2004. — Phorophyte preferences of epiphytic bryophytes in a stream valley in the Carpathian Basin. *Journal of Bryology* 26 (2): 137-146. <https://doi.org/10.1179/037366804225021092>
- TABUA M., RILEY R., RENNER M. A. M., SODERSTROM L., HAGBORG A. & VON KONRAT M. 2017. — Are epiphytic bryophyte communities characterized by changes along an elevational gradient?—A preliminary study on eastern Viti Levu, Fiji Islands. *Bryophyte Diversity and Evolution* 39 (1): 28-37. <https://doi.org/10.11646/bde.39.1.6>
- TROPICOS. N.D. — Missouri Botanical Garden. <http://www.tropicos.org/>
- VAN REENEN G. B. A. & GRADSTEIN S. R. 1983. — Studies on Colombian Cryptogams XX. A transect analysis of the bryophyte vegetation along an altitudinal gradient on the Sierra Nevada de Santa Marta, Colombia. *Acta Botanica Neerlandica* 32 (3): 163-175.
- VAN REENEN G. B. A. 1987. — Altitudinal bryophyte zonation in the Andes of Colombia: A preliminary report. *Symposia Biologica Hungarica* 35: 631-637.
- VAN STEENIS C. G. G. J. 1972. — The Mountain Flora of Java. Leiden, E. J. Brill, 147 p.
- WOLF J. H. D. 1993. — Diversity patterns and biomass of epiphytic bryophytes and lichens along an altitudinal gradient in the Northern Andes. *Annals of the Missouri Botanical Garden* 80 (4): 928-960. <https://doi.org/10.2307/2399938>
- YAMADA I. 1976. — Forest ecological studies of the montane forest of Mt Pangrango, West Java. Stratification and floristic composition of the forest vegetation of the higher part of Mt Pangrango. *South East Asian Studies* 13: 514-534.
- YANG Y., SHEN Z., HAN J. & ZHONGYONG C. 2016. — Plant diversity along the eastern and western slopes of Baima Snow Mountain, China. *Forests* 7 (4): 89. <https://doi.org/10.3390/f7040089>
- ZUUR A. F., IENO E. N. & ELPHICK C. S. 2010. — A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution* 1 (1): 3-14. <https://doi.org/10.1111/j.2041-210X.2009.00001.x>

*Submitted on 13 June 2019;
accepted on 27 January 2020;
published on 6 May 2020.*