

cryptogamie

Bryologie

2021 • 42 • 9

Koponobryum papillosum Printarakul & Chantanaorr., sp. nov. (Pottiaceae, Bryophyta),
a new moss species from northern Thailand

Narin PRINTARAKUL, Arunothai JAMPEETONG, Udon PONGKAWONG,
Phiangphak SUKKHARAK, Ekaphan KRAICHAK,
Kanonrat ADULKITTICHAI & Sahut CHANTANAORRAPINT

DIRECTEUR DE LA PUBLICATION / *PUBLICATION DIRECTOR*: Bruno David,
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Denis LAMY

ASSISTANTE DE RÉDACTION / *ASSISTANT EDITOR*: Marianne SALAÜN (bryo@cryptogamie.com)

MISE EN PAGE / *PAGE LAYOUT*: Marianne SALAÜN

RÉDACTEURS ASSOCIÉS / *ASSOCIATE EDITORS*

Biologie moléculaire et phylogénie / Molecular biology and phylogeny

Bernard GOFFINET

Department of Ecology and Evolutionary Biology, University of Connecticut (United States)

Mousses d'Europe / European mosses

Isabel DRAPER

Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid (Spain)

Francisco LARA GARCÍA

Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid (Spain)

Mousses d'Afrique et d'Antarctique / African and Antarctic mosses

Rysiek OCHYRA

Laboratory of Bryology, Institute of Botany, Polish Academy of Sciences, Krakow (Pologne)

Bryophytes d'Asie / Asian bryophytes

Rui-Liang ZHU

School of Life Science, East China Normal University, Shanghai (China)

Bioindication / Biomonitoring

Franck-Olivier DENAYER

Faculté des Sciences Pharmaceutiques et Biologiques de Lille, Laboratoire de Botanique et de Cryptogamie, Lille (France)

Écologie des bryophytes / Ecology of bryophyte

Nagore GARCÍA MEDINA

Department of Biology (Botany), and Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid (Spain)

COUVERTURE / *COVER*:

Extraits d'éléments de la Figure 1 / Extracts of the Figure 1

Cryptogamie, Bryologie est indexé dans / *Cryptogamie, Bryologie* is indexed in:

- Biological Abstracts
- Current Contents
- Science Citation Index
- Publications bibliographiques du CNRS (Pascal).

Cryptogamie, Bryologie est distribué en version électronique par / *Cryptogamie, Bryologie* is distributed electronically by:

- BioOne® (<http://www.bioone.org>)

Cryptogamie, Bryologie est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Cryptogamie, Bryologie is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi/ The Museum Science Press also publish: *Adansonia*, *Geodiversitas*, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Comptes Rendus Palevol*, *Cryptogamie* sous-sections *Algologie*, *Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2021

ISSN (imprimé / print): 1290-0796 / ISSN (électronique / electronic): 1776-0992

***Koponobryum papillosum* Printarakul & Chantanaorr., sp. nov. (Pottiaceae, Bryophyta), a new moss species from northern Thailand**

**Narin PRINTARAKUL
Arunothai JAMPEETONG
Udon PONGKAWONG**

Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200 (Thailand)

Phiangphak SUKKHARAK

Department of Biology, Faculty of Science, Burapha University, Chonburi 20131 (Thailand)

Ekaphan KRAICHAK

Department of Botany, Faculty of Science, Kasetsart University,
Chatuchak, Bangkok 10900 (Thailand)

Kanonrat ADULKITTICHAJ

Research and development department, Chiangmai Vanussanun Co.,
Ltd., Chiang Mai 50000, (Thailand)

Sahut CHANTANAORRAPINT

SU Herbarium, Division of Biological Science, Faculty of Science,
Prince of Songkla University, Hat Yai, Songkhla 90110 (Thailand)
sahut.c@psu.ac.th (corresponding author)

Submitted on 8 January 2021 | Accepted on 14 April 2021 | Published on 16 June 2021

Printarakul N., Jampeetong A., Pongkawong U., Sukkharak P., Kraichak E., Adulkittichai K. & Chantanaorrapint S. 2021. — *Koponobryum papillosum* Printarakul & Chantanaorr., sp. nov. (Pottiaceae, Bryophyta), a new moss species from northern Thailand. *Cryptogamie, Bryologie* 42 (9): 143–148. <https://doi.org/10.5252/cryptogamie-bryologie2021v42a9>. <http://cryptogamie.com/bryologie/42/9>

ABSTRACT

KEY WORDS
Bryophyte diversity,
Koponobryum bengalense,
pottiaceous moss,
northern Thailand,
new species.

Koponobryum papillosum Printarakul & Chantanaorr., sp. nov., is described as a second species of the genus found in Chiang Mai and Lamphun Provinces, northern Thailand. It is distinguished from *K. bengalense* (Gangulee) Arts by autoicous plants, papillose spores and axillary hairs consisting of one short basal cell and 1–4(5) longer upper cells. A taxonomic description, illustrations and SEM micrographs are presented.

RÉSUMÉ

Koponobryum papillosum Printarakul & Chantanaorr., sp. nov. une nouvelle espèce de mousse du nord de la Thaïlande.

Koponobryum papillosum Printarakul & Chantanaorr., sp. nov. est décrit comme une deuxième espèce du genre trouvée dans les provinces de Chiang Mai et Lamphun, dans le nord de la Thaïlande. Elle se distingue de *K. bengalense* (Gangulee) Arts par des plantes auto-portantes, des spores papillaires et des poils axillaires constitués d'une cellule basale courte et de 1–4(5) cellules supérieures plus longues. Une description taxonomique, des illustrations et des microographies MEB sont présentées.

MOTS CLÉS
Diversité des bryophytes,
Koponobryum bengalense,
mousse pottiacée,
Thaïlande du nord,
espèce nouvelle.

INTRODUCTION

The genus *Koponobryum* Arts was first established by Arts (2001) with a single species, *K. bengalense* (Gangulee) Arts, a species endemic to India. The genus was segregated from *Splachnobryum* Hal. Müll. by the following combination of characters: bordered leaves, the acute leaf apex, the excurrent costa, and the upper lamina cells bearing one central papilla (Arts 2001). Although *Koponobryum* was originally placed in the Splachnobryaceae (Arts 2001), further molecular studies show that the Splachnobryaceae nested in the Pottiaceae (e.g., Werner *et al.* 2004; Frey & Stech 2009; Cox *et al.* 2010).

During the bryological surveys in Chiang Mai and Lamphun Provinces, northern Thailand, some interesting specimens of pottiaceous mosses were collected. They have lanceolate to oblong-lanceolate leaves with acute to acuminate apex and decurrent base, upper lamina cells bearing one central papilla, leaf margins bordered by 1-3 rows of elongate cells, costa excurrent and costa in cross section with one stereid band. This combination of characters implies the genus *Koponobryum*, but the collection did not match *K. bengalense*. It is therefore described here as a species new to science.

MATERIAL AND METHODS

This study is based on recent collections from Thailand. Voucher specimens of the new species are deposited in BKF, CMUB, and PSU herbaria. Morphological and anatomical characters were studied using stereo and compound microscopes. The distinctive characters of the species were illustrated. Mature spores and peristome were and examined with JEOL JSM-5910 LV scanning electron microscope. In addition, detailed descriptions, illustrations, and notes on species of *Koponobryum* are provided.

TAXONOMIC TREATMENT

Family SPLACHNOBRYACEAE A.K. Kop.
Genus *Koponobryum* Arts

Koponobryum papillosum
Printarakul & Chantanaorr., sp. nov.
(Figs 1-3)

DIAGNOSIS. — Similar to *K. bengalense* (Gangulee) Arts, but differing in autoicous plants, papillose spores, and axillary hairs consisting of one short basal cell and 1-4(5) longer upper cells.

TYPE. — Thailand. Chiang Mai Province: Mae Rim District, Doi Suthep-Pui National Park, Mae Sa Falls, 18°54'25.7"N, 98°53'46.6"E, c. 420 m elev., 19.VII.2020, N. Printarakul & K. Adulkittichai 19072020_1 (holo-, CMUB!; iso-, BKF!, PSU!).

ADDITIONAL SPECIMENS EXAMINED. — Thailand. Chiang Mai Province: Mae Rim District, Doi Suthep-Pui National Park, Mae

Sa Falls, c. 420 m elev., on soil under granitic rock, 18°54'25.7"N, 98°53'46.6"E, 4.X.2013, N. Printarakul 6444 (CMUB); Chiang Dao District, Doi Chang Dao Wildlife Sanctuary, Huay Sob On forest protection unit, c. 710 m elev., on soil under calcareous rocks, 19°21'33.7"N, 98°43'07.5"E, 19.VIII.2017, N. Printarakul 7280 (CMUB, PSU). Lamphun Province: Sri Bua Ban District, the Hari-phuchai Education Centre of Chiang Mai University, c. 500 m elev., on soil under calcareous-sandstones, 18°32'17.33"N, 99°07'28.10"E, 22.VIII.2020, N. Printarakul, A. Jampeetong & U. ongkawong 22082020_15 (CMUB).

ETYMOLOGY. — The specific epithet “*papillosum*” refers to the papillose spores.

HABITAT AND DISTRIBUTION. — *Koponobryum papillosum* Printarakul & Chantanaorr., sp. nov. is currently known only from northern Thailand. It grows on disturbed soils on a degraded calcareous substrate, sandstones and granitic rocks in deciduous, dipterocarp-oak, and also in mixed evergreen-deciduous seasonal forests at elevation of 420-710 m.

CONSERVATION STATUS. — IUCN (2012) category Least concern (LC). Although only three locations were found in Thailand, they are within the well-protected areas. Therefore, this species is not under immediate threat.

DESCRIPTION

Plants

Up to 17 mm high (including sporophytes), pale to light green, forming loose turfs.

Rhizoids

Reddish to pale brown, clustered at stem base.

Stems

Erect, unbranched, dark green to reddish brown, in cross section 8-10 layers of cells across, 120-150 µm in diameter; outer cortical cells somewhat smaller with moderately thick-walled, inner cortical cells larger with thicker walls; central stand consisting of smaller and thin-walled cells.

Axillary hairs

Filiform, uniseriate, 2-5(-6) cells long, the basal cell shortest and yellowish to pale brown, the upper cells distinctly longer and colorless.

Leaves

Contorted when dry, erecto-patent to spreading when moist; lanceolate to oblong-lanceolate, 1.1-1.5 × 0.3-0.4 mm; apex acute to acuminate; base slightly decurrent; margins plane above, slightly revolute below, bordered throughout by 1-3 rows of elongate cells, remotely serrulate to denticulate above the midleaf; costa single, stout, short excurrent, ventral and dorsal surfaces smooth, not prominently bulging; costa cross section at midleaf semicircular to elliptic, with one stereid band; lamina unistratose, orange to red with KOH; upper laminal cells subquadrangular, polygonal to rounded-hexagonal, (20-) 25-30 µm in diameter, with a central papilla on both sides, firm to slightly thick-walled; basal marginal cells elongate rectangular, (35-) 40-80 × 12.5-15 µm wide, thin-walled; basal juxta-costal cells broader, rectangular, (37.5-) 50-90 × 25-35 µm, thin-walled.




FIG. 1. — *Koponobryum papillosum* Printarakul & Chantanaorr., sp. nov.: A, plants in natural habitat; B, sporophyte bearing plants; C, leaves; D, leaf apex; E, upper laminal cells and marginal cells; F, leaf bases; G, sporophyte; H, capsule; I, peristome and spores; J, calyptra. Photos by N. Printarakul. All from N. Printarakul & K. Adulkittichai 19072020_1 (CMUB). Scales bars: A, G, H, 1 mm; B, 3 mm; C, J, 0.5 mm; D-F, I, 50 µm.




Fig. 2. – *Koponobryum papillosum* Printarakul & Chantanaorr., sp. nov.: **A**, KOH color reaction of plant; **B**, cross section of stem; **C**, vegetative leaf; **D-G**, cross sections of leaf from upper to base; **H**, perichaetium and perigonium; **I-L**, perichaetal leaves; **M-O**, perigonal leaves; **P**, axillary hair. Photos by N. Printarakul. All from N. Printarakul & K. Adulkittichai 19072020_1 (CMUB). Scale bars: A, 1 mm; B, D-G, P, 25 µm; C, H-O, 0.5 mm.




FIG. 3.—*Koponobryum papillosum* Printarakul & Chantanaorr., sp. nov.: A, plants with sporophytes; B, cross section of stem; C, variations of axillary hairs; D, leaves; E, cross section of midleaf; F, leaf apex; G, leaf base; H, peristome (SEM); I, spore (SEM). A-G. Drawn by N. Printarakul. All from N. Printarakul & K. Adulkittitchai 19072020_1 (CMUB). Scale bars: A, 1 mm; B, C, F, G, 50 µm; D, 0.5 mm; E, 25 µm; H, 10 µm; I, 1 µm.

Asexual reproduction

Not seen.

AUTOICOUS

Perigonia

Lateral on short branches, perigonial leaves ovate to lanceolate, smaller than stem leaves, apex acute to acuminate, margins entire; antheridia 10-15 per perigonium; paraphysis absent.

Perichaetia

Terminal, perichaetal leaves ovate-lanceolate, seldom lingulate, smaller than stem leaves or sometimes rudimentary, apex acute to acuminate or sometimes obtuse, margins entire, weakly sheathing at base; paraphyses absent.

Setae

4-6 mm long, smooth, yellowish brown in upper part, reddish in lower part.

Capsules

Erect, symmetrical, oblong-cylindrical, up to 1.3 mm long, smooth.

Opercula

Conical, reddish brown, c. 0.25 mm long.

Peristome

Single, reddish brown, consisting of 16 linear teeth, 175-190 µm high, papillose.

Spores

Spherical, 12.5-15 µm in diameter, finely papillose.

Calyptra

Cucullate, smooth.

TAXONOMIC NOTES

The new species is characterized by a combination of the following characters: 1) autoicous sexual condition; 2) orange to red coloration in KOH; 3) lanceolate to oblong-lanceolate leaves with acute to acuminate apex and decurrent base; 4) a short-excurrent costa with single stereid band in cross section; 5) leaf bordered by 1-3 rows of elongate cells; 6) leaf margins being serrulate to denticulate above the midleaf; 7) upper laminal cells bearing a central papilla on both sides;

8) filiform axillary hairs composed of 2-5(6) cells long; and 9) oblong-cylindrical capsules with conical operculum; 10) peristome consisting of 16 linear papillose teeth; and 11) papillose spores.

Koponobryum papillosum Printarakul & Chantanaorr., sp. nov. resembles the type species of the genus, *K. bengalense*, in having bordered leaves consisting of 1-3 rows of elongate cells, leaf apex being acute to acuminate, the upper lamina cells having a central papilla on both sides, and central strand distinctly differentiated. *Koponobryum bengalense*, however, differs from the new species by dioicous plants, smooth spores, and axillary hairs consisting of two short basal cells and one longer apical cell.

Acknowledgements

We would like to thank the Department of National Parks, Wildlife and Plant Conservation of Thailand for permission to collect specimens; and R.H. Zander (MO) and anonymous reviewers for their useful suggestions. The first author is deeply grateful to the many years of encouragement given by Dr K. Santanachote, the late Dr B.C. Tan and the late J.F. Maxwell. This research was supported by the Thailand Research Fund (no. DBG6080015) and the plant genetic conservation project under the royal initiation of Her Royal Highness Princess Maha Chackri Sirindhorn-Chiang Mai University Centre (RSPG-CMU).

REFERENCES

- ARTS T. 2001. — A revision of the Splachnobryaceae (Musci). *Lindbergia* 26: 77-96. <https://doi.org/10.2307/20150068>
- COX C. J., GOFFINET B., WICKETT N. J., BOLES S. B. & SHAW A. J. 2010. — Moss diversity: A molecular phylogenetic analysis of genera. *Phytotaxa* 9: 175-195. <https://doi.org/10.11646/phytotaxa.9.1.10>
- FREY W. & STECH M. 2009 — Bryophyta, in FREY W. (ed.), *Sylloabus of Plant Families - A. Engler's Syllabus der Pflanzenfamilien, 13e Part 3: Bryophytes and seedless Vascular Plants*. Borntraeger, Berlin: 116-257.
- IUCN 2012. — *IUCN Red List categories and criteria: version 3.1, second edition*. IUCN, Gland Switzerland and Cambridge, United Kingdom.
- WERNER O., ROS R. M., CANO M. J. & GUERRA J. 2004. — Molecular phylogeny of Pottiaceae (Musci) based on chloroplast rps4 sequence data. *Plant Systematics and Evolution* 243: 147-164. <https://doi.org/10.1007/s00606-003-0076-0>

Submitted on 8 January 2021;
accepted on 14 April 2021;
published on 16 June 2021.