



The last record of an ailuropod bear from the Iberian Peninsula

Juan ABELLA,
Daniel HONTECILLAS,
Alberto VALENCIANO,
Plinio MONTOYA,
Jorge MORALES,
María Dolores PESQUERO &
Luis ALCALÁ



DIRECTEUR DE LA PUBLICATION : Bruno David,
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF* : Didier Merle

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS* : Emmanuel Côté (geodiv@mnhn.fr) ; Anne Mabilie

MISE EN PAGE / *PAGE LAYOUT* : Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD* :

Christine Argot (MNHN, Paris)
Beatrix Azanza (Museo Nacional de Ciencias Naturales, Madrid)
Raymond L. Bernor (Howard University, Washington DC)
Alain Blicq (chercheur CNRS retraité, Haubourdin)
Henning Blom (Uppsala University)
Jean Broutin (UPMC, Paris)
Gaël Clément (MNHN, Paris)
Ted Daeschler (Academy of Natural Sciences, Philadelphie)
Bruno David (MNHN, Paris)
Gregory D. Edgecombe (The Natural History Museum, Londres)
Ursula Göhlich (Natural History Museum Vienna)
Jin Meng (American Museum of Natural History, New York)
Brigitte Meyer-Berthaud (CIRAD, Montpellier)
Zhu Min (Chinese Academy of Sciences, Pékin)
Isabelle Rouget (UPMC, Paris)
Sevket Sen (MNHN, Paris)
Stanislav Štámbek (Museum of Eastern Bohemia, Hradec Králové)
Paul Taylor (The Natural History Museum, Londres)

COUVERTURE / *COVER* :

Made from the Figures of the article.

Geodiversitas est indexé dans / *Geodiversitas is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Physical, Chemical, and Earth Sciences®
- Scopus®

Geodiversitas est distribué en version électronique par / *Geodiversitas is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Geodiversitas* sont référencés par /
Articles and nomenclatural novelties published in Geodiversitas are referenced by:

- ZooBank® (<http://zoobank.org>)

Geodiversitas est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Geodiversitas is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:

Adansonia, Zoosystema, Anthropolozologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Algologie, Bryologie, Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél. : 33 (0)1 40 79 48 05 / Fax : 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2019
ISSN (imprimé / *print*) : 1280-9659/ ISSN (électronique / *electronic*) : 1638-9395

The last record of an ailuopod bear from the Iberian Peninsula

Juan ABELLA

Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona,
c/ Columnes s/n, Campus de la UAB, 08193 Cerdanyola del Vallès, Barcelona (Spain)
and Instituto Nacional de Biodiversidad,
Pje. Rumipamba N. 341 y Av. de los Shyris (Parque La Carolina) Quito (Ecuador)
juan.abella@icp.cat (corresponding author)

Daniel HONTECILLAS

Departamento de Paleobiología, Museo Nacional de Ciencias Naturales-CSIC,
C/José Gutiérrez Abascal, 2, 28006 Madrid (Spain)
daniel.hontecillastamay@gmail.com

Alberto VALENCIANO

Research and Exhibitions Department, Iziko Museums of South Africa,
P.O. Box 61, Cape Town (South Africa)
and Department of Biological Sciences, University of Cape Town, Palaeobiological Research Group,
Private Bag X3, Rhodes Gift 7701, Cape Town (South Africa)
alb3rtovv@gmail.com

Plinio MONTOYA

Departament de Botànica i Geologia, Àrea de Paleontologia,
Universitat de València, Doctor Moliner 50, 46100 Burjassot, València (Spain)
plinio.montoya@uv.es

Jorge MORALES

Departamento de Paleobiología, Museo Nacional de Ciencias Naturales-CSIC,
C/José Gutiérrez Abascal, 2, 28006 Madrid (Spain)
mcnm166@mncn.csic.es

María Dolores PESQUERO

Departamento de Paleobiología, Museo Nacional de Ciencias Naturales-CSIC,
C/José Gutiérrez Abascal, 2, 28006, Madrid (Spain)
and Fundación Conjunto Paleontológico de Teruel-Dinópolis & Museo Aragonés de Paleontología,
Avda. Sagunto s/n, 44002 Teruel (Spain)
pesquero@fundaciondinopolis.org

Luis ALCALÁ

Fundación Conjunto Paleontológico de Teruel-Dinópolis & Museo Aragonés de Paleontología,
Avda. Sagunto s/n, 44002 Teruel (Spain)
alcala@dinopolis.com

Submitted on 1 January 2019 | accepted on 3 June 2019 | published on 2 December 2019

urn:lsid:zoobank.org:pub:4174E526-4A7D-4D2E-9233-6FD43F145F07

Abella J., Hontecillas D., Valenciano A., Montoya P., Morales J., Pesquero M. D. & Alcalá L. 2019. — The last record of an ailuopod bear from the Iberian Peninsula, in Bonis L. de & Werdelin L. (eds), Memorial to Stéphane Peigné: Carnivores (Hy-aenodonta and Carnivora) of the Cenozoic. *Geodiversitas* 41 (23): 797-809. <https://doi.org/10.5252/geodiversitas2019v41a23>. <http://geodiversitas.com/41/23>

ABSTRACT

In this paper we describe Late Miocene (MN13) remains of the genus *Indarctos* Pilgrim, 1913 from the locality of Las Casiones (Teruel, Spain). Although the phylogenetic relationships of this genus are still controversial, the most recent phylogenetic analyses, based on cranial, mandibular and dental characters, include it in Ailuropodinae, thus making the relatives of the giant panda the predominant bears in the carnivoran assemblages for most of the Late Miocene in the Iberian Peninsula. These fossils of *Indarctos punjabiensis* (Lydekker, 1884) represent the last population of this subfamily from the Iberian fossil record, and possibly also from Europe, making this an important advance in our knowledge of the evolutionary history of this group. We also note the replacement of *Indarctos* by *Agriotherium* A. Wagner, 1837 in Iberian faunas, between *c.* 6.3 and *c.* 6.23 Ma.

KEY WORDS

Ursidae,
Ailuropodinae,
Late Miocene,
Las Casiones,
Teruel Basin.

RÉSUMÉ

Le dernier enregistrement d'un ours ailuropode dans la péninsule Ibérique.

Dans cet article nous décrivons les restes fossiles du genre *Indarctos* Pilgrim, 1913 du Miocène supérieur (MN13) de la localité de Las Casiones (Teruel, Espagne). Bien que les relations phylogénétiques de ce genre soient encore controversées, dans l'analyse phylogénétique la plus récente basée sur les caractères crâniens, mandibulaires et dentaires, ils sont inclus dans les Ailuropodinae. Ces parents des pandas géants étaient donc les ours prédominants dans les assemblages de carnivores pendant la majeure partie du Miocène supérieur de la péninsule Ibérique. D'autre part, ces fossiles d'*Indarctos punjabiensis* (Lydekker, 1884) représentent la dernière population de cette sous-famille dans les archives fossiles de la péninsule Ibérique, et éventuellement aussi d'Europe, ils contribuent de manière significative à l'avancement de la connaissance de l'histoire évolutive de cet groupe. En outre, nous signalons le remplacement de *Indarctos* par *Agriotherium* Wagner, 1837 dans les faunes ibériques entre 6.3 et 6.23 Ma.

MOTS CLÉS

Ursidae,
Ailuropodinae,
Miocène supérieur,
Las Casiones,
Teruel Basin.

INTRODUCTION

The Late Turolian (or Ventian, *sensu* Morales *et al.* 2013) fossil locality of Las Casiones, MN13 of the European Continental Neogene Mammal zonation (Mein 1990; Bruijn *et al.* 1992), is located 10 km north of the city of Teruel (Spain) in the district of Villalba Baja, on one of the slopes of a wide ravine near the villages of Celadas and Villalba Baja; more accurately at 40°26'10"N, 1°3'58"W (Alcalá 1994; van Dam *et al.* 2001; Pesquero 2003) (Fig. 1). This locality is included in the local zone M2 of van Dam (1997), and an age of 6.3 Ma has been proposed (van Dam *et al.* 2006: suppl. notes; Gibert *et al.* 2013). The fossils are found in a series of levels of lacustrine origin, mostly composed of green and black clays, with occasional intercalations of fine sand (Pesquero 2003). At least two different levels can be distinguished: Las Casiones (KS) and Las Casiones Superior (KSS), the first of which is the one that has yielded the majority of fossil remains. Several papers have been already written about this locality and its faunal composition. Its geology and stratigraphy was studied by Alcalá (1994) and later by Pesquero (2003), where a more detailed model was proposed.

The latest mammal faunal composition of Las Casiones (KS) is the following: The murids *Stephanomys ramblensis* van de Weerd, 1976, *Apodemus gudrunae* van de Weerd, 1976, *Occitanomys alcalai* Adrover, Mein and Moissenet, 1988, cf. *Rhagapodemus* sp.; the cricetids *Ruscinomys schaubi* Villalta & Crusafont, 1956, *Cricetus* cf. *C. kormosi* Schaub, 1930, *Blancomys sanzi* Adrover, Mein & Moissenet, 1993; the glirid

Eliomys truci Mein & Michaux, 1970; the castorid *Dipoides problematicus* (Schlosser, 1902); the hystricid *Hystrix depereti* Sen, 2001; the erinaceid *Galerix* sp.; the talpids *Desmanella* sp., *Talpa* sp.; the soricids *Archaeodesmana* sp., *Paenelimnoecus* sp., *Blarinella* sp., *Amblycoptus jessiae* Doukas, van den Hoek Ostende, Theocharopoulos & Reumer, 1995; the vespertilionid chiropteran *Vespertilionidae* indet.; the ochotonid lagomorph *Prolagus* sp.; the proboscideans *Tetralophodon longirostris* (Kaup, 1832) and *Zygolophodon turicensis* (Schinz, 1824); the hyracoid *Pliohyrax* sp. cf. *P. graecus* (Gaudry, 1860); two species of the cervid *Pliocervus* Hilzheimer, 1922; the bovids *Tragoportax* sp. and two species of Boselaphini indet.; the hippopotamid *Hippopotamus crusafonti* Aguirre, 1963; the rhinocerotid *Dihoplus schleiermacheri* (Kaup, 1834); three species of the equid *Hipparion* Christol, 1832: *H. primigenium* (von Meyer, 1829), *H. mathewii* (Abel, 1926), and *H. periafricanum* Villalta & Crusafont, 1957; the mustelids *Plesiogulo monspessulanus* Viret, 1939, *Baranogale adroveri* (Petter, 1964), *Mustela* sp., and a Mustelidae indet. aff. *Sabadellictis* sp.; the ursid *Indarctos punjabiensis* (Lydekker, 1884); the hyaenids *Plioviverrops* sp. cf. *P. guerini* (Villalta & Crusafont, 1948) and *Thalassictis hipparionum* (Gervais, 1859); and the felids *Paramachaerodus orientalis* Kittl, 1887, *Amphimachairodus giganteus* Kretzoi, 1929, *Metailurus major* Zdansky, 1924, *Pristifelis attica* Wagner, 1857, and Felinae indet. (Cerdeño 1989; Alcalá 1994; Alcalá *et al.* 1994; van Dam 1997; Alcalá & Montoya 1998a, b; Pesquero 2003; van Weers & Rook 2003; Salesa *et al.* 2012).




FIG. 1. — Schematic geologic map of the northern part of the Teruel basin with the paleontological localities indicated (after Pesquero *et al.* 2003 modified).

TABLE 1. — Measurements (in mm) of the dentition of the specimens of the genus *Indarctos* Pilgrim, 1913 from Las Casiones (length × width).

Specimens	P2	P4	M1	M2
K-174	10.45 × 7.42	—	—	—
K94-1733	—	32.01 × 22.51	—	—
K-162	—	—	29.21 × 28.98	—
K-161	—	—	× 26.89	—
KS-4316	—	—	28.62 × 27.01	—
KS-4315	—	—	—	31.34 × 27.79
KS-4317	—	—	—	31.49 × 28.06

The assignment of the bear fossil remains from Las Casiones to the genus *Indarctos* Pilgrim, 1913 might be considered as a novelty, since Alcalá (1994) described as *Agriotherium* Wagner, 1837 a very worn M1 and a fragment of M1, due to their large width. This identification has subsequently been repeated in many papers. The presence of *Indarctos* in Las Casiones was also tentatively noted by both Alcalá *et al.* (2005) and Salesa *et al.* (2012). This change in taxonomic determination has been possible due to the recovery of the new diagnostic remains described in this paper.

THE GENUS *INDARCTOS* IN THE FOSSIL RECORD

The genus *Indarctos* comprises medium to large-sized bears from the late Miocene, with an omnivorous diet (Montoya *et al.* 2001; Viranta 2004; Abella 2011; Abella *et al.* 2013; Monesillo *et al.* 2014; Domingo *et al.* 2016). Although comprehensively studied, there is not a full consensus about their phylogenetic relationships. Some authors (Hendey 1972; Qiu *et al.* 2014) relate them to the genus *Agriotherium* (considered a Hemicyonidae by the authors of the present paper) and others consider it to be a basal Ursidae (Abella *et al.* 2012, 2014). However, the most recent phylogenetic analyses, based on cranial, mandibular and dental characters, include *Indarctos* spp. in the Ailuropodinae (Abella *et al.* 2012; Qiu *et al.* 2014). This genus was indeed the predominant bear taxon during most of the late Miocene in the Northern Hemisphere (Abella *et al.* 2014).

The first specimen currently referred to the genus *Indarctos* consists of a left mandible fragment from the Greek locality of Pikermi identified as *Hyaenarctos* sp. by Dames (1883), with Weithofer (1888) later referring it to *H. atticus*. However, in 1884 Lydekker created the species *H. punjabiensis* based on a broken maxilla (which included P4-M1) from the Siwaliks in India. In that paper he also assigned a maxilla with P4-M2 and a skull with Upper canine and P4-M2 from other localities to the same genus. The latter specimen has been the cause of most of the later taxonomic confusion in comparisons among species since this skull clearly can be assigned to the genus *Agriotherium*, due to the lack of a clear talon and different relative development of the main cusps in the M2.

Pilgrim (1913) moved some of these remains from the junior synonym genus *Hyaenarctos* (also later synonym of *Agriotherium*) to the newly created genus *Indarctos* based on a maxillary fragment with M2 and broken M1 from the Siwa-

liks. However, it was not until a year later (Pilgrim 1913) that he measured, figured and compared this specimen to other, similar fossils, describing *Indarctos salmontanus* as the type species of the genus (Pilgrim, 1914). Later Baryshnikov (2002) accepted only two species of the genus *Indarctos*, literally he stated: “The genus *Indarctos* comprises two species. *I. arctoides* (Depéret, 1895) was found in Europe, northern Africa and Turkey, dating as the Vallesian (MN 9-11). *I. punjabiensis* (Lydekker, 1884) known from the Turolian (MN 10-13) had a Holarctic range, occurring in Europe, northern Africa, in the central, southern and eastern parts of Asia, and in North America”. Therefore, although indirectly, he synonymized Pilgrim’s type species with *Indarctos punjabiensis*.

The oldest record of *Indarctos* comprises some specimens from MN9 in the Vallesian basin of Vallès-Penedès, at the localities of Can Porull (Villalta & Crusafont 1943, 1948) and later from Can Llobateres (Crusafont & Kurtén 1976). These remains were assigned to *Indarctos vireti* Villalta & Crusafont, 1943, a small sized member of the genus, with elongated molars, more primitive (less reduced) premolars, and well-developed diastemas between them.

Indarctos arctoides (Depéret, 1895), is documented from MN10. This species was first described from the French locality of Montredon, and can be found along the northern edge of the Mediterranean Sea, including Spain, France and Turkey (Tobien 1955; Geraads *et al.* 2005; Abella 2011; Abella *et al.* 2014).

Finally, *Indarctos punjabiensis* (incl. *I. atticus*) has an extensive record in the Iberian Peninsula from MN10 to MN12. Most of these specimens were studied in detail by Montoya *et al.* (2001). Following the arguments of that paper we accept three species of *Indarctos*; four, if we take include the peculiar *I. anthracites* (Weithofer, 1889), which is thought to be an insular form and is only represented by a single published specimen. These are *I. vireti*, *I. arctoides* and *I. punjabiensis* with the last species having the temporal range. The present study increases the stratigraphic range of the genus *Indarctos* in the Iberian Peninsula to the lower part of MN13.

MATERIAL AND METHODS

The fossil remains of *Indarctos punjabiensis* from Las Casiones are part of the collections of the Gobierno de Aragón (Spain), and are housed at the Museo Aragonés de Paleontología, Fundación Conjunto Paleontológico de Teruel-Dinópolis (Teruel, Spain). The studied material was: KS-30: right IV metacarpal; K-162: left M1; K-161: distal fragment of right M1; KS-4315: right M2; KS-4316: left M1; KS-4317: left M2; K94-1733: right P4; K-174: maxillary fragment (P2, root of P1 and P3); KS-4314: premolar (p3?) (Figs 2; 3). Of the fossil remains studied here, only M1 K-161 and K-162 were originally described in Alcalá (1994). For comparison, we studied the the upper dentition of *Indarctos vireti* housed at the Institut Català de Paleontologia- Miquel Crusafont Museu (since the holotype consists of a mandible fragment with p4-m2), cast specimens of *Indarctos arctoides* (paratype) from Montredon housed in the Natural History Museum Basel – Naturhistorisches Museum




FIG. 2. — *Indarctos punjabiensis* (Lydekker, 1884) from Las Casiones: **A-C**, K94-1733, right P4: **A**, lingual view; **B**, labial view; **C**, occlusal view; **D**, K-162, left M1, occlusal view; **E**, KS-4316, left M1, occlusal view; **F**, KS-4317, left M2, occlusal view; **G**, K-161, distal fragment of right M1, occlusal view; **H**, KS-4315, right M2, occlusal view; **I**, K-174, left maxillary fragment with P2 and root of P1 and P3 (occlusal view); **J**, lower premolar, labial view. Scale bar: 5 cm.

Basel, the sample of *I. arctoides* housed at Museo Nacional de Ciencias Naturales-CSIC in Madrid, and the cast of the holotype of *I. punjabiensis* housed at the British Museum of Natural History, London.

The measurements were taken using Mitutoyo Absolute digital calipers to the nearest 0.1 mm (Table 1).

INSTITUTIONAL ABBREVIATIONS

KS and K Las Casiones;
MNCN Museo Nacional de Ciencias Naturales;
NMB Natural History Museum Basel – Naturhistorisches

ICP

Museum Basel;
Institut Català de Paleontologia – Miquel Crusafont
Museu.

SYSTEMATIC PALEONTOLOGY

Order CARNIVORA Bowdich, 1821
Suborder CANIFORMIA Kretzoi, 1943
Infraorder ARCTOIDEA Flower, 1869
Parvorder URSIDA Tedford, 1976
Superfamily URSOIDEA Fischer von Waldheim, 1817



FIG. 3. — *Indarctos punjabiensis* (Lydekker, 1884) from Las Casiones, right IV metacarpal: **A**, plantar view; **B**, medial view; **C**, dorsal view; **D**, lateral view. Scale bar: 5 cm.

Family URSIDAE Fischer von Waldheim, 1817
Subfamily AILUROPODINAE Grevé, 1894
Tribe INDARCTINI Abella, Alba, Robles, Valenciano,
Rotgers, Carmona, Montoya & Morales, 2012
Genus *Indarctos* Pilgrim, 1913

Indarctos punjabiensis (Lydekker, 1884)

Hyaenarctos punjabiensis Lydekker, 1884: 226.

Agriotherium cf. *A. robles* – Alcalá 1994: 103, 104, pl. 3, fig. o.

SYNTYPE. — Fragments of the left and right maxillae with P4-M1, figured by Lydekker (1884: pl. XXX, fig. 2).

TYPE LOCALITY. — Hasnot Siwaliks. Salt Range, Jhelum District. Punjab.

AGE. — Late Miocene. MN 10-13.

EMENDED DIAGNOSIS. — Medium to large sized Ursidae, with marked sexual dimorphism. Plantigrade. Dolichocephalic skull, base of the zygomatic arch coincides with the M1 and the mesial part of M2. The premolars are smaller in relative size and number compared to the other members of the genus, being crowded to some degree. Both the P2 and the P3 are double rooted, with the tendency to be fused in some individuals. The mesio-distal length of the P4 is equal to that of M1, and it has a quite developed parastyle, with

two crists, one lingual and one labial. Wide upper molars, almost as wide as they are long; from sub-rectangular to trapezoid in shape. The talon in M2 is large, but not as long as in the derived Ursinae. Generally low, but robust jaw. Relatively long m1, with equally long talonid and trigonid; tall entoconid, taller than the hypoconid; m1 talonid similar to that of m2. The m3 morphology varies from rounded to elongated.

DESCRIPTION

Upper dentition

The description has been carried out on the possibly associated upper dentition, using both sides when possible.

P1 (Fig. 2I). Only a single root is preserved. Smallest of the upper premolars.

P2 (Fig. 2I). Reduced, ovoid premolar, with only one root. Single cusp with two ridges, one mesial and one distal, that curve towards the lingual side; the distal cusp is blunter than the mesial one. This tooth is rotated relative to the main axis of the maxilla and somehow crowded, due to the shortening of the muzzle.

P3 (Fig. 2I). Only a single root is preserved. Larger than the preceding upper premolars.




Fig. 4. — Bivariate plot of the P4 of different *Indarctos* Pilgrim, 1913 species from various sites.




Fig. 5. — Bivariate plot of the M1 of different *Indarctos* Pilgrim, 1913 species from various sites.




FIG. 6. — Bivariate plot of the M2 of different *Indarctos* Pilgrim, 1913 species from various sites.

P4 (Fig. 2A-C). Complete upper carnassial. Deeply wrinkled enamel. Well-developed parastyle with two ridges, one labial and one lingual. High paracone, almost flat in its lingual part but convex in the labial one. Metastyle lower than paracone, but more sectorial; wide at its base, with two cingula, a weak one in the labial part and a slightly more developed one in the lingual part. Where the two cingula join they create a distal pointed edge. The protocone is complex. It has three cusps, which grow taller towards its distal part, with the latter being the tallest, practically at the level of the notch between paracone and metastyle, and the only one partly surrounded by the labial cingulum of the cusp. The protocone occupies an intermediate position with respect to the paracone, and is projected lingually.

M1 (Fig. 2D). Very worn first upper molar, much more in the distal than in the mesial part. This kind of asymmetric wear is quite common in ursoids, and in this case prevents the description of many of the dental features. It is nearly square in shape. The contact facet for P4 is relatively large

and concave instead of flat, while that for the M2 is located almost at the labio-distal corner of the tooth. Towards the labial side, between the paracone and the metacone, there is an expansion zone that widens towards the metacone; this structure has a slight vertical wear facet.

M1 (Fig. 2E-G). Molars with a clear quadrangular shape, almost as wide as long. The enamel has the typical young ursid roughness. It is barely worn and the roots were not completely formed before death and are not recovered. As in the worn M1, these two also have a medial-lingual thickening of the labial cingulum between the paracone and metacone. The paracone is larger (in both width and height) than the metacone. Both parastyle and metastyle are present, although these are very weak compared to the main cusps. There is a well-developed cingulum occupying almost all of the lingual face. The contact with P4 is vertical and straight. The protocone is large and cristiform. Distal to the protocone there is a robust hypocone, which is taller than the protocone, and although being the small-




FIG. 7. — Other *Indarctos* species for comparison with *I. punjabiensis* (Lydekker, 1884) of Las Casiones: **A**, *I. vireti* Villalta & Crusafont, 1943; **B**, *I. arctoides* (Depéret, 1895) (from type locality); **C**, *I. punjabiensis* (syntype). All images are in occlusal view. Scale bar: 5 cm.

est of the four main cusps. Hypocone is more complex, having a ridge on each side. The mesial ridge is close to the protocone, whereas the lingual one joins the lingual cingulum. The medial one meets the ridge coming down the central valley of the tooth from the metacone and the distal ridge ends at the most distal edge of the tooth. In the labial side there is other cingulum, which is considerably thickened in the distal region.

M2 (Fig. 2F, H). A wide tooth, with large cingula on both in its labial and its lingual side, with the latter the most developed. In occlusal view the talon is slightly rotated labially while the labial cingulum is labially tilted, giving the teeth two different crushing surfaces. The tooth enamel is clearly rough, which is most evident on the lingual surface. The protocone is continued by the ridge that originates at the front of the paracone, continuing throughout the mesial part of the molar towards the lingual side, but not forming a clear parastyle. Unlike M1, the M2 protocone is slightly bilobed in its central part. The paracone is taller than the metacone although the difference is slightly smaller than in the M1. The talon is very rounded and relatively short, and although it joins the metacone, it has a more labial position. It has a ridge that runs from the hypocone

to the metacone, forming a distal semicircle divided into at least six small cusps.

Lower Dentition

px (Fig. 2J). Biradicular premolar. Longer than wide. It has only one cusp in the center of the crown, with two cristids, one slightly mesio-lingual, which approaches the mesial edge, and other one distal.

Postcranial skeleton

Right Mc IV (Fig. 3). This metapodial has a robust general morphology. Epiphyses are wider than the diaphysis, the latter having a homogeneous width in dorsal view throughout its length. However, in lateral view, there is a difference in height because the distal part of the diaphysis is clearly flatter. The proximal epiphysis is robust, convex in the most proximal zone, the surface that serves as articulation for the unciform. Its contact facet for the Mc III has two lobes, a convex dorso-proximal one and a flat palmar-proximal one. The articulation for the Mc V has also two facets, a slightly convex dorsal one and a noticeably concave palmar one. The distal epiphysis is slightly wider and somewhat rounded in its dorsal view, thus being more flattened in lateral view.

SYSTEMATIC DISCUSSION

The fossil remains from Las Casiones show the following features in the dentition that clearly match the morphology of *Indarctos*:

- 1) A two-ridged parastyle on the P4. A mesially complex protocone;
- 2) Complex hypocone on M1;
- 3) Relatively short and robust Mc IV, with relatively large proximal and distal epiphyses that are equal in size. Medial and lateral contact facets of proximal epiphysis separate and distinct from proximal contact facet.

The fossil remains from Las Casiones are amongst the largest of the genus (Figs 4, 5 and 6), and in the Iberian Peninsula second only to the huge remains from Concud. Therefore, they are easily included in *Indarctos punjabiensis*, a species that was probably present in the Iberian Peninsula from MN10 and certainly from MN11: Masia la Roma 604, Terrassa, Crevillente 2, Concud, Rambla de Valdecebro 5 (Montoya *et al.* 2001; Alcalá *et al.* 2005; Abella *et al.* 2012, 2014).

COMPARISON WITH OTHER *INDARCTOS* SPECIES

Indarctos punjabiensis exhibits considerable morphological variability in the dentition, even within samples from the same site (Montoya *et al.* 2001), as shown by the measurements in Table 1 and the size comparison in Figures 4, 5, and 6. We can, however, distinguish it from other species of the genus on to the criteria presented below.

The remains from Las Casiones differ from *Indarctos vireti* (Fig. 7) in the following features (the holotype of this species is a mandible (Can Purull, Villadecavalls, Spain) so the comparison has been carried out using the upper dentition from Can Llobateres):

Overall, the dentition from Las Casiones is much larger than that of *Indarctos vireti*; the P4 of the latter is more elongated, without parastyle and much simpler in terms of the number of cusps in the protocone; the M1, although similar in cusp disposition, is more elongated in the Vallesian species, especially in the talon; the M2 is much more elongated in *I. vireti* than in the Las Casiones specimen.

The fossil remains from Las Casiones differ from the holotype of *Indarctos arctoides* (Montredon, France, MN10), and from the spectacular sample of *Indarctos arctoides* from Batallones 3 (Madrid, Spain) (Fig. 7) in the following features:

The dentition from Las Casiones is larger than that of *Indarctos arctoides*; the P4 of *I. arctoides* (same locality as the holotype) is already as wide as that of Las Casiones or relatively wider, due to the lingual widening of the wall between the paracone and metastyle; the parastyle is not as large and the protocone has fewer cusps, with only two cusps instead of the three in the P4 from Las Casiones; the M1 of both species are almost identical in shape, but the M2 from Las Casiones seems relatively much wider than that of the holotype of *I. arctoides*.

The dentition from Las Casiones is almost identical in size as that of the holotype of *Indarctos punjabiensis* (Hasnot Siwaliks, India. MN10-13) (Fig. 7). Both the P4 and M1 are very

similar in morphology, though the paracone of P4 has fewer cusps in the holotype of *I. punjabiensis*. Comparison of the M2 from Las Casiones to that published by Pilgrim in 1914 under the name of *I. salmontanus* (now *I. punjabiensis*) leads to the conclusion that the latter is slightly more elongated, due to the lesser development of the labial and lingual cingula, but it has almost the same cusp disposition.

BIOSTRATIGRAPHY

As already mentioned, the fossil record of the genus *Indarctos* in the Iberian Peninsula covers much of the late Miocene. Thus, it is represented by *I. vireti* in the Vallesian (MN9 and MN10) of the Vallès-Penedès (Catalonia), by *I. arctoides* in the Vallesian (MN10) of the Batallones sites (Madrid) and by *I. punjabiensis* (formerly *I. atticus*) in the late Vallesian of Terrassa (Vallès-Penedès, Catalonia) and Masía de La Roma 604 (Teruel basin). It is also recorded from the early Turolian (MN11) of Crevillente 2 (Alicante), as well as in other Turolian (MN11 and MN12) localities from the Teruel basin. In addition, the range is now extended to the late Turolian or Ventian (MN13) also from the Teruel basin. More detailed information on the Spanish record of the genus *Indarctos* can be found in Montoya (1994), Montoya *et al.* (2001), Alcalá *et al.* (2005), Abella (2011) and Abella *et al.* (2013, 2014, 2015).

From a biostratigraphic point of view, the material of *Indarctos punjabiensis* in Las Casiones represents the youngest appearance of the genus in the fossil record of the Iberian Peninsula (and perhaps also of Europe), just before its replacement by the hemicyonid *Agriotherium*. Should be mentioned that the *Indarctos* record from Ananiev (Ukraine) could be more or less of the same age (Baryshnikov 2002). Recent works allow us to date, with relatively good precision, the moment of this substitution and its relationship to the faunistic events that took place during the Messinian, although prior to the Messinian Salinity Crisis.

The Las Casiones site lies within MN13, in local zone M2 of van Dam (1997), and van Dam *et al.* (2006: suppl. notes), where it is assigned an age *c.* 6.3 Ma. In addition, it is the reference site for the so-called First African-Iberian Dispersal (Gibert *et al.* 2013) because it records the FAD of Hippopotamidae in Western Europe from Africa. Furthermore, the entry of eastern immigrants, such as the genus *Plesiogulo* and *Hystrix depereti*, takes place during this period.

On the other hand, the FAD of the genus *Agriotherium* in Europe is recorded at the Venta del Moro site (Cuenca del Cabriel, Valencia) (Morales 1984; Montoya *et al.* 2006), with *A. robusti* Morales & Aguirre, 1976. This deposit is located within MN13 in local zone M3, and a recent paleomagnetic study has allowed dating it to *c.* 6.23 Ma (Gibert *et al.* 2013). Because Venta del Moro also records the FAD in Europe of *Paraethomys* and *Paracamelus*, taxa most likely coming from North Africa, it is the reference site for the so-called Second African-Iberian Dispersal (Gibert *et al.* 2013). This dispersal coincides with the so-called 'Paraethomys event' or 'Second Messinian mammalian event' of Agustí *et al.* (2006) and with




FIG. 8. — Biochronology showing the stratigraphic ranges of the genera *Indarctos* Pilgrim, 1913 and *Agriotherium* A. Wagner, 1837 in the Iberian Peninsula, related to the two biotic events recorded during the Messinian. Mammal ages and local zones according to Morales *et al.* 2013. Absolute age after Gibert *et al.* 2013.

the ‘Third dispersal event’ by Made *et al.* (2006). The fact that this faunal event has been evidenced, and named differently by different authors tells us of its importance.

The mammalian fauna with *Agriotherium* from Venta del Moro is, as far as large mammals are concerned, composed of a mixture of taxa already present in the Western European Turolian, together with a contingent of more than 50% that appear first in the fossil record of the Iberian Peninsula (Morales *et al.* 2011), and who are immigrants from the east or from Africa. Thus, the Euroasiatic provenance of *A. roblesii* cannot be disregarded. Therefore, the replacement of *Indarctos* by *Agriotherium* could be linked to this great faunal renewal event, which we now know took place 250 000 years before the onset of the Messinian Salinity Crisis (Agustí *et al.* 2006; Gibert *et al.* 2013) (Fig. 8).

FINAL REMARKS

Although some of the dental features may converge towards those present in some species of *Agriotherium* species, it seems clear that the studied dental remains represent a large-sized *Indarctos*, similar in both size and morphology to the syntype of *I. punjabiensis*. Moreover, the population from Las Casiones

marks the upper range of the record of the genus *Indarctos* in the Iberian Peninsula and probably also in Europe. It belongs to the faunal assemblage prior to the entrance, at c. 6.23 Ma, of the rodent *Paraethomys* and many other immigrants, including the hemicyonid ursoid *Agriotherium*.

Acknowledgements

The authors of this paper would like to express our deep sadness at the passing of our beloved friend and colleague Stéphane, with whom we shared great moments, both during fieldwork and manuscript writing; we would have loved to have had many more moments together. Excavations at Las Casiones, preparation of fossils, and palaeontological research have been made possible thanks to the support of the Dirección General de Patrimonio Cultural, Gobierno de Aragón (Spain). The authors thank the Herrero family (owners of the property where the Las Casiones site is located), J. Esteban, M. Marco, and all members and volunteers from excavation and fossil preparation teams. This study is one of the palaeontological research projects subsidized by the Departamento de Educación, Cultura y Deporte-Gobierno de Aragón and Grupo de Investigación de Referencia E04_17R FOCONTUR cofinanciado por el Departamento de Innovación, Investigación y Universidad

(Gobierno de Aragón) y fondos FEDER Aragón 2014-2020, “Construyendo Europa desde Aragón”; the Instituto Aragonés de Fomento. DH is researcher in formation in the FPU 2013 program granted by the Ministerio de Educación, Cultura y Deporte, Spain). The support of the DST-NFR Centre of Excellence in Palaeosciences (CoE-Pal) toward this research for A.V. (COE2018-09POST and COE2019-PD07) is hereby acknowledged. Opinions expressed and conclusions arrived at, are those of the authors and are not necessarily to be attributed to the CoE. This study was supported by the Spanish Research Projects CGL 2015-68333-P and CGL2016-76431-P (AEI/FEDER, UE), the BSCH-UCM910607 of the Universidad Complutense de Madrid, the Research Groups CSIC 64 1538 and CAM-UCM 910607, and the Generalitat de Catalunya (CERCA Programme, and Beatriu de Pinós contract 2017 BP 00223 from AGAUR to JA).

REFERENCES

- ABELLA J. 2011. — *Indarctos arctoides* Depéret, 1895 (Carnivora, Mammalia) del yacimiento vallesiense de Batallones 3 (cuenca de Madrid) [*Indarctos arctoides* Depéret, 1895 (Carnivora, Mammalia) from the vallesian locality of Batallones 3 (Madrid Basin)]. PhD dissertation, unpublished. Madrid: Universidad Autónoma de Madrid- Museo Nacional de Ciencias Naturales (CSIC) (in Spanish).
- ABELLA J., ALBA D. V., ROBLES J. M., VALENCIANO A., ROTGERS C., CARMONA R., MONTOYA P. & MORALES J. 2012. — *Kretzoiarctos* gen. nov., the oldest member of the giant panda clade. *PLoS ONE* 7: e48985. <https://doi.org/10.1371/journal.pone.0048985>
- ABELLA J., VALENCIANO A., PÉREZ-RAMOS A., MONTOYA P. & MORALES J. 2013. — On the socio-sexual behaviour of the extinct ursid *Indarctos arctoides*: An approach based on its baculum size and morphology. *PLoS ONE* 8: e77711. <https://doi.org/10.1371/journal.pone.0073711>
- ABELLA J., MONTOYA P. & MORALES J. 2014. — Paleodiversity of the Superfamily Ursoidea (Carnivora, Mammalia) in the Spanish Neogene, related to environmental changes. *Journal of Iberian Geology* 40:11-18. https://doi.org/10.5209/rev_JIGE.2014.v40.n1.44083
- ABELLA J., PÉREZ-RAMOS A., VALENCIANO A., ALBA D. M., ERCOLI M. D., HONTECILLAS D., MONTOYA P. & MORALES J. 2015. — Tracing the origin of the panda's thumb. *The science of Nature* 102: 35. <https://doi.org/10.1007/s00114-015-1286-3>
- AGUSTÍ J., GARCÉS M. & KRIJGSMA W. 2006. — Evidence for African-Iberian exchanges during the Messinian in the Spanish mammalian record. *Palaeogeography, Palaeoclimatology, Palaeoecology* 238: 5-14. <https://doi.org/10.1016/j.palaeo.2006.03.013>
- ALCALÁ L. 1994. — Macromamíferos neógenos de la fosa de Alfambra-Teruel [Neogene macromammals from the Alfambra-Teruel grave]. Instituto de Estudios Turolenses, Museo Nacional de Ciencias Naturales – CSIC, Teruel, 554 p. (in Spanish).
- ALCALÁ L. & MONTOYA P. 1998a. — *Hystrix primigenia* (Wagner, 1848) (Rodentia, Mammalia) del Turolense superior (MN 13) de Las Casiones (Teruel, España) [*Hystrix primigenia* (Wagner, 1848) (Rodentia, Mammalia) from the Late Turolian (MN 13) of Las Casiones (Teruel, Spain)]. *Revista Española de Paleontología* 13: 139-147.
- ALCALÁ L. & MONTOYA P. 1998b. — *Hexaprotodon crusafonti* (Hippopotamidae, Mammalia) del Mioceno superior de El Arquillo (Fosa de Teruel) [*Hexaprotodon crusafonti* (Hippopotamidae, Mammalia) from the Late Miocene of El Arquillo (Teruel Grave)]. *Boletín de la Real Sociedad Española de Historia Natural Sección Geológica* 94: 93-100.
- ALCALÁ L., MONTOYA P. & MORALES J. 1994. — New large mustelids from the Late Miocene of the Teruel Basin (Spain). *Comptes rendus de l'Académie des Sciences, série II*, 319: 1093-1100. <https://gallica.bnf.fr/ark:/12148/bpt6k58032150/f233.item>
- ALCALÁ L., VAN DAM J., LUQUE L., MONTOYA P. & ABELLA J. 2005. — Nuevos mamíferos vallesienses en Masía de La Roma (Cuenca de Teruel) [New Vallesian mammals from Masía de La Roma (Teruel Basin)]. *Geogaceta* 37: 199-202.
- BARYSHNIKOV G. F. 2002. — Late Miocene *Indarctos punjabiensis* atticus (Carnivora, Ursidae) in Ukraine with survey of *Indarctos* records from the former USSR. *Russian Journal of Theriology* 1: 83-89. <https://doi.org/10.15298/rusjtheriol.01.2.02>
- BARYSHNIKOV G. F. & TLEUBERDINA P. A. 2017. — Late Miocene *Indarctos* (Carnivora: Ursidae) from the Karabulak Formation of the Kalmakpai river (Zaisan Depression, Eastern Kazakhstan). *Proceedings of the Zoological Institute of the Russian Academy of Sciences* 321: 3-9.
- BONIS L. DE, ABELLA J., MERCERON G. & BEGUN D. R. 2017. — A new late Miocene ailuropodine (Giant Panda) from Rudabanya (North-central Hungary). *Geobios* 50: 413-421. <https://doi.org/10.1016/j.geobios.2017.09.003>
- BOWDICH T. E. 1821. — *An Analysis of the Natural Classifications of Mammalia for Use of Students and Travellers*. J. Smith, Paris, 115 p.
- BRUIJN H. DE, DAAMS R., DAXNER-HÖCK G., FAHLBUSCH V., GINSBURG L., MEIN P. & MORALES J. 1992. — Report of the RCMNS working group on fossil mammals, Reimsburg 1990. *Newsletters on Stratigraphy* 26: 65-118. <https://doi.org/10.1127/nos/26/1992/65>
- CERDEÑO E. 1989. — *Revisión de la sistemática de los rinocerontes del Neógeno de España* [Review of the Systematic of Rhinoceros from the Neogene of Spain]. [PhD dissertation]. Universidad Complutense, Madrid, 429 p.
- CRUSAFONT PAIRÓ M. & KURTÉN B. 1976. — Bears and bear-dogs from the Vallesian of the Vallés-Penedés Basin, Spain. *Acta Zoologica Fennica* 144: 1-29.
- DAMES W. 1883. — Über das Vorkommen von *Hyaenarctos* in den Pliocän-Ablagerungen von Pikermi bei Athen. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin* 1883 (8): 132-139. <https://biodiversitylibrary.org/page/8789424>
- DEPÉRET C. 1895. — Résultats des fouilles paléontologiques dans le Miocène supérieur de la colline de Montredon. *Comptes rendus hebdomadaires des Séances de l'Académie des Sciences* 121: 432-434. <https://biodiversitylibrary.org/page/36256646>
- DOMINGO M. S., DOMINGO L., ABELLA J., VALENCIANO A., BADGLEY C. & MORALES J. 2016. — Feeding ecology and habitat preferences of top predators from two Miocene carnivore-rich assemblages. *Paleobiology* 42: 489-507. <https://doi.org/10.1017/pab.2015.50>
- FISCHER [DE WALDHEIM] G. 1814. — *Zoognosia tabulis synopticis illustrata*. Vol. 3, 3rd ed. Nikolai Sergei Vsevolozsky, Mosquae, 732 p.
- FLOWER W. H. 1869. — On the value of the characters of the base of the cranium in the classification of the order Carnivora, and on the systematic position of Bassaris and other disputed forms. *Proceedings of the Zoological Society of London* 1869: 4-37. <https://biodiversitylibrary.org/page/28662875>
- FRICK C. 1926. — The Hemicyoninae and an American Tertiary Bear. *Bulletin of the American Museum of Natural History* 56: 1-120. <http://hdl.handle.net/2246/1321>
- GERAADS D., KAYA T. & MAYDA S. 2005. — Late Miocene large mammals from Yulafli, Thrace region, Turkey, and their biogeographic implications. *Acta Palaeontologica Polonica* 50 (3): 523-544.
- GIBERT L., SCOTT G. R., MONTOYA P., RUIZ-SÁNCHEZ F. J., MORALES J., LUQUE L., ABELLA J. & LERÍA M. 2013. — Evidence for an African-Iberian mammal dispersal during the pre-evaporitic Messinian. *Geology* 41: 691-694. <https://doi.org/10.1130/G34164.1>
- HENDEY Q. B. 1972. — A Pliocene ursid from South Africa. *Annals of the South African Museum* 59: 115-132. <https://biodiversitylibrary.org/page/40938928>

- HENDEY Q. B. 1980. — *Agriotherium* (Mammalia, Ursidae) from Langebaanweg, South Africa, and relationships of the genus. *Annals of the South African Museum* 81: 1-109. <https://biodiversitylibrary.org/page/40751715>
- KRETZOI M. 1943. — *Kochictis centennii* n. g. n. sp., ein altertümlicher Creodonte aus dem Oberoligozän Siebenbürgens. *Földtani Közlöny* 73: 190-195.
- LYDEKKER R. 1884. — Indian Tertiary and post-Tertiary Vertebrata. Part 6. Siwalik and Narbada Carnivora. *Memoirs of the Geological Survey of India. Palaeontologia Indica* 10: 178-363. <https://biodiversitylibrary.org/page/53674503>
- MADE J. VAN DER, MORALES J. & MONTOYA P. 2006. — Late Miocene turnover in the Spanish mammal record in relation to palaeoclimate and the Messinian Salinity Crisis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 238: 228-246. <https://doi.org/10.1016/j.palaeo.2006.03.030>
- MEIN P. 1990. — Updating of MN zones, in LINDSAY E. H., FAHLBUSCH V. & MEIN P. (eds), *European Neogene Mammal Chronology*. Plenum Press, New York: 73-90. https://doi.org/10.1007/978-1-4899-2513-8_6
- MONESCILLO M. F. G., SALESA M. J., ANTÓN M., SILICEO G. & MORALES J. 2014. — *Machairodus aphanistus* (Felidae, Machairodontinae, Homotherini) from the Late Miocene (Vallesian, MN10) site of Batallones-3 (Torrejón de Velasco, Madrid, Spain). *Journal of Vertebrate Paleontology* 34: 699-709. <https://doi.org/10.1080/02724634.2013.804415>
- MONTOYA P. 1994. — Los macromamíferos del Mioceno superior del área de Crevillente (Alicante) [The macromammals from Late Miocene of the area of Crevillente (Alicante)]. [PhD dissertation] Valencia: Universitat de València (in Spanish).
- MONTOYA P., ALCALÁ L. & MORALES J. 2001. — *Indarctos* (Ursidae, Mammalia) from the Spanish Turolian (Upper Miocene). *Scripta Geologica* 122:123-151.
- MONTOYA P., MORALES J., ROBLES F., ABELLA J., BENAVENT J. V., MARÍN M. D. & RUIZ SÁNCHEZ F. J. 2006. — Las nuevas excavaciones (1995-2006) en el yacimiento del Mioceno final de Venta del Moro, Valencia [The new excavations (1995-2006) in the Late Miocene locality of Venta del Moro]. *Estudios Geológicos* 62: 313-326 (in Spanish). <https://doi.org/10.3989/egol.0662128>
- MORALES J. 1984. — *Venta del Moro: su macrofauna de mamíferos y biostratigrafía continental del Mioceno terminal mediterráneo*. Editorial Universidad Complutense, Madrid, 340 p (in Spanish).
- MORALES J. & AGUIRRE E. 1976. — Carnívoros de Venta del Moro (Valencia) [Carnivorans from Venta del Moro (Valencia)]. *Trabajos sobre Neógeno-Cuaternario* 5: 31-74 (in Spanish).
- MORALES J., MONTOYA P. & ABELLA J. 2011. — Venta del Moro: un yacimiento clave para conocer la historia del límite Mio-Plioceno en el área mediterránea [Venta del Moro: a key locality to know the history of the Mio-Pliocene limit in the Mediterranean area]. *Isurus* 4: 60-71 (in Spanish).
- MORALES J., PELÁEZ-CAMPOMANES P., ABELLA J., MONTOYA P., GIBERT L., SCOTT G., CANTALAPIEDRA J. L. & SANISIDRO O. 2013. — The Ventian mammal age (latest Miocene): present state. *Spanish Journal of Palaeontology* 28: 149-160.
- PESQUERO M. D. 2003. — *Hipparion* del Turoliense superior de Las Casiones (Fosa de Teruel) [*Hipparion* from Late Turolian from Las Casiones (Teruel Grave)]. *Coloquios de Paleontología* Vol. Ext. 1: 511-548 (in Spanish).
- PILGRIM G. E. 1913. — The correlation of the Siwaliks with the Mammal Horizons of Europe. *Records of the Geological Survey of India* 43: 264-326.
- PILGRIM G. E. 1914. — Further description of *Indarctos salmantanus* Pilgrim, the new genus of bear from the Middle Siwaliks, with some remarks on the fossil Indian Ursidae. *Records of the Geological Survey of India* 44: 225-233.
- QIU Z.-X., DENG T. & WANG B.-Y. 2014. — A Late Miocene *Ursavus* skull from Guanghe, Gansu, China. *Vertebrata Palasiatica* 52: 265-302.
- SALESA M. J., PESQUERO M. D., SILICEO G., ANTÓN M., ALCALÁ L. & MORALES J. 2012. — A rich community of felidae (Mammalia, Carnivora) from the late Miocene (Turolian, MN 13) site of Las Casiones (Villalba Baja, Teruel, Spain). *Journal of Vertebrate Paleontology* 32: 658-676. <https://doi.org/10.1080/02724634.2012.649816>
- TEDFORD R. H. 1976. — Relationship of pinnipeds to other carnivores (Mammalia). *Systematic Zoology* 25: 363-374. <https://doi.org/10.2307/2412511>
- TOBIEN H. 1955. — Neue und wenig bekannte Carnivoren aus den Unterpliozänen Dinotheriensanden Rheinheßens. *Notizblatt des Hessischen L.-Amtes für Bodenforschung zu Wiesbaden* 83: 7-31.
- VAN DAM J. A. 1997. — The small mammals from the Upper Miocene of the Teruel-Alfambra region (Spain): paleobiology and paleoclimatic reconstructions. *Geologica Ultraiectina* 156: 1-204.
- VAN DAM J. A., ALCALÁ L., ALONSO ZARZA A., CALVO J. P., GARCÉS M. & KRIJGSMAN W. 2001. — The upper Miocene mammal record from the Teruel-Alfambra region (Spain). The MN system and continental stage/age concepts discussed. *Journal of Vertebrate Paleontology* 21: 367-385. [https://doi.org/10.1671/0272-4634\(2001\)021\[0367:TUMMRF\]2.0.CO;2](https://doi.org/10.1671/0272-4634(2001)021[0367:TUMMRF]2.0.CO;2)
- VAN DAM J. A., ABDUL AZIZ H., ÁLVAREZ-SIERRA M. A., HILGEN F. J., VAN DEN HOEK OSTENDE L., LOURENS L. J., MEIN P., VAN DER MEULEN A. J. & PELÁEZ-CAMPOMANES P. 2006. — Long-period astronomical forcing of mammal turnover. *Nature* 443: 687-691. <https://doi.org/10.1038/nature05163>
- VAN WEERS D. J. & ROOK L. 2003. — Turolian and Ruscinian porcupines (genus *Hystrix*, Rodentia) from Europe, Asia and North Africa. *Paläontologische Zeitschrift* 77: 95-113. <https://doi.org/10.1007/BF03004562>
- VILLALTA Y COMELLA J. F. DE & CRUSAFONT PAIRÓ M. 1943. — *Indarctos vireti*, nova sp., un nuevo úrsido del grupo de los Hemiciónidos del Mioceno superior del Vallés-Penedés [*Indarctos vireti*, nova sp., a new ursid of the hemicyonids group from Late Miocene from Vallés-Penedés]. *Consejo Superior de Investigaciones Científicas* 1943: 45-62 (in Spanish).
- VILLALTA Y COMELLA J. F. DE & CRUSAFONT PAIRÓ M. 1948. — Nuevas aportaciones al conocimiento de los carnívoros pontienses del Vallés-Penedés [New data to the knowledge of the pontian carnivorans from Vallés-Penedés]. *Publicaciones del Instituto de Geología de Barcelona, Miscelánea Almera* 7: 81-121 (in Spanish).
- VIRANTA S. 2004. — Habitat preferences of European Middle Miocene omnivorous ursids. *Acta Palaeontologica Polonica* 49: 325-327.
- WAGNER A. 1837. — First Part of the nineteenth Volume of Asiatic Researches; or Transactions of the Society, instituted in Bengal, for enquiring into the History, etc. [book review]. *Gelehrte Anzeigen* 5 (170): 334-336.
- WEITHOFER A. 1888. — Beiträge zur Kenntniss der Fauna von Pikermi bei Athen. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients* 6: 225-292.
- ZDANSKY O. 1924. — Jungtertiäre Carnivoren Chinas. *Palaeontologia Sinica*, ser. C, 2: 1-155.

Submitted on 1 January 2019;

accepted on 3 June 2019;

published on 2 December 2019.