

Systematics, palaeoecology and taphonomy of Turonian oysters from the northern Gabon Coastal Basin

Benjamin MUSAVU MOUSSAVOU

Université des Sciences et Techniques de Masuku et URESTE, Faculté des Sciences,
Département de Géologie, boîte postale 943, Franceville (Gabon)
musavu@yahoo.fr

Published on 30 June 2017

urn:lsid:zoobank.org:pub:F4D33955-5823-49E9-BC9A-26DFB03038A7

Musavu Moussavou B. 2017. — Systematics, palaeoecology and taphonomy of Turonian oysters from the northern Gabon Coastal Basin. *Geodiversitas* 39 (2): 213-224. <https://doi.org/10.5252/g2017n2a3>

ABSTRACT

The systematic, paleoecology and taphonomy of the Turonian (Azilé Formation) oysters fauna from the geological section PK 12, South of Libreville – Gabon, are carried out. Twelve species from seven genera are described. Genera *Curvostrea* Vyalov, 1936 ; *Gryphaeostrea* Conrad, 1865 ; *Gyrostroma* Mirkamalov, 1963 ; *Ilymatogyra* Stenzel, 1971 ; *Ostrea* Linnaeus, 1758 ; *Rhynchostreon* Bayle, 1878 and all species are recorded herein for the time in Gabonese coastal basin. The assemblage is dominated by representatives of the genus *Ostrea* and can be regarded as parautochthonous. The oyster shells are thin and thick with medium to large size. The paleoecological study indicates a nearshore shallow marine environment with high energy, nutrient-rich and well oxygenated water column. Taphonomic observations show that shells of oyster are affected by disarticulation. Bioerosion, traces of predators and encrustation are scarcity or absence. Abrasion, breakage and high fragmentation of shell are generally lacking.

KEY WORDS

Taphonomy,
paleoecology,
systematics,
Cretaceous,
Azilé Formation,
Turonian,
oysters,
Gabon coastal basin.

RÉSUMÉ

Systématique, paléoécologie et taphonomie des huîtres du Turonien du bassin côtier Nord Gabon.

Une étude intégrant la systématique, la taphonomie et la paléoécologie des huîtres du Turonien (Formation de l'Azilé) du Sud de Libreville au Gabon a été menée à partir de la coupe du PK 12. Douze espèces appartenant à sept genres ont été décrites. Les genres *Curvostrea* Vyalov, 1936 ; *Gryphaeostrea* Conrad, 1865 ; *Gyrostroma* Mirkamalov, 1963 ; *Ilymatogyra* Stenzel, 1971 ; *Ostrea* Linnaeus, 1758 ; *Rhynchostreon* Bayle, 1878 et l'ensemble des espèces sont signalés pour la première fois au sein du bassin côtier gabonais. L'assemblage est dominé par les représentants du genre *Ostrea* et est interprété comme étant parautochtone. L'étude paléoécologique indique un environnement marin de faible profondeur à niveau d'énergie élevé, riche en nutriment avec une colonne d'eau bien oxygénée. Les observations taphonomiques montrent que les coquilles des huîtres sont désarticulées. Les traces de prédateurs, les incrustations et les bioérosions sont rares à absentes. Les abrasions, les cassures et des grandes fragmentations des coquilles sont généralement absentes.

MOTS CLÉS

Taphonomie,
paléoécologie,
systématique,
Crétacé,
Formation de l'Azilé,
Turonien,
huître,
bassin côtier du Gabon.

INTRODUCTION

The marine Cretaceous sedimentary deposits exposed in the Gabonese coastal basin exhibit a rich and diversified assemblage of macrofossils which has been previously studied by several authors such as: Kossmat (1893), Lombard (1930), Furon (1931, 1950), Choubert (1935), Darteville & Brébion (1956), Darteville & Freneix (1957), Freneix (1959, 1966), Weydert & Collignon (1981), Meister *et al.* (1996, 2003), Musavu Moussavou *et al.* (2013a, b; 2014a, b), Musavu Moussavou & Mabicka Obame (2015), Musavu Mousavou (2015). Oysters have been mentioned but they have received little attention.

In fact, the systematic descriptions of Gabonese coastal oysters have been carried out only by Lombard (1930), Darteville & Freneix (1957) and Musavu Moussavou *et al.* (2013a, 2014a). There are no known works on their taphonomy and palaeoecology. Therefore the oyster fauna and its ecology are still unknown, despite their presence in the deposits.

This paper is an attempt to improve this situation based on oyster collected from the so-called PK 12 road cut section which located at Libreville region at GPS coordinates 0°24.080'N, 9°31.611'E.

GEOLOGICAL AND GEOGRAPHICAL SETTING

The Libreville region is situated in North-Western of Gabon (Fig. 1). It belongs to the Gabon sedimentary coastal basin, which extends 800 km along the West African Coast between 1°N and 4°S latitude with a surface of 90 000 km² (Teisserenc & Villemin 1990). The age of the deposits in the Libreville region is assigned to the Cenomanian?-Turonian by ammonites (Hourcq & Hausknecht 1954; Meister *et al.* 2003). In the Libreville region, the sedimentary sequence, which belongs to Azilé Formation (Turonian), is mainly composed of limestone, marls and clays with locally inter-bedded sandstone (e.g. Hourcq & Hausknecht 1954; Gérard 1958; Weydert 1981; Mbina MOUNGUENGUI 1998). Also small coral bioherms have been reported within the limestone (Weydert 1981). The deposits contain a varied marine fauna (Lombard 1930; Hourcq & Hausknecht 1954; Darteville & Brébion 1956; Hudeley & Belmonte 1970; Meister *et al.* 1996, 2003; Mbina MOUNGUENGUI 1998; Chevalier *et al.* 2002; Musavu Moussavou *et al.* (2013a, b; 2014a). In total seven species of oysters have been reported from the Mesozoic of the Libreville region (Table 1).

The PK 12 section (Figs 1; 2) studied here is located at 12 Km in east of Libreville town along national road 1. The sedimentary succession of this section ranges from Upper Cenomanian to Lower/Middle Turonian. Cenomanian is represented only by 1 meter of red argillite. In Turonian, we distinguish two sedimentary series separated by one angular unconformity. Series situated below consists of argillaceous siltstone, greyish to purplish-blue argillite and fine to medium sandstone beds. There are mud ball, sand ball, current ripple, trough bedding, channels and flasers. Series situated above

consisting on greyish to purplish-blue argillite, argillaceous siltstone, white-grey carbonated sandstone, white-grey argillaceous sandstone beds.

MATERIAL

Study of oysters was carried out based on a total of 204 specimens collected from PK 12 section. All specimens are collected from above series within the bed 30 (Fig. 2). From the collected specimens, 149 (89 right valves and 60 left valves) are disarticulated and 55 are articulated. In general, all specimens are well-preserved.

The material is deposited at the collection of the Geology Department, Faculty of Sciences Masuku University of Franceville (Gabon), under collection number MDG/LBV/Lm.

SYSTEMATIC PALEONTOLOGY

In the descriptions, the terms small, medium and large are defined by the following size ranges: small, up to 10.0 mm long; medium, 10.1 to 30 mm; large 30.1 mm and over. Follow abbreviations are used LV for Left valve, RV for Right valve. The classification used here is that of Carter *et al.* (2011).

Class BIVALVIA Linnaeus, 1758
Order OSTREIDA Férussac, 1822
Family GRYPHAEIDAE Vyalov, 1936
Genus *Gryphaeostrea* Conrad, 1865

Gryphaeostrea sp.
(Fig. 3D-F)

MATERIAL. — 3 specimens (MDG/LBV/Lm-7.1 to Lm-7.3).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell is large, inequivalve, elongated to oval in outline. LV is strongly convex, inflated. RV is only slightly convex. LV beak is opisthogyral spiral. Anterior wall of LV is spirally curved. Attachment area is small to large, depressed occupies the posterodorsal portion of the left valve. Ornamentation of LV consists of growth lamellae which adjoining the attachment area. RV is flat to slightly convex. Ornamentation consists of dense, fine growth squamae.

REMARK

The northern Gabon Coastal Basin specimens show great similarities with young specimen of *Gryphaeostrea vomer* Stephenson, 1941 from the Upper Cretaceous which illustrated and described by Pugaczewska (1977). It differs from the present material by having larger space between growth lamellae.




FIG. 1. — Location map of Gabon and Libreville (A) and PK 12 section (B).

TABLE 1. — List of oyster fauna of Libreville. After Lombard (1930), Darteville & Freneix (1957), Musavu Moussavou *et al.* (2013a, 2014a) and the present study.

No. Species	Lombard (1930)	Darteville & Freneix (1957)	Musavu Moussavou <i>et al.</i> 2013a	Musavu Moussavou <i>et al.</i> 2014a	This study
1 <i>Curvostrea tevesthensis</i> (Coquand, 1862)	—	—	—	—	x
2 <i>Exogyra olisiponensis</i> Sharpe, 1850	—	x	—	—	—
3 <i>Gryphaeostrea</i> sp.	—	—	—	—	x
4 <i>Gyrostrea delettrei</i> (Coquand, 1862)	—	—	—	—	x
5 <i>Gyrostrea</i> sp. A	—	—	—	—	x
6 <i>Gyrostrea</i> sp. B	—	—	—	—	x
7 <i>Ilymatogyra</i> (<i>Afrogyra</i>) <i>africana</i> (Lamarck, 1801)	—	—	—	—	x
8 <i>Liostrea</i> ? aff. <i>roachensis</i> (Fourtau, 1917)	x	—	—	—	—
9 <i>Liostrea</i> ? <i>cellea</i> (De Stefani, 1913)	—	x	—	—	—
10 <i>Lopha lombardi</i> Darteville & Freneix, 1957	—	x	x	—	—
11 <i>Lopha</i> ? <i>dieneri</i> (Blanckenhorn, 1890)	—	x	—	—	—
12 <i>Ostrea</i> sp. A	—	—	—	—	x
13 <i>Ostrea</i> sp. B	—	—	—	—	x
14 <i>Ostrea</i> sp. C	—	—	—	—	x
15 <i>Ostrea</i> sp. D	—	—	—	—	x
16 <i>Pycnodonta vesicularis</i> (Lamarck, 1806)	—	x	—	—	—
17 <i>Pycnodonte</i> sp.	—	—	—	—	x
18 <i>Rastellum</i> sp.	—	—	x	x	—
19 <i>Rhynchostreon</i> cf. <i>suborbiculatum</i> (Lamarck, 1801)	—	—	—	—	x




FIG. 2. — Lithostratigraphy of the PK 12 section plotted against occurrence of oysters, gastropods and bivalve.

Genus *Ilymatogyra* Stenzel, 1971

Ilymatogyra (*Afrogyra*) *africana* (Lamarck, 1801) (Fig. 3I-M)

Gryphaea africana Lamarck, 1801: 399, pl. 189, figs 5, 6.

Ostrea auressensis Coquand, 1862: 233, pl. 22, figs 12, 13.

Ostrea africana – Coquand 1869: 134, pl. 39, figs 5, 6; pl. 55, figs 10-12.

Exogyra africana – Trevisan 1937: 72, pl. 3, fig. 18; pl. 5, figs 1-4.

Ilymatogyra (*Afrogyra*) *africana* 'forma crassa' – Malchus 1990: 121-124, pl. 7, figs 6, 8, 9, 11, 12, 14-20; pl. 8, figs 1-3.

Ilymatogyra (*Afrogyra*) *africana* – Seeling & Bengtson 1999: 58, fig. 9d-g. — Berndt 2002: 110; pl. 3, figs 5-7. — Ayoub-Hannaa 2011: 84, pl. 7, figs 4-6; text-fig. 3.7.

MATERIAL. — 6 specimens (MDG/LBV/Lm-8.1 to Lm-8.6).

OCCURRENCE. — Cenomanian of North Africa (Coquand 1862; Ayoub-Hannaa 2011), Europe (Trevisan 1937), Brazil (Seeling & Bengtson 1999). In Gabon, this species is recorded for the first time.

DESCRIPTION

Shell is large, inequivalve, elongated to tear-shaped in outline. Umbo is broken. Left valve is convex. This ornamentation consists of strong scaly growth lamellae. Right valve is flat to moderately convex ornament consisting of fine growth line which becoming lamellose towards ventral margin.

REMARK

Malchus (1990) erected two varieties of *Ilymatogyra* (*Afrogyra*) *africana*, 'forma typica' and 'forma crassa'. For difference between these two varieties, the reader is referred to Malchus (1990). All the specimens from the northern Gabon Coastal Basin seem to correspond to 'forma crassa' *sensu* Malchus (1990).

Genus *Rhynchostreon* Bayle, 1878

Rhynchostreon cf. *suborbiculatum* (Lamarck, 1801) (Fig. 3B, C)

Gryphaea cf. *suborbiculata* Lamarck, 1801: 398, figs in Knorr (1801: pl. 62; figs 1, 2).

Rhynchostreon cf. *columbum mermeti* – Freneix 1972: 89, pl. 5, figs 4, 5.

Rhynchostreon cf. *suborbiculatum* – Fischer 1980: 250, pl. 121; figs 7, 8. — Malchus 1990: 132-133. — Videt 2003: 24, pl. 5, figs 1-8.

Rhynchostreon mermeti – Malchus 1990: 128, pl. 8, figs 15-17; pl. 9, figs 5-21.

MATERIAL. — 1 specimen (MDG/LBV/Lm-9).

OCCURRENCE. — Upper Cretaceous of Europe (Fischer 1980; Videt 2003), North Africa (Freneix 1972; Malchus 1990). In Gabon, this species is recorded for the first time.




FIG. 3. — **A**, *Gyrostrea* A MDG/LBV/Lm-13; **B**, **C**, *Rhynchostreon* cf. *suborbiculatum* (Lamarck, 1801) MDG/LBV/Lm-9; **D**–**F**, *Gryphaeostrea* sp.; **D**–**E**, MDG/LBV/Lm-7.1, **F**, MDG/LBV/Lm-7.2; **G**, **H**, *Curvostrea tevesthensis* (Coquand, 1862) MDG/LBV/Lm-11; **I**–**M**, *Ilymatogyra* (*Afrogyra*) *africana* (Lamarck, 1801); **I**, **J**, MDG/LBV/Lm-8.3, **K**, MDG/LBV/Lm-8.2, **L**–**M**, MDG/LBV/Lm-8.1; **N**, **Q**, *Ostreidea* gen. et sp.; indet MDG/LBV/Lm-34; **O**, **P**, *Pycnodonte* sp. MDG/LBV/Lm-10. Scale bar: 1 cm.

DESCRIPTION

Shell, represented only by left valve, is large, inequivalve, elliptical shape in outline. Umbo is small, very opisthogyrate.

Anterior margin is curved into approximately semicircular shape. Posterior margin is slightly curved. Attachment area is small. Ornament consists of ribs in posterodorsal part.

REMARK

This morphotype is doubtfully assigned to *R. suborbiculatum* (Lamarck, 1801) because of the presence of ribs in postero-dorsal part of left valve.

R. suborbiculatum shows great similarities to *R. mermeti* (Coquand, 1862). For discussion concerning this topic, the reader is referred to Freneix (1972), Freneix & Viaud (1986), Malchus (1990), Aqrabawi (1993), Cooper (1995), Seeling & Bengtson (1999). In our material we have only one specimen. For this reason we prefer to consider *R. mermeti* as a synonym *R. suborbiculatum*.

Genus *Pycnodonte* Fischer de Waldheim, 1835

Pycnodonte sp.
(Fig. 3O, P)

MATERIAL. — 1 specimen (MDG/LBV/Lm-10).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell, represented only by one left valve, is large, suborbicular in outline. Umbo is small. Anterior margin is curved into approximately semicircular shape. Posterior margin is straight, curving into ventral margin which is evenly rounded. Attachment area is very large (25 mm in diameter). Surface ornament consists of growth lamellae.

REMARKS

Because of the poor preservation and the small number of specimens, a species designation is not possible; however, the outline and ornamentation seem to show affinity with *Pycnodonta vesicularis* (Lamarck, 1806).

Family FLEMINGOSTREIDAE Stenzel, 1971

Genus *Curvostrea* Vyalov, 1936

Curvostrea tevesthensis (Coquand, 1862)
(Fig. 3G, H)

Ostrea tevesthensis Coquand, 1862: 227, pl. 19, figs 7-13. — Basse 1932: 6.

Liostrea tevesthensis – Pervinquier 1912: 69.

Curvostrea tevesthensis – Malchus 1990: 156. — Cooper 2002: 36, fig-text. 5.

MATERIAL. — 1 specimen (MDG/LBV/Lm-11).

OCCURRENCE. — Cretaceous of North Africa (Pervinquier 1912; Malchus 1990), Southeast Africa (Cooper 2002). In Gabon, this species is recorded for the first time.

DESCRIPTION

Shell is large, inequival, inequilateral, elongated to oval in outline. LV is strongly convex, inflated. RV is only slightly convex. LV beak is opisthogyally spiral. Anterior margin of LV is spirally curved. Its posterior margin is straight, curving into ventral margin which evenly rounded. Attachment area

is small. LV ornamentation consists of commarginal growth striae and rugae separated by wide interspaces. RV is ornamented by concentric fine growth squamae.

REMARK

Curvostrea rouvillei (Coquand, 1862) differs from the present material in being broader and more curved. *Curvostrea bourguignati* (Coquand, 1869) is more quadrate.

Genus *Gyrostrea* Mirkamalov, 1963

Gyrostrea delettrei (Coquand, 1862)
(Fig. 4G-N)

Ostrea delettrei Coquand, 1862: 224, pl. 18, figs 1-7. — Fourtau 1917: 34; pl. 1, figs 1-4.

Exogyra delettrei – Pervinquier 1912: 186, pl. 12, figs 18, ?19. — Trevisan 1937: 74, pl. 5, figs 5-10.

Crassostrea (*Gyrostrea*) *delettrei tarfayensis* – Freneix 1972: 98, pl. 8, figs 5-8; pl. 9, figs 1-7.

Gyrostrea delettrei – El-Sheikh *et al.* 1998: pl. 1, fig. B. — Ayoub-Hannaa 2011: 92, pl. 8, figs 3, 4.

MATERIAL. — 10 specimens (MDG/LBV/Lm-12.1 to Lm-12.10).

OCCURRENCE. — This species has been recorded from Cenomanian to Turonian of North Africa (Pervinquier 1912; Freneix 1972; El-Sheikh *et al.* 1998; Ayoub-Hannaa 2011), Italy (Trevisan 1937). In Gabon this is the first record.

DESCRIPTION

Shell is medium to large, inequivalve, inequilateral, elongated, subtrigonal to elliptical shape in outline. Left and right valves are respectively convex and flat. Umbo is small and opisthogyrate. Anterior margin curved into approximately semicircular shape. Posterior margin is straight or curved. Ventral margin is evenly rounded. Attachment area is rounded and large. LV ornamentation consists of radial and continuous ribs towards umbonal area where are crossing by concentric growth lamellae. Towards ventral margin, ornamentation consists only of widely spaced imbricating concentric growth lamellae separated by wide interspaces. RV is ornamented by concentric fine growth squamae.

REMARK

The specimens of *Gyrostrea delettrei* (Coquand, 1862) from Gabon Coastal Basin show some similarity to *Ostrea roachensis* (Fourtau 1917: 50, pl. 3, fig. 1), but differ in having a variable size shape, size of the attachment area.

Gyrostrea sp. A
(Fig. 3A)

MATERIAL. — 1 specimen (MDG/LBV/Lm-13).

OCCURRENCE. — Turonian of Gabon (this study).




FIG. 4. — **A-D**, *Ostrea* sp. D: **A, B**, MDG/LBV/Lm-33.1; **C, D**, MDG/LBV/Lm-33.2; **E**, *Ostrea* sp. C MDG/LBV/Lm-32.1; **F**, *Gyrostrea* sp. B MDG/LBV/Lm-14; **G-N**, *Gyrostrea delectrei* (Coquand, 1862); **G, H**, MDG/LBV/Lm-12.1; **I, J**, MDG/LBV/Lm-12.2; **K, L**, MDG/LBV/Lm-12.3, M-N. MDG/LBV/Lm-12.4; **O-R**, *Ostrea* sp. B; **O, P**, MDG/LBV/Lm-31.1; **Q, R**, MDG/LBV/Lm-31.2; **S-U**, *Ostrea* sp. A, S-T. MDG/LBV/Lm-15.1; **U**, MDG/LBV/Lm-15.2. Scale bar: 1 cm.

DESCRIPTION

Shell, represented by LV, is medium, inequilateral, elongated to elliptical shape in outline. Anterior and posterior margin are curved giving semicircular shape of shell. At-

tachment area is small. Surface rough with growth squamulae and few rough rounded irregular ribs well separated from each other.

REMARKS

This species differs to *Gyrostrea delettrei* (Coquand, 1862) in having small irregular ribs.

Gyrostrea sp. B
(Fig. 4F)

MATERIAL. — 1 specimen (MDG/LBV/Lm-14).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell, represented only by LV, is large, inequilateral, elongated to oval in outline. Valve is strongly convex, inflated. Beak is opisthogyally spiral. Anterior margin is spirally curved. Its posterior margin is straight, curving into ventral margin which evenly rounded. Attachment area is small. Ornamentation consisting of radial ribs near the umbonal area with concentric growth lamellae which becoming concentric ribs toward central part and margin ventral.

REMARK

Gyrostrea sp. B differs from *Gyrostrea* sp. A in having concentric ribs.

Family OSTREIDAE Rafinesque, 1815
Genus *Ostrea* Linnaeus, 1758

Ostrea sp. A
(Fig. 4S-U)

MATERIAL. — 10 specimens (MDG/LBV/Lm-15.1 to Lm-15.10).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell is large, inequivalve, inequilateral, elongated, subtrigonal to suborbicular in outline. Umbo is small and prominent. Anterior and ventral margin are rounded. Posterior margin straight is angulated toward ventral margin. Attachment area is small. LV is slightly inflated. RV is flat. Ornamentation of LV consisting of numerous radial subacute ribs. This ribs start at the umbonal area, occasionally carrying small spines near ventral margin, and crossed by growth lamellae at irregular intervals. RV is ornamented by concentric fine growth squamae.

REMARK

Ostrea sp. A differs from *Ostrea lateralis* Nilsson, 1827 in having an angulated posterior margin without fold.

Ostrea sp. B
(Fig. 4O-R)

MATERIAL. — 81 specimens (MDG/LBV/Lm-31.1 to Lm-31.81).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell is medium to large, inequivalve, inequilateral, elongated, suborbicular to elliptical shape in outline. LV slightly inflated and RV flat. Umbo is small and opisthogyrate. Ligamental area is narrow and small. Anterior margin is curved. Posterior margin is straight or curved. Ventral margin is evenly rounded. Attachment area is medium, depressed occupies the posterodorsal portion of the LV. Its ornamentation consists of numerous radial ribs which are crossing by concentric growth lamellae at irregular intervals. Ribs start at the umbonal area, occasionally carrying small spines near ventral margin. RV ornamented by concentric fine growth squamae.

REMARKS

Ostrea tunetana Munier-Chalmas, 1881 shows some similarity with our material but differs in having longer and more pointed umbo and straight ligamental area.

Ostrea sp. C
(Fig. 4E)

MATERIAL. — 3 specimens (MDG/LBV/Lm-32.1 to Lm-32.3).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell, represented only by one LV, is medium, inequilateral, elongated to elliptical shape in outline. Umbo is small and opisthogyrate. Ligamental area is larger than high. Anterior margin and posterior margin are subparallel. Ventral margin is evenly rounded. Attachment area is long and suboval occupies the posterodorsal part of valve. Ornamentation consists of numerous radial ribs which interrupted by free standing frilled delicate growth squamae.

REMARKS

This species differs from *Ostrea* sp. B, in having a long and suboval attachment area and large ligamental area.

Ostrea sp. D
(Fig. 4A-D)

MATERIAL. — 25 specimens (MDG/LBV/Lm-33.1 to –33.25).

OCCURRENCE. — Turonian of Gabon (this study).

DESCRIPTION

Shell is medium to large, inequivalve, inequilateral, elongated to elliptical shape in outline. LV is slightly inflated and RV flat. Umbo is small and opisthogyrate. Ligamental area is small. Anterior margin is curved. Posterior margin is straight or curved. Ventral margin evenly rounded. Attachment area is medium, depressed occupies the posterodorsal portion of the LV. Its ornamentation consists of numerous radial ribs which are crossing by concentric growth lamellae at irregular intervals. Ribs start at the umbonal area, occasionally carrying small spines near ventral margin. RV is ornamented by concentric fine growth squamae.

TABLE 2. — Summary of life-habit and feeding mode of the oysters identified in this study. Abbreviations: **EC**, epifaunal cemented, **EF**, epifaunal free mobile; **S**, suspension feeders.

Families	Species	Life habit	Feeding mode	Abundance
Gryphaeidae	<i>Gryphaeostrea</i> sp.	EC	S	3
	<i>Ilymatogyra</i> (<i>Afrogyra</i>) <i>africana</i> (Lamarck, 1801)	EC	S	6
	<i>Pycnodonte</i> sp.	EC	S	1
	<i>Rhynchostreon</i> cf. <i>suborbiculatum</i> (Lamarck, 1801)	EC/EF	S	1
Flemingostreidae	<i>Curvostrea tevesthensis</i> (Coquand, 1862)	EC	S	1
	<i>Gyrostrea delectrei</i> (Coquand, 1862)	EC/EF	S	10
	<i>Gyrostrea</i> sp. A	EC/EF	S	1
	<i>Gyrostrea</i> sp. B	EC/EF	S	1
Ostreidae	<i>Ostrea</i> sp. A	EC	S	10
	<i>Ostrea</i> sp. B	EC	S	81
	<i>Ostrea</i> sp. C	EC	S	3
	<i>Ostrea</i> sp. D	EC	S	25

REMARKS

Ostrea sp. D differs to *Ostrea delectrei* (Coquand 1869: 143, pl. 47, figs 1-6) in being smooth towards umbo and in having small space between concentric growth lamellae.

PALAEOECOLOGY

DESCRIPTION

The fossil oyster assemblage from PK 12 section, which found only within bed 30 (Fig.), is rich in specimens but relatively poor in species diversity. We have identified twelve species from seven genera, namely *Curvostrea tevesthensis* (Coquand, 1862), *Gryphaeostrea* sp., *Gyrostrea delectrei* (Coquand, 1862), *Gyrostrea* sp. A, *Gyrostrea* sp. B, *Ilymatogyra* (*Afrogyra*) *africana* (Lamarck, 1801), *Ostrea* sp. A, *Ostrea* sp. B, *Ostrea* sp. C, *Ostrea* sp. D, *Pycnodonte* sp., *Rhynchostreon* cf. *suborbiculatum* (Lamarck, 1801). This oyster assemblage is dominated by representatives of *Ostrea* genus with 119 specimens (Table 2). The oysters are not associated with other organisms. Assemblage is characterized by thin and thick shells which are medium to large in size. All left valves have attachment area and prominent ornamentation. In comparison with previous studies of oyster fauna from Gabonese coastal basin (Lombard 1930; Darteville & Freneix 1957; Musavu Moussavou *et al.* 2013a, 2014a), oysters studied herein are provided 12 species *versus* 7 species in Gabonese coastal basin area. In addition, all species are recorded for the time in Gabonese coastal basin.

INTERPRETATION

Many studies have treated the distribution patterns of marine macroinvertebrates (e.g., Fürsich 1984, 1994; Berndt 2002; Delvene 2003; Fürsich *et al.* 2004; Harzhauser & Mandic 2004; Zuschin *et al.* 2004a, b; Fürsich & Thomsen 2005; Schneider *et al.* 2009; Ayoub-Hannaa 2011; Ayoub-Hannaa & Fürsich 2012; Bengtson *et al.* 2014). The parameters controlling faunal distribution are substrate, water energy, salinity, temperature, nutrient supply and oxygen availability. The palaeoautecology of oysters is summarised in Table 2. It is based on information by Berndt (2002), Delvene (2003), Ayoub-Hannaa (2011), Ayoub-Hannaa & Fürsich (2012), Sorensen *et al.* (2012).

According to Morrison & Brand (1986), bivalves living in cold and quiet water have thin, chalky shell, whereas those from warmer climates have a thicker with more consistent growth lines. In PK 12 section, thin and thick shells of oysters are together. Because left valves have a prominent ornamentation, we consider that oyster assemblage studied herein evolved in shallow-water where are exposed to sunlight.

With respect to substrate and sedimentation rate, the absence of other organisms in bed 30, such as infaunal organisms, the presence of attachment area on all left valve indicate a firm substrate and relatively low sedimentation rate (Stenzel 1971; Ayoub-Hannaa 2011; Ayoub-Hannaa & Fürsich 2012). On the other hand, the studied oysters would have tended to sink their shells into a firm substrate without the danger of ultimately burial (Abdel Aal & El-Hedeny 1998). The absence of infaunal organisms may be also due to early diagenetic dissolution of aragonitic shells.

All oysters are suspension-feeders. This means that the water energy level was sufficiently high to keep organic matter in suspension and refers also to a nutrient-rich and well oxygenated water column (Berndt 2002; Delvene 2003; Fürsich & Thomsen 2005; Ayoub-Hannaa 2011; Ayoub-Hannaa & Fürsich 2012). The absence of deposit feeders suggests probably a very small amount of organic matter in the sediment or organics in the water column (Ayoub-Hannaa & Fürsich 2012). This, suggest also an environment at least intermittently influenced by waves and/or currents (Fürsich & Heinberg 1983).

TAPHONOMY

DESCRIPTION

In PK 12 section, 149 disarticulated and 55 articulated specimens are collected. Highly fragmented of shells are generally lacking. The disarticulated left and right oyster valves are almost equal in proportion. Oyster shells are non-bioeroded, scarcity encrusted (4.4 %), rarely fragmented (0.98%) and presented rare traces of predation (0.98%).

INTERPRETATION

Many studies have used taphonomic features in order to follow the evolution of shell since its death until its final burial (e.g. Powell *et al.* 1989; Kidwell & Bosence 1991; Brett & Baird 1993; El-Hedeny 2005; Mekawy 2013; Hernandez-Ocana *et al.* 2015). Taphonomic process commonly described includes decomposition, dissolution, abrasion, bioerosion, fragmentation and the biological and hydraulic reworking of skeletal remains.

According to Boucot *et al.* (1958), the degree of disarticulation, along with abrasion and breakage, is a reliable indication of the degree of transport a bivalve has undergone since death. In this study, the stage of disarticulation of valves and fragmentation of shells, suggests that oyster community from PK12 section is transported under high-energy conditions, but of short duration and a short distance after death (Boucot *et al.* 1958; Seeling & Bengtson 1999). Therefore the assemblage of oysters can be regarded as parautochthonous.

The good quality of oyster shells, absence of bioerosion and scarcity of encrustation suggest a rapid burial rate after death and a relatively low sedimentation rate (Ayoub-Hannaa 2011; Ayoub-Hannaa & Fürsich 2012; Mekawy 2013). Only 2 (0.98%) from 204 of studied oyster specimens are presented traces of predation. This means scarcity of predators during lives of organisms. This is corroborated by the absence of other organisms in assemblage. Because the scarcity of predators, we regarded the high frequency of disarticulated oyster shells as due only to very weak dysodont dentition (El-Hedeny 2005).

CONCLUSIONS

The Turonian oysters of PK 12 section are characterized by thin and thick shells which are medium to large in size. The fauna is dominated by representatives of *Ostrea*. All morphotypes are recorded here for the first time in the Gabonese coastal basin. In comparison with previous studies of oyster fauna from Gabonese coastal basin, oysters from PK 12 section have a higher diversity and differ on their systematic content.

Paleoecologically, oysters were living in nearshore shallow, high-energy marine environments which are rich in nutrient and have a well oxygenated water column. After their death, they have been rapidly buried.

Taphonomically, the studied oysters are affected by disarticulation. Bioerosion, traces of predators and encrustation are scarcity or absence. Abrasion, breakage and high fragmentation of shell are generally lacking. The assemblage of oysters can be regarded as parautochthonous.

Acknowledgements

I am particularly grateful to Peter Bengtson and the Journal editorial board for their constructive comments, which greatly improved the manuscript.

REFERENCES

- ABDEL AAL A. A. & EL-HEDENY M. M. 1998. — On the variability of the Campanian *Pycnodonte* (*Phygraea*) *vesiculare* (Lamark). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1: 42-54.
- AQRABAWI M. 1993. — Oysters (Bivalvia-Pteriomorpha) of the Upper Cretaceous rocks of Jordan: palaeontology, stratigraphy and comparison with the Upper Cretaceous oysters of northwest Europe. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg* 75: 1-136.
- AYOUB-HANNA W. S. 2011. — *Taxonomy and Palaeoecology of the Cenomanian-Turonian Macro-Invertebrates from Eastern Sinai, Egypt*. Thèse de Bayerischen Julius-Maximilians-Universität Würzburg, 386 p.
- AYOUB-HANNA W. S. & FÜRSICH F. T. 2012. — Palaeoecology and environmental significance of benthic associations from the Cenomanian-Turonian of eastern Sinai, Egypt. *Beringeria* 42: 93-138.
- BASSE E. 1932. — Paléontologie de Madagascar. XVIII. Faune malacologique du crétacé supérieur du sud ouest de Madagascar. *Annales de Paléontologie* 21: 91-104.
- BENGTSON P., AYOUB-HANNA W. S. & FÜRSICH F. T. 2014. — Taxonomy and palaeoecology of Pinna (P) cretacea (Scholtheim, 1813) from the Upper Cretaceous of the Sergipe basin, Brazil. *Revista Brasileira de Paleontologia* 17 (3): 289-306. <https://doi.org/10.4072/rbp.2014.3.02>
- BERNDT R. 2002. — *Palaeoecology and Taxonomy of the Macrobenthic Fauna from the Upper Cretaceous Ajlun Group, Southern Jordan*. Unpublished PhD Thesis Würzburg University Germany, 222 p.
- BOUCOT A. J., BRACE W. & DEMAR R. 1958. — Distribution of brachiopod and pelecypod shells by currents. *Journal of Sedimentary Petrology* 28: 321-332. <https://doi.org/10.1306/74D707EA-2B21-11D7-8648000102C1865D>
- BRETT C. E. & BAIRD G. C. 1993. — Taphonomic approaches to temporal resolution in stratigraphy, in KIDWELL S. M. & BEHRENSMEYER A. K. (eds), Taphonomic approaches to time resolution in fossil assemblages. *Paleontological Society, Short Course* 6: 250-274.
- CARTER J. G., ALTABA C. R., ANDERSON L. C., ARAUJO R., BIAKOV A. S., BOGAN A. E., CAMPBELL D. C., CAMPBELL M., CHEN J. H., COPE J. C. W., DELVENE G., DIJKSTRA H. H., FANG Z. J., GARDNER R. N., GAVRILOVA V. A., GONCHAROVA I. A., HARRIES P. J., HARTMAN J. H., HAUTMANN M., HOEH W. R., HYLLEBERG J., JIANG B.-Y., JOHNSTON P., KIRKENDALE L., KLEEMANN K., KOPKA J., KRÍŽ J., MACHADO D., MALCHUS N., MÁRQUEZ-ALIAGA A., MASSE J.-P., MCROBERTS C. A., MIDDELFART P. U., MITCHELL S., NEVESSKAJA L. A., ÖZER S., POJETA JR. J., POLUBOTKO I. V., MARIA PONS J., POPOV S., SÁNCHEZ T., SARTORI A. F., SCOTT R. W., SEY I. I., SIGNORELLI J. H., SILANTIEV V. V., SKELTON P. W., STEUBER T., WATERHOUSE J. B., WINGARD G. L. & YANCEY T. 2011. — A synoptical classification of the Bivalvia (Mollusca). *Paleontological Contributions* 4: 1-47. <https://doi.org/10.17161/PC.1808.8287>
- CHEVALIER L., MAKANGA J. F. & THOMAS R. J. 2002. — *Carte géologique de la République gabonaise au 1/1 000 000*. Notice explicative. Council for geoscience, 195 p.
- CHOUBERT B. 1935. — Sur les terrains crétacés de la zone côtière du Gabon. *Comptes Rendus de l'Académie des Sciences de Paris* 201: 401-403.
- COOPER M. R. 1995. — Exogyrid oysters (Bivalvia: Gryphaeidae) from the Cretaceous of southeast Africa. Part 1. *Durban Museum Novitates* 20: 1-48.
- COOPER M. R. 2002. — Palaeolophid and liostreine oysters (Bivalvia: Ostreidae) from the Cretaceous of southeast Africa, with comments on oyster biostratigraphy and biofacies. *Durban Museum Novitates* 27: 28-60.
- COQUAND H. M. 1862. — *Géologie et paléontologie de la région sud de la province de Constantine*. Arnaud, Marseille, 343 p.

- COQUAND H. M. 1869. — *Monographie du genre 'Ostrea'. Terrain crétacé*. H. Seren, Marseille, 215 p. <http://gallica.bnf.fr/ark:/12148/bpt6k56522857>
- DARTEVELLE E. & BRÉBION P. 1956. — Mollusques fossiles du Crétacé de la côte occidentale d'Afrique du Cameroun à l'Angola. I. Gastéropodes. *Annales du Musée royal du Congo belge*, Série 8, *Sciences géologiques* 15, 128 p.
- DARTEVELLE E. & FRENEIX S. 1957. — Mollusques fossiles du Crétacé de la côte occidentale d'Afrique du Cameroun à l'Angola. II. Lamellibranches. *Annales du Musée royal du Congo belge*, Série 8, *Sciences géologiques* 20, 271 p.
- DELVEY G. 2003. — Middle and Upper Jurassic bivalve associations from the Iberian Range (Spain). *Geobios* 36: 519-531. [https://doi.org/10.1016/S0016-6995\(03\)00058-5](https://doi.org/10.1016/S0016-6995(03)00058-5)
- EL-HEDENY M. M. 2005. — Taphonomy and Paleoecology of the Middle Miocene oysters from Wadi Sudr, Gulf of Suez, Egypt. *Revue de Paléobiologie* 24 (2): 719-733. <http://hdl.handle.net/123456789/11307>
- EL-SHEIKH H. A., ABDELHAMID M. A. & EL QOT G. M. 1998. — Macrofossils and foraminiferal biostratigraphy and paleoecology of some Cenomanian-Santonian sequences in north and west central Sinai, Egypt. *Egyptian Journal of Geology* 42: 471-495.
- FISCHER J. C. 1980. — *Fossiles de France et des régions limitrophes. Guides géologiques régionaux*. Masson, Paris, 444 p.
- FOURTAU R. 1917. — Catalogue des invertébrés fossiles de l'Égypte représentés dans la collection du Musée de Géologie au Caire. Terrains Crétacés 2^e partie: Mollusques Lamellibranches. *Geological Survey of Egypt, Palaeontological Series* 3: 1-108
- FRENEIX S. 1959. — Mollusques fossiles du Crétacé de la côte occidentale d'Afrique du Cameroun à l'Angola. III. Conclusions stratigraphiques et paléontologiques. *Annale du Muséum du Royaume du Congo belge*, Série 8, *Sciences géologiques* 24, 126 p.
- FRENEIX S. 1966. — Faunes de Bivalves et corrélation des faunes marines du Crétacé des bassins côtiers de l'Ouest africain- Bassins sédimentaires du littoral africain, in REYRE D. (ed.), *Symposium New Delhi, 1964. Association des Services géologiques africains*, Paris 1: 52-78.
- FRENEIX S. 1972. — Les mollusques bivalves du Bassin côtier de Tarfaya (Maroc méridional). *Notes et Mémoires du Service géologique du Maroc* 228: 49-255.
- FRENEIX S. & VIAUD J. M. 1986. — Huîtres du Crétacé supérieur du Bassin de Challans-Commequiers (Vendée). Biostratigraphie, taxinomie, paléobiologie. *Bulletin trimestriel de la Société géologique de Normandie et Amis du Muséum du Havre* 73 (1-2): 13-19.
- FURON R. 1931. — Sur la géologie du Gabon (A.E.F.). *Comptes Rendus de l'Académie des Sciences de Paris* 192: 168-169.
- FURON R. 1950. — *Géologie de l'Afrique*. Payot, Paris, 1 vol., 350 p.
- FÜRSICH F. T. 1984. — Palaeoecology of boreal invertebrate faunas from the Upper Jurassic of central Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology* 48: 309-346. [https://doi.org/10.1016/0031-0182\(84\)90050-6](https://doi.org/10.1016/0031-0182(84)90050-6)
- FÜRSICH F. T. 1994. — Palaeoecology and evolution of Mesozoic salinity-controlled benthic macroinvertebrate associations. *Lethaia* 26: 327-346. <https://doi.org/10.1111/j.1502-3931.1993.tb01540.x>
- FÜRSICH F. T., OSCHMANN W., PANDEY D. K., JAITLEY A. K., SINGH I. B. & LIU C. 2004. — Palaeoecology of middle to lower Upper Jurassic macrofaunas of the Kachchh Basin, western India: an overview. *Journal of the Palaeontological Society of India* 49: 1-26.
- FÜRSICH F. T. & HEINBERG C. 1983. — Sedimentology, biostratigraphy, and palaeoecology of an Upper Jurassic offshore sand bar complex. *Bulletin of the Geological Society of Denmark* 32: 67-95.
- FÜRSICH F. T. & THOMSEN E. 2005. — Jurassic biota and biofacies in erratics from the Sortland area, Vesterålen, northern Norway. *Norges geologiske undersøkelse Bulletin* 443: 37-53.
- GÉRARD G. 1958. — *Carte géologique de l'Afrique équatoriale française au 1/2 000 000. Notice explicative*. Gouvernement général de l'Afrique équatoriale française, Direction des Mines et de la Géologie, 198 p.
- HARZHAUSER M. & MANDIC O. 2004. — The muddy bottom of Lake Pannon a challenge for dreissenid settlement (Late Miocene; Bivalvia). *Palaeogeography, Palaeoclimatology, Palaeoecology* 204: 331-352. [https://doi.org/10.1016/S0031-0182\(03\)00735-1](https://doi.org/10.1016/S0031-0182(03)00735-1)
- HERNANDEZ-OCANA M. I., QUIROZ-BARROSO S. A. & SOUR-TOVAR F. 2015. — Tafonomía y Paleocología de las ostras de la formación San Juan Raya, Aptiense del sureste de Puebla, Mexico. *Boletín Geológico y Minero* 126 (1): 37-62.
- HOUREQ V. & HAUSKNECHT J.-J. 1954. — *Notice explicative sur la faille Libreville-Ouest et carte géologique de reconnaissance au 1/500 000*. Gouvernement général de l'Afrique équatoriale française, Paris, 25 p.
- HUDELEY H. & BELMONTE Y. 1970. — *Carte géologique de la République gabonaise au 1/100 000. Notice explicative*. Bureau de Recherche géologiques et minière, Mémoire 72, 191 p.
- KIDWELL S. M. & BOSENCE D. W. 1991. — Taphonomy and time-averaging of marine shelly fauna, in ALLISON P. A. & BRIGGS D. E. G. (eds), *Taphonomy: Releasing the Data Locked in the Fossil Record*. Springer, London: 115-209.
- KOSSMAT F. 1893. — Über einige Kreideversteinerungen vom Gabun. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften* 102: 575-590.
- LAMARCK J. B. P. 1801. — *Système des animaux sans vertèbres*. Deterville, Paris, 432 p. <http://www.biodiversitylibrary.org/bibliography/14255>
- LOMBARD J. 1930. — Céphalopodes et Lamellibranches crétacés du Congo Français. *Bulletin de la Société géologique de France* 4 (30): 277-322.
- MALCHUS N. 1990. — Revision der krei-Austern (Bivalvia: Pteriormorphia) Ägyptens (Biostratigraphie, Systematik). *Berliner Geowissenschaftliche Abhandlungen (Reihe A)* 125: 1-231.
- MBINA MOUNGUENGUI M. 1998. — *Dynamique sédimentaire et fluctuations eustatiques au cours du Cénomaniens et du Turonien basal dans le Nord du bassin côtier gabonais*. Thèse de l'Université de Bourgogne, 376 p.
- MEISTER C., MBINA MOUNGUENGUI M. & LANG J. 1996. — Les ammonites Cénomano-Turonien du Gabon: intérêt pour la liaison Téthys-Atlantique Sud et corrélations. 39^e Congrès Brésilien de Géologie. IGCP projet 3881 (South Atlantic Mesozoic Correlations SAMC), News 5, Abstract, 405-407.
- MEISTER C., MBINA MOUNGUENGUI M. & LANG J. 2003. — Les ammonites du Cénomano-Turonien du bassin côtier nord-gabonais: systématique et intérêt paléogéographique pour la liaison Téthys-Atlantique Sud. *Revue de Paléobiologie* 22 (1): 341-335.
- MEKAWY M. S. 2013. — Taphonomy of Aptian-Albian beds in the Gebel Mistan, Maghara area, Northern Sinai, Egypt. *Journal of Earth Science & Climatic Change* 4: 135. <https://doi.org/10.4172/2157-7617.1000135>
- MORRISON R. & BRAND U. 1986. — Paleocene 5: Geochemistry of recent marine invertebrates. *Geoscience Canada* 13 (4): 237-254.
- MUNIER-CHALMAS E. 1881. — Note paléontologique sur les fossiles recueillis par M. le Commandant Roudaire dans son expédition scientifique en Tunisie. Extraits de la Mission de M. le Commandant Roudaire dans les chotts tunisiens (1878-1879). Paris, G. Chamerot, 79 p.
- MUSAVU MOUSSAVOU B. 2015. — Bivalves (Mollusca) from the Coniacian-Santonian Anguille Formation from Cap Esterias, Northern Gabon, with notes on paleoecology and paleobiogeography. *Geodiversitas* 37 (3): 315-324. <https://doi.org/10.5252/g2015n3a2>
- MUSAVU MOUSSAVOU B. & MABICKA OBAME R. 2015. — Taxonomy and paleoecology of Albian Madiela Formation gastropods, Gabonese coastal basin: N'Toum quarry section. *Bulletin de la Société géologique de France* 186 (1): 35-42. <https://doi.org/10.2113/gssgfbull.186.1.35>
- MUSAVU MOUSSAVOU B., NDONG ONDO S. M. & MAKAYA M'VOUBOU 2013a. — Turonian bivalves from the Coastal Basin of Gabon, South of Libreville. *Bulletin de l'Institut scientifique, Rabat, section Sciences de la Terre* 35: 1-8.

- MUSAVU MOUSSAVOU B., MAKAYA M'VOUBOU & MABICKA OBAME R. 2013b. — Contribution à la connaissance de la famille Aporrhaidae Gray 1850 dans le sous bassin nord du bassin côtier gabonais. *Africa Geosciences Review* 20 (3-4): 97-103.
- MUSAVU MOUSSAVOU B., MAKAYA M'VOUBOU & NDONG ONDO S. M. 2014a. — Turonian bivalves from Gabonese coastal basin: Belle vue 3 section, South of Libreville region. *Bulletin de la Société géologique de France* 185 (1): 43-50. <https://doi.org/10.2113/gssgfbull.185.1.43>
- MUSAVU MOUSSAVOU B., NDONG ONDO S. M. & MAKAYA M'VOUBOU 2014b. — Les gastéropodes du Turonien de la coupe de Bellevue 3 dans la région de Libreville Sud au sein du bassin côtier gabonais. *Revue de Paléobiologie, Genève* 33 (2): 419-429.
- NILSSON S. 1827. — *Petrefacta Suecana Formationis Cretacea, descripta et iconibus illustrata. Pars prior, Vertebrata et Mollusca sistens*. Ex Officina Berlingiana (Londini Gothorum [Lund]). <https://doi.org/10.5962/bhl.title.106190>
- PERVINQUIÈRE L. 1912. — Étude de la paléontologie tunisienne, II, gastéropodes et lamellibranches des terrains crétacés. *Carte géologique de la Tunisie*. J. Lamarre et Cie, Paris, 352 p.
- POWELL E. N., STAFF G. M., DAVIES D. J. & CALLENDER W. R. 1989. — Macro-benthic death assemblages in modern marine environments: Formation, interpretation and application: *Critical Reviews in Aquatic Sciences* 1: 555-589.
- PUGACZEWSKA H. 1977. — The upper Cretaceous ostreidae from the middle Vistula region (Poland). *Acta Palaeontologica Polonica* 22 (2): 187-204.
- SEELING J. & BENGTSON P. 1999. — Cenomanian oyster from the Sergipe Basin, Brazil. *Cretaceous Research* 20: 747-765. <https://doi.org/10.1006/cres.1999.0190>
- SCHNEIDER S., BERNING B., BITNER M. A., CARRIOL R.-P., JÄGER M., KRIWET J., KROH A. & WERNER W. 2009. — A parautochthonous shallow marine fauna from the Late Burdigalian (early Ottnangian) of Gurlarn (Lower Bavaria, SE Germany): Macrofaunal inventory and paleoecology. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 254: 63-103. <https://doi.org/10.1127/0077-7749/2009/0004>
- SORENSEN A. M., SURLYK F. & JAGT J. W. M. 2012. — Adaptive morphologies and guild structure in a high-diversity bivalve fauna from an early Campanian rocky shore, Ivö Klack (Sweden). *Cretaceous Research* 33: 21-41. <https://doi.org/10.1016/j.cretres.2011.07.004>
- STENZEL H. B. 1971. — Oysters, in MOORE R. C. (ed.), *Treatise on Invertebrate Paleontology, Part N. Mollusca 6, Bivalvia, vol 3*. Geological Society of America & University of Kansas, Boulder & Lawrence: 953-1224.
- TEISSERENC P. & VILLEMIN J. 1990. — Sedimentary Basin of Gabon – Geology and Oils Systems, in EDWARDS J. D. & SANTOGROSSI A. (eds), *Divergent/Passive margin basins. American Association of Petroleum Geologists Bulletin, memoir* 48: 117-199.
- TREVISAN L. 1937. — La fauna ei giacimenti del Cenomaniano di facies Africana della Sicilia occidentale. *Mémoire dell'Institut Geologico della Universita di Padova* 12: 1-132.
- VIDET B. 2003 — *Dynamique des paléoenvironnements à huîtres du Crétacé supérieur Nord-Aquitain (SO France) et du Mio-Pliocène Andalou (SE Espagne)*. Thèse de l'Université de Rennes 1, 263 p.
- WEYDERT P. 1981. — Les faciès péri-récifaux du turonien de Libreville (Gabon): significations paléocéologiques au cours de l'ouverture de l'Atlantique. *Comptes Rendus de l'Académie des Sciences, Paris* 295: 95-100.
- WEYDERT P. & COLLIGNON M. 1981. — *Schindewolfites gabonensis* nov. sp. et *furoniceras?*, ammonites du Turonien de Libreville (Gabon). Doctorat Laboratoire de Lyon, h.s. 6: 293-299.
- ZUSCHIN M., HARSHAUSER M. & MANDIC O. 2004a. — Spatial variability within a single parautochthonous Paratethyan tidal flat deposit (Karpatian, Lower Miocene-Kleinebersdorf, Lower Austria). *Courier Forschungsinstitut Senckenberg* 246: 153-168.
- ZUSCHIN M., HARSHAUSER M. & MANDIC O. 2004b. — Taphonomy and Palaeoecology of the Lower Badenian (Middle Miocene) molluscan assemblages at Grund (Lower Austria). *Geologica Carpathica* 55: 117-128.

Submitted on 25 October 2015;
accepted on 29 October 2016;
published on 30 June 2017.