

First record of the new *Neoplanorbulinid* species (Foraminifera) from the Early Oligocene in Turkey, Malatya Basin, Eastern Taurids

Fatma GEDIK

General Directorate of Mineral Research and Exploration,
Geological Research Department, 06800 Ankara (Turkey)
fatmagedik@yahoo.com

Published on 30 June 2017

urn:lsid:zoobank.org:pub:A7EF2FC3-B9F5-429D-ABA4-798B54B6091F

Gedik F. 2017. — First record of the new *Neoplanorbulinid* species (Foraminifera) from the Early Oligocene in Turkey, Malatya Basin, Eastern Taurids. *Geodiversitas* 39 (2): 273–284. <https://doi.org/10.5252/g2017n2a6>

ABSTRACT

The genus *Neoplanorbulinella* Matsumaru, 1976 was firstly identified in the Lower Miocene of Saipan and Mariana Islands in the Western Pacific. The present paper reports on its occurrence in the Lower Oligocene of Malatya Basin (Eastern Taurids, East Turkey). *Neoplanorbulinella* is characterized by a conical test, having several sized equatorial chambers irregularly arranged around the cone, and lateral chambers underdeveloped in umbilical cavity. The new planorbulinid species, namely *Neoplanorbulinella matsumarui* n. sp. and *Neoplanorbulinella malatyaensis* n. sp., are recorded in the Muratlı Formation (Malatya Basin, Turkey), which is dated to the Rupelian-early Chattian. The new species are associated to an assemblage of the SB 21–22 larger foraminiferal zone. The new species are described, and their affinities are discussed.

RÉSUMÉ

Signalement d'une nouvelle espèce de Neoplanorbulinid (Foraminifera) de l'Oligocène inférieur du bassin de Malatya (Taurides orientales, Est de la Turquie).

Le genre *Neoplanorbulinella* Matsumaru, 1976 a d'abord été identifié dans le Miocène inférieur des îles de Saipan et de Mariana dans le Pacifique occidental. Le présent article rend compte de sa présence dans l'Oligocène inférieur du bassin de Malatya (Taurides orientales, Est de la Turquie). *Neoplanorbulinella* est caractérisée par un test conique, présentant plusieurs chambres équatoriales de taille irrégulière disposées autour du cône, et des chambres latérales moins développées dans la cavité ombilical. Les nouvelles espèces de Planorbulinidae, à savoir *Neoplanorbulinella matsumarui* n. sp. et *Neoplanorbulinella malatyaensis* n. sp., ont été trouvées dans la formation de Muratlı (bassin de Malatya, Turquie), qui est datée du Rupélien-Chattien inférieur. Les nouvelles espèces sont associées à un assemblage de la zone de grands foraminifères SB 21–22. Les nouvelles espèces sont décrites, et leurs affinités discutées.

KEY WORDS

Foraminifera,
Systematics,
Oligocene,
Malatya Basin,
Turkey,
new species.

MOTS CLÉS

Foraminifera,
Systématique,
Oligocène,
Bassin de Malatya,
Turquie,
espèces nouvelles.

INTRODUCTION

The Oligocene is one of the most significant time intervals for the geology of Turkey as Neo-Tethys came to end in this time through the collision of the African and Eurasian plates (Şengör & Yılmaz 1981; Rögl 1999). In addition to the initiating Taurus mountain range, which is a part of the Alp-Himalayan orogenic belt, the closure of the Neo-Tethys Ocean caused new basins to form and/or existing Anatolian basins to change their characteristics. One of changed basins was the Malatya Basin which is the object of this study (Fig. 1). The Malatya Basin is a major interior basin in East Anatolia, approximately 1000 km² wide, exposing Upper Cretaceous-Cenozoic volcano-sedimentary rocks up to 4500 m thick (Çağlar 2009).

Several geological studies with different aims have been done in the investigated area by the following authors: Ayan 1961; Akkuş 1971; Yoldaş 1972; Kurtman 1978; Örcen 1986; Karaman *et al.* 1993; Alkan 1997; Çağlar 2009; Gedik 2010, 2014, 2015 and Karadenizli *et al.* 2016. In this long stratigraphic succession, Karaman *et al.* (1993) reported that the Muratlı Formation is Oligocene in age based on the foraminiferal data. For some other students (Yoldaş 1972; Örcen 1986) the age of the Muratlı Formation is Aquitanian. The unit starts with a brecciated limestone at the base, followed upward by clastic limestones, marls and limestones with abundant benthic foraminiferal assemblages of Oligocene age.

Oligocene shallow marine sedimentary successions of the Neo-Tethys realm contain diagnostic hyaline larger foraminifera such as miogypsinids, lepidocyclinids and nummulitids that are used for biostratigraphic dating and regional correlations (Drooger 1993; BouDagher-Fadel 2008). Furthermore, in the Oligocene of the Near and Middle East, diverse associations of porcelaneous larger foraminifera are known to be common and widespread since the work of Henson (1948a, b; 1950). Especially, the representatives of the subfamily Archaiasinae are common and associated with *Peneroplis glynnjonesi* (Henson, 1950), *P. thomasi* Henson, 1950, *Praerhapydionina delicata* Henson, 1950 and various species of *Austrotrillina* Parr, 1942, *Borelis* de Montfort, 1808 and *Bullalveolina* Reichel, 1936. As Henson (1950) demonstrated, these associations are clearly linked to a particular, very shallow, often restricted marine carbonate facies (see also Bassi *et al.* 2007).

The section measured in this study represents an uninterrupted Oligocene sequence, chiefly composed of carbonate/clastic succession, and includes large benthic foraminifera. In addition to this, this section is important in containing porcelaneous large foraminifera associated with *Penarchaias glynnjonesi* (Henson, 1950), *Austrotrillina brunni* Marie, 1955, various species of *Borelis* de Montfort, 1808, two new Neoplanorbulinid species (see here below), indicative of very shallow-water carbonate facies, and the hyaline group with miogypsinids, lepidocyclinids and nummulitids, which characterizes shallow-water marine environments.

This study presents a systematic description of two new Neoplanorbulinid species, obtained from the Rupelian-lower Chattian limestones of the Karamağara section (Figs 2; 4). In a previous study, Sirel & Gedik (2011) reported on the occurrence of a new miogypsinid genus and species, *Post-miogypsinella intermedia* Sirel & Gedik, 2011, from the late Chattian shallow marine limestone of the Karamağara section, associated with foraminiferal species.

The interpretation of structural elements is given in Fig. 5 for the new Neoplanorbulinid species as well as for the already known Early Miocene species described by K. Matsumaru.

MATERIAL AND METHODS

This study is based on samples collected along the Karamağara section in the Muratlı Formation. The examined section is located to the northeast of Akçadağ, 35 km west of Malatya (Fig. 2). In this section, strata comprise limestones, sandy/clayey limestones and marls. The foraminifera were recovered mainly from the limestones. The foraminiferal fauna was studied in non-oriented thin sections because the indurated carbonate rocks do not allow specimens to be freed.

ABBREVIATIONS

HYM	Karamağara section sample numbers;
SB	shallow marine benthic foraminiferal zones;
MSS	measured stratigraphical section.

STRATIGRAPHY AND LITHOLOGY OF THE KARAMAGARA SECTION

This studied Karamağara section (NE-SW) lies to the north of Karamağara village, situated on the 1/25 000 topographic sheet of Malatya L40-a1 (bottom coordinates X1: 16 599; Y1: 57 032 and the top X2: 12 705; Y2: 58 026) (Fig. 1). Sirel & Gedik (2011) and Gedik (2015) already provided a detailed lithological description of this section and some biostratigraphical data. Oligocene-Early Miocene deposits with abundant shallow-water foraminiferal species are well developed in the Karamağara section. The sample numbers and stratigraphic distribution of the foraminiferal species are shown in Figure 4. The following lithologic units are observed from bottom to top.

The lithologic succession of the Karamağara section begins with the breccia limestones without fossils. The first fossiliferous unit extends from the samples 35 to 40 and is composed of shallow-water sandy and clayey limestones, which include *Borelis pygmaea* (Hanzawa, 1930), *Borelis merici* Sirel & Gündüz, 1981, *Austrotrillina brunni* Marie, 1955, *Archaias hensoni* Smout & Eames, 1958, *Penarchaias glynnjonesi* (Henson, 1950), *Planorbulina bronnimanni* Bignot & Decrouez, 1982, *Amphistegina* sp. This assemblage indicates a Rupelian-early Chattian age (SB 21-22).

The second unit extends from the samples 40B to 42 and consists of alternations of marls and limestones. The




FIG. 1. — **A**, Tectonic map of Turkey showing the major tectonic structures (modified from Bozkurt 2001); **B**, limits of the Sivas-Erzincan, GAD (Gürün-Akışla-Divriği) and Malatya basins.




FIG. 2. — Geological map of the study area (simplified from Bilgiç 2002).

basal marl beds contain planktonic foraminiferal species, namely, *Globigerinoides primordius* Blow & Banner, 1962, *Tenuitellinata juvenilis* (Bolli, 1957), *Globigerinita incrusta*

Akers, 1955, *Globigerina leroi* Blow & Banner, 1962, *G. occlusa* Blow & Banner, 1962, *G. praebulloides* Blow, 1959, *G. gnaucki* Blow & Banner, 1962, *G. ouachitaensis*

Upper System	System	Series	Formation	Thickness (m)	Lithology	Explanation
TERTIARY	QUATER.					Debris and alluvials
						Unconformity
	PLIO-QUATER.		BEYLER DERESI	500		Conglomerate, sandstone
						Unconformity
	PLIOCENE		SULTAN SUYU	300		Conglomerate, sandstone, lacustrine limestone
						Unconformity
	MIOCENE	UPPER MIOCENE LOWER PIOCENE	HAÇOVA VOLKANICS	?		Tuff, agglomerate, basalt
						Unconformity
						Yellowish white, occasionally massive, very thick bedded limestone with much algal, reef, macrofossil molluscs with basal conglomerate.
	MIOCENE	LOWER MIOCENE	PETEK KAYA	400-750		Gray, medium to thick horizontal bedded marl
						Dirty white, pale brown, fine to medium bedded detritic limestone
						Detritic, clayey limestone
TERTIARY	OLIGOCENE		MURATLI	100-130		Brecciated limestone
						Dirty white, dirty yellow, thick to very thick bedded limestone
	EOCENE	MIDDLE-UPPER EOCENE	YEŞİL YURT	1000		Alternation of dirty yellow, pale brown sandstone, siltstone, shale marl, limestone
						Limestone
						Brown conglomerate
						Unconformity

FIG. 3. — Generalized stratigraphy of Cenozoic deposits in the Malatya Basin (modified from Karaman *et al.* 1993).

Howe & Wallace, 1932, *G. cf. anguliseturalis* Bolli, 1957, *Globorotaloides cf. suteri* Bolli, 1957, *Paragloborotalia* spp., *Globogadrina* sp., indicating the Chattian-Aquitanean interval. The limestone beds located at the top of the unit include the shallow-water benthic foraminiferal species *Post-miogypsinella intermedia* Sirel & Gedik, 2011, *Miogypsinella akcadagensis* (Gedik & Sirel, 2009), *M. borodinensis* Tan,

1936, *Miogypsinodella* sp. Considering the stratigraphic range of *M. borodinensis* Tan, 1936 and *M. akcadagensis* (Gedik & Sirel, 2009), this unit was dated as late Chattian (SB 23).

The third unit extends from the samples 42A to 44D. The marly horizon located at the bottom contains planktonic foraminiferal species, such as *Globoturborotalia euapertura*




Fig. 4. — Stratigraphic distribution of larger benthic and planktonic foraminifera and nannoplankton species in the Karamağara section, NE Akçadağ, W of Malatya, Eastern Turkey.




FIG. 5. — Diagnostic structural elements of the *Neoplanorbulinella*: **A**, *Neoplanorbulinella saipanensis* Matsumaru, 1976 (1976: 201, pl. 6, figs 1, 3-5); **B**, *Neoplanorbulinella malatyaensis* n. sp. (axial section); **C**, *Neoplanorbulinella malatyaensis* n. sp. (holotype, equatorial section); **D**, *Neoplanorbulinella matsumarui* n. sp. (axial section). Abbreviations: **Pr**, protoconch; **Dt**, deuteroconch; **Eqc**, equatorial chamber; **Ltc**, lateral chamber; **Spc**, spiral chamber; **St**, stolon.

(Jenkins, 1960), *Globigerina ouachitaensis* Howe & Wallace, 1932, *G. gnaucki* Blow & Banner, 1962, *G. occlusa* Blow & Banner, 1962, *G. leroyi* Blow & Banner, 1962, *G. praebulloides* Blow, 1959, *G. cf. angulisuturalis* Bolli, 1957, *G. ciperoensis* Bolli, 1954, *Globoquadrina dehiscens* (Chapman, Parr & Collins, 1934), *G. venezuelana* (Hedberg, 1937), *G. praedehiscens* Blow & Banner, 1962, *G. selli* Borsetti, 1959, *Globigerinoides primordius* Blow & Banner, 1962, and the calcareous nannoplankton species *Discoaster deflandrei* Bramlette & Riedel, 1954, *Coccolithus eopelagicus* (Bramlette & Riedel, 1954), *Cyclicargolithus abisectus* (Müller, 1970), *C. floridanus* (Roth & Hay, 1967), *Reticulofenestra hampdenensis* Edwards, 1973, *R. umbilica* (Levin, 1965), *R. reticulata* (Gartner & Smith, 1967), *R. hillae* Burky & Percival, 1971, *Zygrhablithus bijugatus* (Deflandre, 1959), *Lanternithus minutus* Stradner, 1962, *Ericsonia formosa* (Kamptner, 1963), *E. subdisticha* (Roth & Hay, 1967), *Cruciplacolithus tenuis* (Stradner, 1961), *Braarudosphaera*

bigelowi (Gran & Braarud, 1935), *Helicosphaera perchenielseniae* Haq, 1971, *H. euphratis* Haq, 1966, *H. obliqua* Bramlette & Wilcoxon, 1967, *H. kamptneri* Hay & Mohler, 1967, *H. recta* Haq, 1966, *Coronocyclus nitescens* (Kamptner, 1963), *Dictyococcites bisectus* (Hay, Mohler & Wade, 1966), *Pontosphaera multipora* (Kamptner, 1948), *P. plana* (Bramlette & Sullivan, 1961), *Micrantholithus crenulatus* (Bramlette & Sullivan, 1961), *Sphenolithus moriformis* (Brönnimann & Stradner, 1960), *S. predistentus* Bramlette & Wilcoxon, 1967, *S. distentus* (Martini, 1965), *S. delphix* Burky, 1973, *Sphenolithus* sp. all dated to the Early Miocene. The limestone beds located at the top of the unit do not contain any characteristic foraminiferal species.

The fourth unit, from the samples 44D to 48, is composed of reefal limestones, which contain *Miogypsina globulina* (Michelotti, 1841), *M. polymorpha* (Rutten, 1912), *M. cf. thecideaformis* (Rutten, 1912), *Miolepidocyclina* sp. and other foraminiferal taxa of Burdigalian (SB 25).

SYSTEMATICS

This study follows the systematic classification of Loeblich & Tappan (1987). All holotypes and paratypes are stored in the collection of the Natural History Museum of MTA (General Directorate of Mineral Research and Exploration of Turkey, Ankara), under no. MTA 2006/FG16.

Order FORAMINIFERIDA Eichwald, 1830
Suborder ROTALIINA Delage & Herouard, 1896
Family PLANORBULINIDAE Schwager, 1877

Genus *Neoplanorbulinella* Matsumaru, 1976

TYPE SPECIES. — *Neoplanorbulinella saipanensis* Matsumaru, 1976 by original designation.

Neoplanorbulinella matsumarui n. sp.
(Fig. 6A-D)

HOLOTYPE. — Axial section, Fig. 6A (sample no. HYM-36A/5/2).

TYPE LOCALITY. — Karamağara village, NE Akçadağ, W Malatya, Eastern Turkey, HYM-first unit.

AGE. — Rupelian-Early Chattian (SB 21-22).

DIAGNOSIS. — *Neoplanorbulinella* with small, high conical test with almost right apical angle, having equatorial chambers around the cone, and lateral chambers underdeveloped in umbilical cavity, and calcareous wall.

ASSEMBLAGE. — *Borelis pygmaea*, *B. merici*, *Penarchaias ghyngijonesi*, *Archaias hensoni*, *Neoplanorbulinella malatyaensis* n. sp., *Neoplanorbulinella* sp., *Malatyna* sp., *Borelis* sp., *Miogyssinella* sp. and *Asterigerina* sp.

ETYMOLOGY. — This species is dedicated to Professor Kunitaru Matsumaru to acknowledge his contribution to micropaleontological research.

DESCRIPTION

Only megalospheric form is observed. This form has a high conical test with small size. The basal diameter of the cone reaches 0.5 mm, and the maximum cone height is 0.3 mm. The apical angle is about 90°. The large, spherical megalosphere (its diameter 30-92 µm) is situated at the top of the cone, followed by arcuate equatorial chambers. The size of the equatorial chambers increases regularly from apex to the base.

COMPARISON

Neoplanorbulinella matsumarui n. sp. differs from *N. saipanensis* Matsumaru, 1976 in having high conical test and about 90° apex angle. *N. saipanensis* has low conical test and about 112° apex angle (Matsumaru 1976). In addition, the equatorial and

lateral chambers of *N. saipanensis* are arranged irregularly. The equatorial chambers of the new species are arranged regularly, and the connection of chambers is provided with stolons.

Neoplanorbulinella malatyaensis n. sp.
(Fig. 6E-H)

HOLOTYPE. — Equatorial section, Fig. 6F (sample no. HYM-36A/5/1).

TYPE LOCALITY. — Karamağara village, NE Akçadağ, W Malatya, Eastern Turkey, HYM-first unit.

AGE. — Rupelian-Early Chattian (SB 21-22).

DIAGNOSIS. — *Neoplanorbulinella* with large, low conical to lenticular test, having equatorial chambers around the cone, lateral chambers underdeveloped in umbilical cavity, and calcareous wall.

ASSEMBLAGE. — The same assemblage as for *Neoplanorbulinella matsumarui* n. sp.

ETYMOLOGY. — From Malatya, a city of eastern Turkey.

DESCRIPTION

Only megalospheric form is observed. This form has a low conical to lenticular test of large size. The basal diameter of the cone reaches 1.18 mm and the cone maximum height is 0.42 mm. The apical angle is about 140°. The large, spherical megalosphere (its diameter 65 µm) is situated at the top of the cone, and it is followed by the second chamber (its diameter 78 µm; Fig. 6F) and arcuate equatorial chambers. The umbilical cavity is very wide and its diameter is 0.85 mm. Lengthened umbilical cavity is filled by underdeveloped lateral chambers (Fig. 6H).

COMPARISON


Neoplanorbulinella malatyaensis n. sp. differs from *N. matsumarui* n. sp. in possessing larger and low conical to lenticular test, greater apex angle, almost flat base of cone, larger proloculus and greater number and larger equatorial chambers, which are observed in axial sections.

Neoplanorbulinella sp.
(Fig. 6I)

DESCRIPTION

Only one specimen is observed. This form has a very high conical test with large size. The basal diameter of the cone reaches 0.8 mm and the cone height is 0.8 mm. The apical angle is 90°. The spherical megalosphere (diameter 65 µm) is situated at the top of the cone, followed by arcuate equatorial chambers. The size of the equatorial chambers increases gradu-

Fig. 6. — Selected benthic foraminiferal species from the studied section in the Malatya Basin, Turkey: **A-D**, *Neoplanorbulinella matsumarui* n. sp.; **A**, holotype, axial section, megalospheric form, showing spherical megalosphere followed by equatorial chambers, umbilical cavity is filled 5 row lateral chambers (HYM-36A/5/2); **B**, axial section, megalospheric form (HYM-36A/11/1); **C, D**, axial sections (HYM-36A); **E-H**, *Neoplanorbulinella malatyaensis* n. sp.; **E**, axial section, megalospheric form, (HYM-36A/8/1); **F**, holotype, equatorial section, megalospheric form (HYM-36A/5/1); **G, H**, axial sections, megalospheric forms, showing small spherical megalosphere followed by equatorial chambers (HYM-36A/8 and HYM-36A/3/2); **I**, *Neoplanorbulinella* sp., axial section, showing its very high conical test with equatorial chambers and coarser structure of lateral chambers in the umbilical cavity (HYM-36A/10/4); **J-M**, *Austrotrillina brunni* Marie, 1955, **J**, axial section (HYM-35/1/1); **K**, subaxial section (HYM-36A/3/1); **L**, axial section, megalospheric form (HYM-36A/1); **M**, subaxial section, (HYM-36A/10/1); **N**, *Planorbulina bronnimanni* Bignot & Decrouez, 1982,



axial section, showing protoconch followed by planispiral chambers (HYM-36A/6/1); **O**, *Borelis merici* Sirel & Gündüz, 1981, non-centered axial section (HYM-36A/5/4); **P**, *Borelis pygmaea* (Hanzawa, 1930), axial section, megalospheric form (HYM-36A/1/1); **Q-R**, *Penarchaias glynnjonesi* (Henson, 1950), axial section, megalospheric forms showing protoconch followed by planispiral chambers and chambers with uncoiled stage (HYM-36A/3 and HYM-36A). Scale bar: A-I, O-S, 0.25 mm; J-N, 0.5 mm.

ally from the apex to the base. The diameter of umbilical cavity is 0.42 mm, which is filled by numerous lateral chambers.

COMPARISON

Neoplanorbulinella sp. resembles *N. malatyaensis* n. sp. and *N. matsumarui* n. sp. in general appearance, but is distinguished by having very high conical test and larger size. In addition, it differs from *N. malatyaensis* n. sp. and *N. matsumarui* n. sp. in possessing numerous equatorial chambers and coarser structure of lateral chambers in the umbilical cavity. *Neoplanorbulinella* sp. is found in the Karamağara section together with *N. matsumarui* n. sp. and *N. malatyaensis* n. sp.

Genus *Planorbulina* d'Orbigny, 1826

TYPE SPECIES. — *Planorbulina mediterraneensis* d'Orbigny, 1826 by original designation.

Planorbulina bronnimanni Bignot & Decrouez, 1982 (Fig. 6N)

Planorbulina bronnimanni Bignot & Decrouez, 1982: 144, pl. 1, figs 1-9; pl. 2, figs 1-5; pl. 3, figs 1-3; plate 4, figs 1-6, pl. 5, figs 1-9; plate 6, figs 1-8. — Sirel & Acar 1993: 181, 183, pl. 2, fig. 21.

DESCRIPTION

The test is discoidal with calcareous wall and its diameter varies between 0.55-1.89 mm. It is dimorphic and there are chambers in megalospheric forms with trochospiral whorls after the proloculus. The following chambers are irregularly arranged in different sizes. As it is clearly seen in well-oriented sections, the connection between chambers is provided by the basal stolons.

Suborder MILIOLINA Delage & Herouard, 1896 Family SORITIDAE Ehrenberg, 1839 Subfamily ARCHAISINAE Cushman, 1927

Genus *Penarchaias* Hottinger, 2007

TYPE SPECIES. — *Peneroplis glynnjonesi* Henson, 1950 by original designation.

Penarchaias glynnjonesi (Henson, 1950) (Fig. 6Q, R)

Peneroplis glynnjonesi Henson, 1950: 35, pl. 9, figs 8, 9.

Peneroplis aff. *glynnjonesi* – Sirel 2004: 36, pl. 35, figs 2, 3, 5, 10.

Penarchaias glynnjonesi – Hottinger 2007: 11, 12, pl. 1, fig. 3; pl. 6, figs 2, 10; pl. 7, figs 7, 8; pl. 8, figs 6, 10; pl. 9, figs 6; pl. 12, figs 2; pl. 13, figs 3; pl. 14, figs 13; pl. 15, figs 1-5, 9.

DESCRIPTION

Only megalospheric form is observed. The test is lenticular, its diameter 0.9 mm in early stage, while in adult forms the

diameter and the thickness of the test ranges from 1.31 to 1.36 mm and 0.47 to 0.52 mm, respectively. Spheric megalosphere is followed by planispiral early chambers, and later chambers become uniserial. Interior skeletal apertures are observed on the septum (Hottinger 2007: pl. 7, figs 7, 9). These apertures, which are one of the most important characteristics of the genus, are well observed in Figure 6R, S.

Family AUSTROTRILLINIDAE Loeblich & Tappan, 1986

Genus *Austrotrillina* Parr, 1942

TYPE SPECIES. — *Trillina howchini* Schlumberger, 1893 by original designation.

Austrotrillina brunni Marie, 1955 (Fig. 6J-M)

Austrotrillina brunni Marie, 1955: 203, pl. 9, figs 4-8. — Adams 1968: 85, pl. 6, figs 6, 8. — Sirel 2003: 294, pl. 10, figs 10-16.

DESCRIPTION

Megalospheric form

The test is small and the peripheral margin is rounded in transverse sections (Fig. 6L). The length of the test ranges from 0.62 to 0.75 mm and the width from 0.39 to 0.49 mm. The spheric megalosphere (0.11 to 0.19 mm in diameter) is followed by small undivided chambers arranged in triloculine mode. Later adult chambers with fine subepidermal partitions are also lined up in triloculine pattern. Two types of subepidermal partitions form small alveolar compartments.

Microspheric form

The small test is triangular with rounded margin in transverse sections. The length of this species is unknown, the width of the test is 0.8 mm. Very small, spherical microsphere is followed by the early chambers arranged in quinqueloculine mode, later chambers arranged in triloculine pattern.

Family ALVEOLINIDAE Ehrenberg, 1839

Genus *Borelis* de Montfort, 1808

TYPE SPECIES. — *Nautilus melo* Fichtel & Moll, 1798 by original designation.

Borelis pygmaea (Hanzawa, 1930) (Fig. 6P)

Borelis (Fasciolites) pygmaea Hanzawa, 1930: 94, pl. 26, figs 14, 15.

Nealveolina pygmaea – Bakx 1932: 237, pl. 3, figs 18, 19. — Bursch 1947: 28, pl. 1, figs 11, 15, 19. — Adams 1965: 25.

Borelis pygmaea – Sirel 2003: 298, pl. 11, figs 1-7.

DESCRIPTION

The specimens of the megalospheric form are medium sized and fusiform. Their axial diameter is 2.10 mm and equatorial diameter 0.68 mm at the 6th whorls. Index of elongation is 3.08. The small spherical megalosphere (its diameter is 0.65 mm; Fig. 6P) is followed by tightly coiled four or five and loosely arranged adult whorls. The whorls are coiled tightly at the equatorial region, but they are arranged loosely at axial region, so that the axial thickening becomes higher at the poles of the test. The subspheric to oval chamberlets are arranged tightly and their size becomes large at the axial regions.

Borelis merici Sirel & Gündüz, 1981
(Fig. 6O)

Borelis merici Sirel & Gündüz, 1981: 73, 74, pl. 1, figs 9-13. — Sirel 2003: 299, pl. 11, figs 8, 9.

DESCRIPTION

The test is slightly elongated oval with an axial diameter of 1.01-1.41 mm and equatorial diameter of 0.45-0.78 mm. Index of elongation is between 1.79-2.71. The first 4-5 whorls which follow the proloculus are devoid of an axial thickening, so that the shape of the test seems spheric in this stage. Axial thickening increases at the last 4 whorls, therefore the shape of the test becomes elongated oval in the adult stage. Dimorphism is faint.

CONCLUSION

The genus *Neoplanorbulinella* Matsumaru, 1976 was firstly identified in the Lower Miocene of Saipan and Mariana Islands in the Western Pacific. It is discovered for the first time in the Lower Oligocene deposits of Malatya Basin in eastern Anatolia (Turkey), and is not known there from older or younger strata. The present study provides the description of two new species from the Karamağara stratigraphic section, *Neoplanorbulinella matsumarui* n. sp. and *Neoplanorbulinella malatyaensis* n. sp. *N. matsumarui* n. sp. differs from the type species *N. saipanensis* Matsumaru, 1976 in being characteristically small, and in having high conical test with almost right apical angle. The other species *N. malatyaensis* n. sp. has a larger, low conical to lenticular test. In addition, the associated foraminiferal taxa are listed and the key species are described and illustrated (Fig. 6J-S).

Acknowledgements

The author is grateful to Dr. Aynur Hakyemez and Aysegül Aydın (MTA, Ankara) for identification of planktonic foraminiferal and calcareous nannoplankton assemblages, respectively, and to Sevkettin Sen (MNHN, Paris) for advice on the presentation of the manuscript. Special thanks go to Dr Ercüment Sirel (Ankara University), for his support, advice and encouragements he has given me during the studying process. The author also likes to thank two anonymous reviewers, who greatly improved the manuscript.

REFERENCES

- ADAMS C. G. 1965. — The Foraminifera and stratigraphy of the Melinau Limestone, Sarawak and its importance in Tertiary correlation. *Quarterly Journal of the Geological Society of London* 121: 283-338. <https://doi.org/10.1144/gsjgs.121.1.0283>
- ADAMS C. G. 1968. — A revision of the foraminiferal genus *Austrorillina* Parr. *Bulletin of the British Museum (Natural History), Geology* 16: 73-97.
- AKKUŞ M. F. 1971. — Darende-Balaban Havzasının (Malatya, ESE Anadolu) jeolojik ve stratigrafik incelenmesi [Stratigraphic and geological investigations of Darende-Balaban Basin (Malatya, ESE Anatolia)]. *Maden Tetkik ve Arama Dergisi* 76: 1-60.
- ALKAN H. 1997. — *Malatya Baseninin Jeolojisi ve Petrol Olanakları* [The geology of Malatya Basin and its petrol opportunities]. TPAO Reports 3766, Ankara (unpublished).
- AYAN T. 1961. — *Malatya kuzeyindeki Hekimhan-Ebreme köyü bölgesinin (K39-c3) detay jeolojisi ve petrol imkanları* [Detailed geology and petrol opportunities of Hekimhan-Ebreme village (North Malatya)]. Maden Tetkik ve Arama Raporu 4186, Ankara, (unpublished).
- BAKX L. A. J. 1932. — De genere *Fasciolites* en *Neoalveolina* in het Indo-Pacifische gebied. *Verhandelingen Geologisch Mijnbouwkundig Genootschap voor Naderland en Kolonien, Geologische serie* 9: 205-266.
- BASSI D., HOTTINGER L. & NEBELSICK J. H. 2007. — Larger Foraminifera from the Upper Oligocene of the Venetian Area, North-East Italy. *Palaeontology* 50: 845-868. <https://doi.org/10.1111/j.1475-4983.2007.00677.x>
- BIGNOT G. & DECROUEZ D. 1982. — Un Planorbulinidae (Foraminifera) nouveau du Priabonien et de l'Oligocène de l'Europe méridionale et occidentale. *Revue de Paléobiologie* 1 (2): 141-163.
- BİLGİÇ T. 2002. — 1/500 000 ölçekli Türkiye Jeoloji Haritaları Serisi, Sivas Paftası [1/500 000 scale geology maps series of Turkey]. Maden Tetkik ve Arama Genel Müdürlüğü yayınları, Ankara.
- BOZKURT E. 2001. — Neotectonics of Turkey – a synthesis. *Geodinamica Acta* 14: 3-30. <https://doi.org/10.1080/09853111.2001.11432432>
- BOUDAGHER-FADEL M. K. 2008. — *Evolution and geological significance of larger benthic foraminifera*. Developments in Palaeontology and Stratigraphy, Volume 21, Elsevier, Amsterdam, 571 p.
- BURSCH J. G. 1947. — Micropaläontologische Untersuchungen des Tertiärs von Gros Key (Molukken). *Mémoires suisses de Paléontologie* 65: 1-69.
- ÇAĞLAR (KAYA) M. 2009. — Benthic foraminiferal biostratigraphy of the Tertiary sediments from the Elazığ and Malatya basins, Eastern Turkey. *Journal of the Geological Society of India* 74 (2): 209-222. <https://doi.org/10.1007/s12594-009-0123-8>
- DELAGE Y. & HÉROUARD É. 1896. — *Traité de Zoologie concrète*, Vol. 1, La cellule et les protozoaires. Schleicher Frères, Paris. <https://doi.org/10.5962/bhl.title.11672>
- DROOGER C. W. 1993. — Radial foraminifera; morphometrics and evolution. *Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen* 41: 1-242. <https://doi.org/10.2113/gsjfr.24.4.312>
- GEDİK F. 2010. — *Benthic Foraminiferal Description and Biostratigraphy of Oligo-Miocene Shallow-Water Sediments of Malatya Basin*. PhD Thesis, 184 p., 25 pl., Ankara (unpublished).
- GEDİK F. 2014. — Benthic foraminiferal fauna of Malatya Oligo-Miocene Basin (Eastern Taurids, Eastern Turkey). *Bulletin of the Mineral Research and Exploration* 149: 95-139. <https://doi.org/10.19111/bmre.91054>
- GEDİK F. 2015. — Benthic foraminiferal biostratigraphy of Malatya Oligo-Miocene succession (Eastern Taurids, Eastern Turkey). *Bulletin of the Mineral Research and Exploration* 150: 19-50.
- HANZAWA S. 1930. — Note on Foraminifera found in the Lepidocyclina-limestone from Pabehasan, Java. *Science Reports of the Tohoku Imperial University, 2nd series, Geology* 14 (1): 85-96.

- HENSON F. R. S. 1948a. — *Larger Imperforate Foraminifera of South-Western Asia*. Order of the Trustees of the British Museum, London, 126 p., 16 figs, 16 pls.
- HENSON F. R. S. 1948b. — Preliminary note on the structure of the foraminiferal genus *Archaias* and related forms. *Quarterly Journal of the Geological Society* 104 (1-4): 323-324. <https://doi.org/10.1144/GSL.JGS.1948.104.01-04.13>
- HENSON F. R. S. 1950. — *Middle Eastern Tertiary Peneroplidae (Foraminifera) with Remarks on the Phylogeny and Taxonomy of the Family*. PhD Thesis, Leiden University, West Yorkshire Printing Co., Wakefield: 1-70, 10 pls.
- HOTTINGER L. 2007. — Revision of the foraminiferal genus *Globoreticulina* Rahaghi, 1978, and of its associated fauna of larger foraminifera from the late Middle Eocene of Iran. *Carnets de Géologie/Notebooks on Geology*: article 2007/06.
- KARADENİZLİ L., VAROL B., SARAÇ G. & GEDİK F. 2016. — Late Eocene-Early Miocene palaeogeographic evolution of Central Eastern Anatolian basins, the closure of the Neo-Tethys Ocean and continental collision. *Journal of the Geological Society of India* 88 (6): 773-798. <https://doi.org/10.1007/s12594-016-0545-z>
- KARAMAN T., POYRAZ N., BAKIRHAN B., ALAN İ., KADINKIZ G., YILMAZ H. & KILINÇ F. 1993. — *Malatya-Doğanşehir-Çelikhan dolayının jeolojisi* [Geologic vicinity of Malatya-Doğanşehir-Çelikhan]. Maden Tetkik ve Arama Raporu, 9587, Ankara (unpublished).
- KURTMAN F. 1978. — Gürün bölgesinin jeolojisi ve tektonik özellikleri [The geology and tectonic specialities of Gürün region]. *Maden Tetkik ve Arama Dergisi* 91: 1-12.
- LOEBLICH A. R. & TAPPAN H. 1987. — Foraminiferal genera and their classification. Van Nostrand Reinhold, New York, vol. I: 1-970; vol. II: 1-212, 847 pls. <https://doi.org/10.1007/978-1-4899-5760-3>
- MARIE P. 1955. — Quelques formes nouvelles de polypiers et de foraminifères de l'Oligocène et du Miocène du NW de la Grèce. *Bulletin de la Société géologique de France* 6 (5): 193-205.
- MATSUMARU K. 1976. — Larger foraminifera from the islands of Saipan and Guam, Micronesia, in TAKAYANAGI Y. & SAITO T. (eds), *Progress in Micropaleontology*. Microplaleontology Press Special Publication, New York, American Museum of Natural History: 190-213.
- MATSUMARU K. 1996. — *Tertiary Larger Foraminifera (Foraminifera) from the Ogasawara Islands, Japan*. Paleontological Society of Japan, vol. 36, 239 p.
- ÖRÇEN S. 1986. — Medik-Ebreme (KB Malatya) dolayının biyostratigrafisi ve paleontolojisi [Medik-Ebreme's (NW Malatya) Palaeontology and Biostratigraphy]. *Maden Tetkik ve Arama Dergisi* 105/106: 39-68.
- RÖGL F. 1999. — Mediterranean and Paratethys Palaeogeography during the Oligocene and Miocene, in AGUSTI J., ROOK L. & ANDREWS P. (eds), *The Evolution of Neogene Terrestrial Ecosystems in Europe*. Cambridge University Press, Cambridge: 8-22.
- SİREL E. 2003. — Foraminiferal description and biostratigraphy of the Bartonian, Priabonian and Oligocene shallow-water sediments of the southern and eastern Turkey. *Revue de Paléobiologie* 22 (1): 269-339.
- SİREL E. 2004. — Türkiye'nin Mesozoyik ve Senozoyik yeni bentik Foraminiferleri [Turkey's Mesozoic and Cenozoic new benthic foraminifers]. *Jeoloji Mühendisleri Odası Yayınları/Chamber of Geological Engineers of Turkey* 84: 1-219.
- SİREL E. & ACAR S. 1993. — *Malatyna*, a new foraminiferal genus from the Lutetian of Malatya region (East Turkey). *Geologia croatica* 46 (2): 181-188.
- SİREL E. & GEDİK F. 2011. — *Postmiogypsinella*, a new Miogypsinidae (Foraminifera) from the late Oligocene in Malatya Basin, Turkey. *Revue de Paléobiologie* 30 (2): 591-603.
- SİREL E. & GÜNDÜZ H. 1981. — Description of new *Borelis* species from the Hatay (S of Turkey) and Elazığ region (E of Turkey). *Bulletin of the Mineral Research and Exploration Institute of Turkey* 92: 70-74.
- ŞENGÖR A. M. C. & YILMAZ Y. 1981. — Tethyan evolution of Turkey: a plate tectonic approach. *Tectonophysics* 75: 181-241. [https://doi.org/10.1016/0040-1951\(81\)90275-4](https://doi.org/10.1016/0040-1951(81)90275-4)
- YOLDAŞ R. 1972. — *Malatya kuzeyinin jeolojisi ve petrol olanakları* [The geological and petroleum opportunities of Northern Malatya]. Maden Tetkik ve Arama Raporu 4936, Ankara (unpublished).

Submitted on 2nd May 2016;
accepted on 10 November 2016;
published on 30 June 2017.