

Erratum: volume 41 (6) 2020: 119-132, May 4, 2020.

***Kwanghwana miscanthi* Karun., C.H.Kuo & K.D.Hyde,
gen. et sp. nov. (Phaeosphaeriaceae, Pleosporales)
on *Miscanthus floridulus* (Labill.) Warb.
ex K.Schum. & Lauterb. (Poaceae)**

Anuruddha KARUNARATHNA

Department of Plant Medicine, National Chiayi University,
300 Syuefu Road, Chiayi City 60004 (Taiwan)
and Centre of Excellence in Fungal Research, Mae Fah
Luang University, Chiang Rai 57100 (Thailand)
and Department of Entomology and Plant Pathology, Faculty of Agriculture,
Chiang Mai University, Chiang Mai 50200 (Thailand)
and Key Laboratory for Plant Diversity and Biogeography of East Asia,
Kunming Institute of Botany, Chinese Academy of Sciences,
132 Lanhei Road, Kunming 650201 (China)
and World Agroforestry Centre, East and Central Asia,
132 Lanhei Road, Kunming 650201, Yunnan (China)

Rungtiwa PHOOKAMSAK

Key Laboratory for Plant Diversity and Biogeography of East Asia,
Kunming Institute of Botany, Chinese Academy of Sciences,
132 Lanhei Road, Kunming 650201 (China)
and World Agroforestry Centre, East and Central Asia,
132 Lanhei Road, Kunming 650201, Yunnan (China)

Ruvishika S. JAYAWARDENA

Centre of Excellence in Fungal Research, Mae Fah Luang University,
Chiang Rai 57100 (Thailand)

Kevin D. HYDE

Centre of Excellence in Fungal Research, Mae Fah
Luang University, Chiang Rai 57100 (Thailand)
and Key Laboratory for Plant Diversity and Biogeography of East Asia,
Kunming Institute of Botany, Chinese Academy of Sciences,
132 Lanhei Road, Kunming 650201 (China)
and World Agroforestry Centre, East and Central Asia,
132 Lanhei Road, Kunming 650201, Yunnan (China)

Chang H. KUO

Department of Plant Medicine, National Chiayi University,
300 Syuefu Road, Chiayi City 60004 (Taiwan)
chkuo@mail.ncyu.edu.tw (corresponding author)

Published on 17 June 2020

Karunaratna A., Phookamsak R., Jayawardena R. S., Hyde K. D. & Kuo C. H. 2020. — Erratum: volume 41 (6) 2020: 119-132, 4 May 2020. *Kwanghwana miscanthi* Karun., C.H.Kuo & K.D.Hyde, gen. et sp. nov. (Phaeosphaeriaceae, Pleosporales) on *Miscanthus floridulus* (Labill.) Warb. ex K.Schum. & Lauterb. (Poaceae). *Cryptogamie, Mycologie* 41 (9): 157-160. <https://doi.org/10.5252/cryptogamie-mycologie2020v41a9>. <http://cryptogamie.com/mycologie/41/9>

Aux pages 121, 124 et 126, l'identifiant holotype donné est incorrect. Il a été remplacé par MFLU 18-1952. La Figure 1 de la page 125 est également incorrecte. Ici, nous l'avons remplacée par la figure correcte. De même, l'identificateur d'holotype dans la légende de la Figure 2 a été corrigé.

In pages 121, 124 and 126, the given holotype identifier is incorrect. They were replaced by MFLU 18-1952. Figure 1 on page 125 is also incorrect. Here, we replaced it with the correct figure. Also, the holotype identifier in the legend of Figure 2 has been corrected.

TAXONOMY

In this section, the new genus (*Kwanghwana* Karun., C.H.Kuo & K.D.Hyde, gen. nov.) is described and the new species (*Kwanghwana miscanthi* Karun., C.H.Kuo & K.D.Hyde, sp. nov.) are illustrated and described from holotype (holo-, MFLU[MFLU 18-1952]), and detailed illustrations are provided for the new species.

Genus *Kwanghwana* Karun., C.H.Kuo & K.D.Hyde

Kwanghwana miscanthi Karun., C.H.Kuo &
K.D.Hyde, sp. nov.
(Fig. 2)

INDEX FUNGORUM NUMBER. — IF557101.

FACESOFFUNGI NUMBER. — FoF 07776.

ETYMOLOGY. — Refers to the generic name for the host.

HOLOTYPE. — MFLU 18-1952.

CULTURE CHARACTERS. — Colonies on PDA slow growing at 20-25°C, circular, white at the growing edge, light to dark pink in the centre; reverse white at the edge, the centre dense, convex or dome-shaped to umbonate, with entire edge, glabrous with tiny granular on surface, no pigment produced.

MATERIAL EXAMINED. — **Taiwan**, Chiayi Province, Kwang Hwa, alt. 1330 m, 23°30.084'N, 120°41.73.5'E on *Miscanthus floridulus* (Poaceae) leaf sheath, 17.III.2018, A. Karunarathna, AKTW 05 (holo-, MFLU[MFLU 18-1952]), ex-type living culture, FU31017, NCYUCC 19-0372.

Published on 17 June 2020.




Fig. 1. — Phylogram generated from RAxML analysis based on a combined dataset of LSU, SSU, ITS and TEF1- α partial sequences. Bootstrap support values for maximum likelihood higher than 60% and Bayesian posterior probabilities (BPP) greater than 0.90 are displayed as above the nodes respectively. The ex-type strains are in bold. New strain is indicated in red. The tree is rooted to *Leptosphaeria dolium* (Pers.) Ces. & De Not. (CBS 505.75) and *Paraleptosphaeria dryadis* (Johanson) Gruyter, Aveskamp & Verkley (CBS 643.86).




FIG. 2.— *Kwanghwana miscanthi* Karun., C.H.Kuo & K.D.Hyde, sp. nov. (MFLU 18-1952, holotype): **A, B**, appearance of ascomata on the host *Miscanthus floridulus* (Labill.) Warb. ex K.Schum. & Lauterb. (Poaceae) leaf sheath; **C**, section through ascoma; **D**, section through peridium at the apex; **E**, section through peridium at the base; **F**, hamathecium; **G**, Pseudoparaphyses; **H-K**, asci; **L-P**, ascospores; **Q**, germinating ascospore; **R, S**, culture characteristics on PDA (r = from above, s = from below). Scale bars: C, 20 µm; D-F, 25 µm; G-K, 20 µm; L-Q, 25 µm.