

Annotated checklist of the Hemiptera Heteroptera
of the Site of Community Importance and Special
Area of Conservation “Alpi Marittime” (NW Italy)

Margherita NORBIATO,
Paride DIOLI & Fabio CIANFERONI

DIRECTEUR DE LA PUBLICATION : Bruno David
Président du Muséum national d'Histoire naturelle

RÉDACTRICE EN CHEF / *EDITOR-IN-CHIEF* : Laure Desutter-Grandcolas

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS* : Anne Mabilie (zoosyst@mnhn.fr), Emmanuel Côté

MISE EN PAGE / *PAGE LAYOUT* : Anne Mabilie

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD* :

James Carpenter (AMNH, New York, États-Unis)
Maria Marta Cigliano (Museo de La Plata, La Plata, Argentine)
Henrik Enghoff (NHMD, Copenhagen, Danemark)
Rafael Marquez (CSIC, Madrid, Espagne)
Peter Ng (University of Singapore)
Norman I. Platnick (AMNH, New York, États-Unis)
Jean-Yves Rasplus (INRA, Montferrier-sur-Lez, France)
Jean-François Silvain (IRD, Gif-sur-Yvette, France)
Wanda M. Weiner (Polish Academy of Sciences, Cracovie, Pologne)
John Wenzel (The Ohio State University, Columbus, États-Unis)

COUVERTURE / *COVER* :

Eurydema fieberi Fieber, 1837. Photo: S. Bambi.

Zoosystema est indexé dans / *Zoosystema is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Zoosystema est distribué en version électronique par / *Zoosystema is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Zoosystema* sont référencés par /
Articles and nomenclatural novelties published in Zoosystema are referenced by:

- ZooBank® (<http://zoobank.org>)

Zoosystema est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris / *Zoosystema is a fast track journal published by the Museum Science Press, Paris*

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:

Adansonia, Geodiversitas, Anthropolozologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Algologie, Bryologie, Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)
Tél. : 33 (0)1 40 79 48 05 / Fax : 33 (0)1 40 79 38 40
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2019
ISSN (imprimé / *print*) : 1280-9551 / ISSN (électronique / *electronic*) : 1638-9387

Annotated checklist of the Hemiptera Heteroptera of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (NW Italy)

Margherita NORBIATO

Natural History Museum of the University of Florence, Zoological Section “La Specola”,
Via Romana 17, I-50125 Florence (Italy)
margherita.norbiato@libero.it

Paride DIOLI

Natural History Museum, Corso Venezia 55, I-20121 Milan (Italy)
paridedioli@virgilio.it

Fabio CIANFERONI

Natural History Museum of the University of Florence, Zoological Section “La Specola”,
Via Romana 17, I-50125 Florence (Italy)
and Research Institute on Terrestrial Ecosystems, CNR – National Research Council of Italy,
Via Madonna del Piano 10, I-50019 Sesto Fiorentino, Florence (Italy)
cianferoni.fabio@gmail.com, fabio.cianferoni@unifi.it

Submitted on 10 January 2018 | accepted on 28 September 2018 | published on 9 July 2019

urn:lsid:zoobank.org:pub:6B6BCC2F-9968-4D52-AA99-C24BAF56C360

Norbiato M., Dioli P. & Cianferoni F. 2019. — Annotated checklist of the Hemiptera Heteroptera of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (NW Italy). *Zoosystema* 41 (15): 269-312. <https://doi.org/10.5252/zoosystema2019v41a15>. <http://zoosystema.com/41/15>

ABSTRACT

We provide the first annotated checklist of the Heteroptera (Insecta: Hemiptera) of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (IT1160056), based on literature records and unpublished material collected between 2012-2015 within two European projects in which the Alpi Marittime Nature Park was involved. The literature review provided a list of 62 species, recorded for the study area between 1884 and 1990, scattered in 14 papers. The new data here provided raise the total number of species occurring in this region to 208, grouped in 25 families and 135 genera. For each species reported in the catalogue, previous literature records, new data and the known distribution are provided. Among the most interesting findings, *Salda henschii* (Reuter, 1891) which was recently recorded for the first time in Italy, and *Sternodontus obtusius* Mulsant & Rey, 1856 which is rediscovered in the Piedmont region after more than a half century.

RÉSUMÉ

Liste commentée des Hemiptera Heteroptera du Site d'importance communautaire et Zone spéciale de conservation «Alpi Marittime» (Nord Ouest de l'Italie).

La première liste raisonnée des Hétéroptères du Site d'Importance Communautaire et Zone spéciale de Conservation «Alpi Marittime» (IT1160056) est fournie, basée à la fois sur les données de la littérature et sur du matériel inédit récolté pendant la période 2012-2015 dans le cadre de deux projets européens dans lesquels le Parc naturel des «Alpi Marittime» était impliqué. La recherche

KEY-WORDS

Heteroptera,
Maritime Alps,
SW-Alps,
Piedmont,
Italy,
faunistics,
new records.

MOTS CLÉS
Heteroptera,
Alpes Maritimes,
Alpes nord-occidentales,
Piémont,
faunistique,
signalisations nouvelles.

PAROLE CHIAVE
Heteroptera,
Alpi Marittime,
Alpi nord-occidentali,
Piemonte,
faunistica,
nuove segnalazioni.

bibliographique a permis de lister 62 espèces sur la zone d'étude, signalées entre 1884 et 1990 dans 14 publications. Les nouvelles données font passer le nombre des espèces connues à 208, regroupées en 25 familles et 135 genres. Pour chaque espèce mentionnée dans le catalogue sont incluses les données bibliographiques, les nouvelles signalisations et la distribution géographique connue. Parmi les découvertes les plus intéressantes, on peut citer *Salda henschii* (Reuter, 1891), récemment signalée pour la première fois en Italie et *Sternodontus obtusus* Mulsant & Rey, 1856, qui n'avait pas été retrouvée dans le Piémont depuis plus d'un demi-siècle.

RIASSUNTO

Checklist ragionata degli Hemiptera Heteroptera del Sito di Importanza Comunitaria e Zona Speciale di Conservazione "Alpi Marittime" (Italia Nord Occidentale).

Viene fornita la prima checklist ragionata degli eterotteri del Sito di Importanza Comunitaria e Zona Speciale di Conservazione "Alpi Marittime" (IT1160056), basata sia su dati di letteratura che su materiale inedito raccolto nel periodo 2012-2015 nell'ambito di due progetti europei in cui il Parco Naturale delle Alpi Marittime era coinvolto. La ricerca bibliografica ha consentito di elencare 63 specie per l'area di studio, segnalate tra il 1884 e 1990 e suddivise in 14 pubblicazioni. I nuovi dati riportati fanno salire il numero di specie conosciute a 208, appartenenti a 25 famiglie e 135 generi. Per ogni specie inserita nel catalogo vengono forniti dati bibliografici, nuove segnalazioni e distribuzione geografica. Tra le specie più interessanti, *Salda henschii* (Reuter, 1891), recentemente segnalata per la prima volta in Italia, e *Sternodontus obtusus* Mulsant & Rey, 1856, ritrovato in Piemonte dopo più di mezzo secolo.

INTRODUCTION

The Maritime Alps region has been defined as a key area for understanding the dynamics that have shaped the fauna of the Italian peninsula as well as that of western Europe in general, and this is due to the peculiar faunistic diversity (Minelli *et al.* 2006). The reason for this extraordinary biodiversity is related to the particular biogeographical frame in which the area is located, between the mountain systems of Piedmont and Liguria regions and the Mediterranean Sea, straddling the border between the Mediterranean, Alpine and continental zones. In the NW part of Italy, the mountain chain peaks are located close to the seaside and this fact contributes to create very steep ecological gradients both in the northern Ligurian maritime slope and in the southern Piedmontese continental one. Moreover, the wide altitudinal range (650 to 3297 m a.s.l.) and the geological and glacial history of the area contribute significantly in making the Maritime Alps one of the most important European hot spots of biodiversity (Bonelli *et al.* 2015, Isaia *et al.* 2015). The Maritime Alps, the southernmost part of the Alpine chain, divide the Piedmontese plains from the Côte d'Azur and are between two very popular crossings since antiquity: the Colle di Tenda and the Colle della Maddalena. Both sides of the Maritime Alps are subject to protection: actually, on the French side, there is the Mercantour National Park; on the Italian side, the Alpi Marittime Nature Park. The two Parks border over 35 kilometers and together form a protected area of over 100 000 hectares.

In the four-year period 2012-2015, different entomological research has been carried out, mainly by the first author, in the Site of Community Importance (SCI) and Special Area of Conservation (SAC) "Alpi Marittime" (IT1160056). Most

of the material was collected within the framework of two different European projects. The "ATBI + M" (All Taxa Biodiversity Inventory + Monitoring) project, coordinated by the European Distributed Institute of Taxonomy (EDIT), funded in 2006 in the frame of the Sixth Framework Programme of the European Commission (Deharveng *et al.* 2015; Villemant *et al.* 2015). All Taxa Biodiversity Inventories (ATBI) are intensive community efforts to identify and record all living species that exist within a given area. The second was the Interreg ALCOTRA 2007-2013 "Gestione comune del paesaggio transfrontaliero delle Alpi Meridionali" ("Joint development of the cross-border landscape of the Southern Alps"). Further material from other collectors was also added.

This contribution represents the results of the research conducted mainly from 2012 and 2015 within the Site of Community Importance and Special Area of Conservation IT1160056 "Alpi Marittime" (IT1160056) and it also provides the first checklist of the Heteroptera for this area. Based on intense field activity and on an extensive bibliographic search, the work aims to increase the knowledge of the Heteropteran fauna of this study area and to provide a basis for future studies of this group of insects.

MATERIAL AND METHODS

STUDY AREA

The Site of Community Importance and Special Area of Conservation "Alpi Marittime" (IT1160056) (Fig. 1), located in the Cuneo province, is the largest protected area in Piedmont. It covers a surface of 33 673 hectares and includes the municipalities of Aisone, Entracque, Limone Piemonte,




Fig. 1. — Study area in NW Italy (Site of Community Importance and Special Area of Conservation “Alpi Marittime” IT1160056).

Roaschia, Valdieri, Vernante, and includes part of the Valle Stura (Demonte), almost all the Valle Gesso, the upper part of Valle Vermentagna (Colle di Tenda) and the upper Val Grande (Palanfrè). The altitudinal range goes from about 700 m to the 3297 m of the Maritime Alps highest peak, the “Cima Argentera”. The habitats found in the area vary with the elevation: at the bottom of the valleys there are hardwood forests (mainly beech forests – *Fagus sylvatica* L.), followed by small spruce and larch forests; increasing the altitude we can find large areas of shrubs and bushes, and finally the open habitats of the higher areas are composed by grasslands, pastures and rocky lands. The history of the current protected area began with the unification of two protected areas instituted in 1979 and 1980: respectively, the “Parco Naturale dell’Argentera” and the “Riserva naturale speciale Boschi e Laghi di Palanfrè”, this incorporation created the so-called “Parco Naturale delle Alpi Marittime” in 1995, covering a surface of 28 000 hectares. In 2010 there has been a new delimitation of the Sites of Community Importance and Special Areas of Conservation in the Piedmont region, and this has led to increase the authority of the Park, which now has been expanded to the whole surface included within the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (IT1160056). In 2016, the “Parco Naturale delle Alpi Marittime” has been included in the “Ente di Gestione delle Aree Protette delle Alpi Marittime” (management body), which includes several protected areas, including the area studied in this work.

In this paper we consider all the territories within the SCI and SAC areas, including the “Parco Naturale delle Alpi Marittime”. Considering their closeness to the study area, several localities cited in literature and falling just outside the borders, were also included in this work.

This research was mainly carried out in 10 sites (Table 1) located within the protected area. These areas consist on different macrohabitats: forests, grasslands and pastures, wetlands and aquatic environments. In Table 1 a description of the investigated areas is provided.

SOURCES

The fieldwork was preceded by an extensive bibliography search. Neither for the Site of Community Importance, Special Area of Conservation “Alpi Marittime” (IT1160056), nor for the “Parco Naturale delle Alpi Marittime”, a complete faunistic study of the Heteroptera has been carried out until now.

The main source used about Italian Heteroptera is the volume of the Fauna d’Italia by Servadei (1967). To date, though the volume is outdated, it represents the most complete bibliographic source about this group of insects for Italy. In order to identify any references reported for the territory taken into consideration by this search, we checked all the literature indicated by Servadei (1967) for “Piedmont”. Then we consulted a hundred works, of which only 14 reported data locations falling within the borders of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (IT1160056).

TABLE 1. — List of the collecting sites.

Exposure	Altitudinal range	Municipality	Area
N; NE; E	850-1360 m	Entracque	Fus Carbonaro forest
W	1450-1550 m	Valdieri	Valletta forest
E; W	1180-1200 m	Entracque	Suffiet bridge
SW	980 m	Valdieri	Sant'Anna di Valdieri and Gesso stream
N; N-E	1600-2050 m	Limone Piemonte	Vallone della Perla (Fig. 2A, B)
E-S-E; N; N-E	1600-2000 m	Valdieri	Colle dell'Arpione (Fig. 2C)
S-E; E	1500-2200 m	Vernante	Palanfrè (Fig. 2D, E)
E; W	1680-1740 m	Valdieri	Pian della Casa and Gias della Casa (Fig. 2F)
S; NW	1400-1800 m	Valdieri	Pian del Valasco (Fig. 2G)
W	1400-2060 m	Entracque	Vej del Bouc lake (Fig. 2H)

Most of the bibliographic data originated from general works; in particular, the first records came from Ferrari (1884), who reported data of specimens collected by G. and L. Doria and in one case reports “Alpi Marittime”, although this is a doubtful record because we are not sure if the record fall within the protected area; however we decided to keep it in the list. Other old records come from Griffini (1893) who studied Pentatomidae Leach, 1815 and Coreidae Leach, 1815 of Piedmont region, including some data that fall within the current protected area; the last works about Heteroptera of the nineteenth century were those of Noelli (1897a; 1897b), who contributed to the knowledge of this group with his studies on Reduviidae Latreille, 1807 and Lygaeidae Schilling, 1829 of Piedmont region. In relatively more recent years, there have been works who focused on the analysis of some families of Italian Hemiptera and included data for Piedmont (e.g. Tamanini 1951, 1956); others, like those of Mancini (1952, 1959, 1963), consisted of “Miscellanea Emitterologica” with data from various Italian regions and, finally, a paper of Mancini (1963) on Heteroptera in Liguria, from which some useful data were found for Piedmont. In addition, two papers by Servadei (1955, 1957) on other Italian places provided records which can be attributed to the investigated area. Some records have been found also in the “Faune de France” volumes by Péricart (1972, 1983, 1987, 1990, 1999a, b, c). These are the main sources that gave most of the data.

The examination of literature provided, finally, a list of 63 species, scattered in 14 papers, published between 1884 and 1999.

In the list, in some sporadic cases, records of localities for which it was not clear if they would fall perfectly within the territory of the protected area were also added; these are very old records of the years prior to 1900. They were nevertheless considered, as it is most likely that they can actually fall in the territory concerned.

Some reports of Heteroptera regarding this territory were taken from the website of the European project ATBI (All Taxa Biodiversity Information), which was mentioned in the introduction (Deharveng *et al.* 2015; Villemant *et al.* 2015).

Finally, the examination of some old collections of the Museum of Natural History “Giacomo Doria” of Genoa allowed us to identify some unpublished data, which, once checked, have been reported in the list under the heading “unpublished data”. In addition, the visit to this museum allowed also to locate the specimens related to some old published records for the investigated area, that it was possible to confirm or correct.

Nomenclature and systematic follow the “Catalogue of the Heteroptera of the Palaearctic Region” (Aukema & Rieger 1995, 1996, 1999, 2001, 2006; Aukema *et al.* 2013), with the exception of the superfamily Lygaeoidea Schilling, 1829, for which we chose to follow Henry (1997). The checklist, at infraorder level, follows the taxonomic arrangement of Schuh & Slater (1995). Superfamilies, families and species are, however, in alphabetical order. For each species, the unpublished data and bibliographic ones are shown. For the new data we provide the following information: data of locality, geographical coordinates (WGS84), altitude metres a.s.l. (when available), date of collection, collector, number of specimens, and the collection in which the specimens are deposited. For each species, its original name of description, with author, year and the page of description is given. Then we indicate the chorotype according to Vigna Taglianti *et al.* (1993, 1999), followed by a paragraph “Remark”, in which additional information is provided, such as the presence of the species in Italy, if it is a new record for the study area or for Italy, ecological notes (e.g. host plants) or, in some cases, systematic notes.

Concerning the chorotypes, the Alpine one was not indicated in Vigna Taglianti *et al.* (1993, 1999), but we created it (eventually to abbreviate as ALPI) for the species *Horwathia lineolata* (A. Costa, 1852) and *Acalypta visolensis visolensis* Péricart, 1976, whose distribution range could not be associated with any of the categories mentioned in that paper.

The terrestrial Heteroptera (Cimicomorpha Leston, Pendergrast & Southwood, 1954; Pentatomomorpha Leston, Pendergrast & Southwood, 1954) have been identified by P. Dioli, the aquatic Heteroptera *s.l.* (Gerromorpha Popov, 1971; Nepomorpha Popov, 1968; Leptopodomorpha Popov, 1971) have been identified by F. Cianferoni.

ABBREVIATIONS

Collections

coll. AC	Attilio Carapezza collection, Palermo;
coll. AS	Antonio Servadei collection, Padua (now in the Natural History Museum of Verona);
coll. CR	Christian Rieger collection, Nürtingen;
coll. FC	Fabio Cianferoni collection, Florence;
coll. LT	Livio Tamanini collection, Rovereto (now in the Civic Museum of Rovereto);
coll. PD	Paride Dioli collection, Sondrio;
MZR	Zoology Museum of Rome;
MSNM	Natural History Museum of Milan;
MNHN	National History Museum of Paris;
MSNG	Natural History Museum “Giacomo Doria”, Genoa;
MSNV	Natural History Museum of Verona;
MZUF	Natural History Museum of the University of Florence, Zoological Section “La Specola”, Florence.



FIG. 2. — **A**, La Perla small lake; **B**, La Perla Valley; **C**, Pastures of the “Colle dell’Arpione”; **D**, Palanfrè forest; **E**, Pastures of Palanfrè; **F**, Pian della Casa and Gias della Casa small ponds; **G**, Pian del Valasco; **H**, Vej del Bouc lake. Photos: M. Norbiato.

Collectors abbreviations

CR	Christian Rieger;
FC	Fabio Cianferoni;
FT	Fabio Terzani;
LB	Luca Bartolozzi;
MD	Marco Dogliotti;
MN	Margherita Norbiato;
PD	Paride Dioli;
SR	Saverio Rocchi.

Other abbreviations

dx	right;
juv.	juvenile specimen;
leg.	legit/legerunt;
loc.	locality.

ANNOTATED CHECKLIST

Infraorder GERROMORPHA Popov, 1971
Superfamily GERROIDEA Leach, 1815
Family GERRIDAE Leach, 1815

Gerris (Gerris) costae (Herrich-Schäffer, 1850)

Hydrometra costae Herrich-Schäffer, 1850: 71.

LITERATURE. — Carapezza & Faraci (2005, 2007 [sub *Gerris (Gerris) costae costae*]): San Giacomo di Entracque, Retreat, 1800 m, 1964, MSNV.

NEW DATA. — Entracque, Vei del Bouc Lake, 2054 m, 44.14209°N, 7.42284°E (WGS84), 23.VIII.2013, MN, MD leg., 2 specimens juv., MZUF. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, FC, FT, PD, SR, MN, LB, MD leg., 4 ♂, 4 ♀, 2 juv., coll. FC. — Limone Piemonte, SE del Bric Campanino, little lakes, 2125 m, 44.153°N, 7.604°E (WGS84), 27.VII.2015, FC, FT, PD, SR, MN, LB, MD leg., 5 ♂, 8 ♀, 1 nymph ♂, 1 nymph ♀, coll. FC. — Valdieri, Gias della Casa, wet area, c. 1660 m, 44.1777°N, 7.2673°E (WGS84), 28.VII.2015, FC, SR, FT, PD, MD, MN, LB leg., 2 ♀, 1 ♂, 1 nymph ♀, coll. FC. — Valdieri, Gias della Casa, pebble ponds e Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, FC, SR, FT, MD, PD, MN, LB leg., 2 ♂, 3 ♀, coll. FC. — Valdieri, Pian del Valasco, 1747 m, 44.19608°N, 7.23051°E (WGS84), 20.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri (CN), Pian del Valasco, 1747 m, 5.VIII.2014, 2 specimens, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 2 specimens, MZUF.

CHOROTYPE. — Turano-European.

Family VELIIDAE Amyot & Serville, 1843

Velia (Plesiovelia) currens (Fabricius, 1794)

Gerris currens Fabricius, 1794: 193.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1400 m, 1964, MSNV; Limone Piemonte, L. Tamanini leg., coll. LT; Limone Piemonte, A. Servadei leg., 1952; coll. AS.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, FC, FT, MD, PD, SR, MN, LB leg., 2 juv., coll. FC. — Entracque, Trinità, right tributary of Bousset stream, 11.VI.2011, FC leg., 2 ♀, coll. FC. — Entracque, Tetti Violin, Torrente del Vallone Balme di Gherra, 1040 m, 10.VI.2011, FC leg., 7 ♀, 1 ♂, coll. FC. — Entracque, Trinità, torrente del Vallone del Prêr, c. 1090 m, 44.2168°N, 7.4311°E (WGS84), 28.VII.2015, FC, FT, SR, MD, PD, MN, LB leg., 2 juv., coll. FC. — Limone Piemonte, 25.VIII.1989, F.T. leg., 1 ♂, MZUF. — Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, FC, FT, SR, MD, PD, MN, LB leg., 1 ♂, 1 ♀, 1 nymph ♂, coll. FC.

CHOROTYPE. — Alpino-Appenninico Endemic.

Infraorder NEPOMORPHA Popov, 1968
Superfamily CORIXOIDEA Leach, 1815
Family CORIXIDAE Leach, 1815

Arctocorisa carinata (C. R. Sahlberg, 1819)
(Fig. 3)

Corixa carinata C. R. Sahlberg, 1819: 12.

NEW DATA. — Col di Boaria, La Perla small lake, 1500 m, 07.VII.2001, A. Carapezza leg., 6 ♂, 8 ♀, coll. AC. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, FC. SR, FT, PD, LB, MN, MD leg., 15 ♂, 3 ♀, 10 juv., coll. FC. — Limone Piemonte, SE del Bric Campanino, wet area (little lakes and ponds), 2125 m, 44.153°N, 7.604°E (WGS84), 27.VII.2015, FC. SR, FT, PD, LB, MN, MD leg., 10 ♂, 8 ♀, 24 juv., coll. FC.

CHOROTYPE. — Sibero-European.

Sigara (Vermicorixa) lateralis (Leach, 1817)

Corixa lateralis Leach, 1817: 17.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1400 m, 1964, MSNV.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, FC, FT, PD, SR, MB, LB, MD leg., 1 ♀, coll. FC.

CHOROTYPE. — Palaearctic.

Superfamily NOTONECTOIDEA Latreille, 1802
Family NOTONECTIDAE Latreille, 1802

Notonecta (Notonecta) viridis Delcourt, 1909

Notonecta viridis Delcourt, 1909: 379.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1400 m, 1964, MSNV.

NEW DATA. — Alpi Marittime, V. Ghiliani leg., [sub "*glauca*"], 1 ♂, 1 ♀, MSNG. — *idem*, [sub "*viridis*"], 2 ♂, MSNG. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, FC, SR, FT, MD, MN, LB leg., 1 ♂, 1 ♀, coll. FC.

CHOROTYPE. — Turano-European.




FIG. 3. — Male (A) and female (B) of *Arctocoris carinata* (C. R. Sahlberg, 1819). Scale bar: 1 mm. Photo: G. Strauss.

Infraorder LEPTOPODOMORPHA Popov, 1971
Superfamily SALDOIDEA Amyot & Serville, 1843
Family SALTIDAE Amyot & Serville, 1843

Chartoscirta cocksii (Curtis, 1835)

Acanthia cocksii Curtis, 1835: text to plate 548.

LITERATURE. — Carapezza & Faraci (2005, 2007): Colle Tenda, 1300 m, VI.1881, D. Baudi leg., 1 specimen, MSNG.

CHOROTYPE. — West-Palaeartic.

Macrosaldula scotica (Curtis, 1835)

Acanthia scotica Curtis, 1835: text to plate 548.

LITERATURE. — Faraci & Rizzotti Vlach (1992), Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1400 m, 12.VIII.1964, G. Osella leg., 6 specimens, MSNV; Colle di Tenda, 1320 m,

VII.1892, A. Dodero leg., 1 specimen, MSNG; Limone Piemonte, 1010 m, VIII.1892, A. Dodero leg., 1 specimen, MSNG; Mancini (1952 [sub *Saldula scotica*]): Limone Piemonte; Mancini (1963 [sub *Saldula scotica*]): Colle di Tenda; Péricart (1990): “Alpes Maritimes”.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, FC, FT, MD, PD, SR, MN, LB leg., 1 ♀, coll. FC. — Sant’Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, FC, FT, SR, MB, LB leg., 2 ♂, coll. FC. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, FC, SR, FT, MD, PD, MN, LB leg., 1 ♀, coll. FC.

CHOROTYPE. — West-Palaeartic.

Macrosaldula variabilis (Herrich-Schäffer, 1835)

Salda variabilis Herrich-Schäffer, 1835: 61.

NEW DATA. — Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, FC, FT, SR, MB, LB leg., 1 ♂, coll. FC.

CHOROTYPE. — Turano-European-Mediterranean.

Salda henschii (Reuter, 1891)
(Fig. 4)

Acanthia henschii Reuter, 1891: 23.

LITERATURE. — Cianferoni *et al.* (2017): Valdieri, Gias della Casa, small pools in a peat bog near Torrente Gesso della Valletta (stream), c. 1675 m a.s.l., 44.174°N, 7.267°E (WGS84), 28.VII.2015, FC leg., 5 ♂, 7 ♀, coll. FC; *idem*, SR leg., 1 ♀, coll. FC; Valdieri, Pian del Valasco, peat bog, c. 1775 m, 44.196°N, 7.230°E (WGS84), 27.VII.2012, MN leg., 1 ♂, MZUF.

CHOROTYPE. — Central-European.

REMARK. — This species was recently recorded for the first time in Italy (Cianferoni *et al.* 2017).

Saldula arenicola (Scholtz, 1847)

Salda arenicola Scholtz, 1847: 110.

NEW DATA. — Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, FC, FT, SR, MN, LB leg., 1 ♂, 2 ♀, coll. FC.

CHOROTYPE. — Palaearctic.

Saldula c-album (Fieber, 1859)

Salda c-album Fieber, 1859: 236.

LITERATURE. — Faraci & Rizzotti Vlach (1992), Carapezza & Faraci (2005, 2007): Colle di Tenda, 1300 m, VI.1881, D. Baudi leg., 1 specimen, MSNG.

NEW DATA. — Entracque, Tetti Porcera, right tributary of Bousset stream, Porcera bridge, 1080 m, 11.VI.2011, FC leg., 4 ♂, 3 ♀, coll. FC. — Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, FC, FT, SR, MN, LB leg., 2 ♀, coll. FC.

CHOROTYPE. — European.

Saldula melanoscela (Fieber, 1859)

Salda melanoscela Fieber, 1859: 236.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, FC, FT, MD, PD, SR, MN, LB leg., 1 ♀, coll. FC.

CHOROTYPE. — Asiatic-European.

Saldula orthochila (Fieber, 1859)

Salda orthochila Fieber, 1859: 234.

LITERATURE. — Faraci & Rizzotti Vlach (1992), Carapezza & Faraci (2005, 2007): Pian del Valasco (Valdieri), 1350 m, VII.1966, 1 specimen, MSNV; Valle Gesso, Soria Retreat, 1800 m, VII.1974, 1800 m, G. Osella leg., 2 specimens, MSNV.

NEW DATA. — Limone Piemonte, SE del Bric Campanino, wet area (little lakes and ponds), 2125 m, 44.153°N, 7.604°E (WGS84), 27.VII.2015, FC, FT, MD, PD, SR, MN, LB leg., 1 ♀, coll. FC.

CHOROTYPE. — Asiatic-European.

Saldula saltatoria (Linnaeus, 1758)

Cimex saltatorius Linnaeus, 1758: 448.

LITERATURE. — Faraci & Rizzotti Vlach (1992), Carapezza & Faraci (2005, 2007): Valle Gesso, Soria Retreat, 12.VII.1964, 1950 m, G. Osella leg., 1 specimen, MSNV; San Giacomo di Entracque, 17.VII.1964, 1400 m, M. Olmi leg., 1 specimen, MSNV.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, FC, FT, PD, SR, MN, LB, MD leg., 2 ♂, 2 ♀, 2 juv., coll. FC. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, FC, SR, FT, MD, PD, MN, LB leg., 1 ♀, coll. FC.

CHOROTYPE. — Holarctic.

Infraorder CIMICOMORPHA
Leston, Pendergrast & Southwood, 1954
Superfamily CIMICOIDEA Latreille, 1802
Family ANTHOCORIDAE Fieber, 1837

Anthocoris nemoralis (Fabricius, 1794)

Acanthia nemoralis Fabricius, 1794: 76.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF; Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European-Mediterranean.

Anthocoris nemorum (Linnaeus, 1761)

Cimex nemorum Linnaeus, 1761: 254.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 12 specimens, MZUF, MNHN; Limone Piemonte, 1250 m, 15.VII.1978, R. Poggi leg., [Alpi Liguri (CN), Dint. Piemonte, 15.VII.1978, R. Poggi], 1 specimen, MSNG. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MNHN. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, 1 juv., MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 speci-



FIG. 4. — *Salda henshii* (Reuter, 1891). Scale bar: 1 mm. Photo: F. Cianferoni.

men, MZUF. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410° E (WGS84), 09.IX.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Malaise trap, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 1 specimen, 1 juv., MZUF.

CHOROTYPE. — Asiatic-European.

Anthocoris simulans Reuter, 1884

Anthocoris minki var. *simulans* Reuter, 1884: 194.

LITERATURE. — Péricart (1972): Limone Piemonte.

CHOROTYPE. — Sibero-European.

Dufouriellus ater (Dufour, 1833)

Xylocoris ater Dufour, 1833: 104.

NEW DATA. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Orius vicinus (Ribaut, 1923)

Triphleps vicinus Ribaut, 1923: 529.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Orius minutus (Linnaeus, 1758)

Cimex minutus Linnaeus, 1758: 446.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF, MNHN. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Orius niger (Wolff, 1811)

Salda nigra Wolff, 1811: 167.

NEW DATA. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 3 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Temnostethus pusillus (Herrich-Schäffer, 1835)

Anthocoris pusillus Herrich-Schäffer, 1835: 60.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Superfamily MIROIDEA Hahn, 1831
Family MIRIDAE Hahn, 1831

Adelphocoris lineolatus (Goeze, 1778)

Cimex lineolatus Goeze, 1778: 267.

NEW DATA. — Valdieri, Valasco, 1450 m, 44.204067°N, 7.256410°E (WGS84), 31.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1450 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus

Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 28.VII.2015, LB & MN leg., 7 specimens, MZUF, MNHN. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, MN & LB leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Adelphocoris seticornis (Fabricius 1775)

Cimex seticornis Fabricius, 1775: 725.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Adelphocoris vandalicus (Rossi, 1790)

Cimex vandalicus Rossi, 1790: 249.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 3 specimens, MZUF.

CHOROTYPE. — Mediterranean.

Agnocoris rubicundus (Fallén, 1807)

Lygaeus rubicundus Fallén, 1807: 84.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Apolygus lucorum (Meyer-Dür, 1843)

Capsus lucorum Meyer-Dür, 1843: 46.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 2 specimens, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Asiatic-European.

Alloeotomus gothicus (Fallén, 1807)

Lygaeus gothicus Fallén, 1807: 93.

LITERATURE. — Mancini (1952, 1963), Carapezza & Faraci (2005, 2007): Colle di Tenda.

CHOROTYPE. — European.

Bryocoris pteridis (Fallén, 1807)

Capsus pteridis Fallén, 1807: 105.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Calocoris roseomaculatus (De Geer, 1773)

Cimex roseomaculatus De Geer, 1773: 293.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, MZUF; Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 7 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1636 m, 44.27444°N, 7.30426°E (WGS84), 22.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 05.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré 1679 m, 44.2041°N, 7.48709°E (WGS84), 27.VII.2013, MN, MD leg., 5 specimens, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 15 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, 1800 m, 08.VIII.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — European.

Calocoris affinis (Herrich-Schäffer, 1835)

Capsus affinis Herrich-Schäffer, 1835: 49, 82.

NEW DATA. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1300 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN, leg., 1 specimen, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.59948°E (WGS84), 11.VIII.2014, MN, MD leg., 4 specimens, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.59948°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 24.VII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Europeano-Mediterranean.

Calocoris alpestris (Meyer-Dur, 1843)

Capsus pabulinus var. *alpestris* Meyer-Dur, 1843: 50.

NEW DATA. — Valdieri, Pian del Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 27.VII.2012, MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 3 specimens, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 4 specimens, MZUF. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

REMARK. — Disjunct distribution of Boreo-Alpine type, but limited only to Europe.

Campyloneura virgula (Herrich-Schäffer, 1835)

Capsus virgula Herrich-Schäffer, 1835: 48.

NEW DATA. — Limone Piemonte, SE del Bric Campanino, wet area (little lakes and ponds), 2125 m, 44.153°N, 7.604°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — West-Palaearctic.

Capsodes flavomarginatus (Donovan, 1798)

Cimex flavomarginatus Donovan, 1798: 79.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — West-European.

Capsus ater (Linnaeus, 1758)

Cimex ater Linnaeus, 1758: 447.

NEW DATA. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — European.

Charagochilus gyllenhalii (Fallén, 1807)

Lygaeus gyllenhalii Fallén, 1807: 88.

NEW DATA. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Charagochilus weberi Wagner, 1953

Charagochilus weberi Wagner, 1953: 84.

LITERATURE. — Servadei (1957): Limone Piemonte.

CHOROTYPE. — Palaearctic.

REMARK

This is a doubtful taxon, that could be attributed to *Charagochilus spiralifer* Kerzhner, 1988 (see Simon 1992, Kment & Banar 2012).

Chlamydatus pullus (Reuter, 1870)

Agalliaestes pullus Reuter, 1870: 324 ["224"].

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Closterotomus annulus (Brullé, 1832)

Miris annulus Brullé, 1832: 78.

NEW DATA. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.61385°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 01.IX.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — East-European.

Closterotomus biclavatus (Herrich-Schäffer, 1835)

Capsus biclavatus Herrich-Schäffer, 1835: 48, 82.

NEW DATA. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, coll. PD.

CHOROTYPE. — West-Palaearctic.

Closterotomus norwegicus (Gmelin, 1790)

Cimex norwegicus Gmelin, 1790: 2176.

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — West-Palaearctic.

REMARK. — It is a very common West-Palaearctic species, introduced in North America and Australia (Carapezza 2007).

Closterotomus fulvomaculatus (De Geer, 1773)

Cimex fulvomaculatus De Geer, 1773: 294.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Compsidolon absinthii (Scott, 1870)

Agalliaestes absinthii Scott, 1870: 100.

NEW DATA. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Turano-European.

Cremnocephalus albolineatus (Reuter, 1875)

Cremnocephalus albolineatus Reuter, 1875c: 87.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Criocoris crassicornis (Hahn, 1834)

Phytocoris crassicornis Hahn, 1834: 90.

NEW DATA. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 ♂, 1 ♀, MZUF. — Valdieri, Pian del Valasco, 1766 m, 44.19608°N, 7.23051°E (WGS84), 27.VII.2012, MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 ♀, MZUF.

CHOROTYPE. — Asiatic-European.

Deraeocoris (Knightocapsus) lutescens (Schilling, 1837)

Phytocoris lutescens Schilling, 1837: 84.

LITERATURE. — Carapezza & Faraci (2005, 2007): Limone Piemonte, 1892, MSNG; Limone Piemonte, 1952, coll. AS.

CHOROTYPE. — West-Palaearctic.

Deraeocoris (Deraeocoris) ribauti Wagner, 1943

Deraeocoris luctuosus f. *ribauti* Wagner, 1943: 7.

LITERATURE. — Carapezza & Faraci (2005, 2007): Valdieri, 1963, MSNV; Tamanini (1977), Carapezza & Faraci (2005, 2007): Valdieri, VII-VIII.1977, G. Osella leg., MSNV.

CHOROTYPE. — West-European.

Deraeocoris (Deraeocoris) ruber (Linnaeus, 1758)

Cimex ruber Linnaeus, 1758: 446.

LITERATURE. — Mancini (1963): Colle di Tenda; Carapezza & Faraci (2005, 2007): Entracque, 1978, coll. AS; Entracque, San Giacomo, 1700 m, 1966, MSNV; Limone Piemonte, 1952, coll. AS; Valdieri, Sant'Anna, 1974, coll. AS.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 3 specimens, MZUF. — Valdieri, Madonna del Colletto, 1250 m, 44.28286°N, 7.38510°E (WGS84), 02.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — West-Palaearctic.

Deraeocoris (Deraeocoris) scutellaris (Fabricius, 1794)

Lygaeus scutellaris Fabricius, 1794: 180.

LITERATURE. — Mancini (1952, 1963): Colle di Tenda.

CHOROTYPE. — Asiatic-European.

Deraeocoris (Camptobrochis) serenus
Douglas & Scott, 1868

Camptobrochis serenus Douglas & Scott, 1868: 135.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1600 m, 1966, MSNV; Limone Piemonte, 1952, coll. AS.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-Mediterranean.

Dicyphus epilobii Reuter, 1883

Dicyphus epilobii Reuter, 1883a: 50, 52.

LITERATURE. — Tamanini (1956): Limone Piemonte, VIII, A. Servadei leg.

CHOROTYPE. — European.

Dicyphus (Brachyceraea) annulatus ononidis
(Wagner, 1951)

Brachyceraea ononidis Wagner, 1951: 24, 35.

LITERATURE. — Tamanini (1956): Limone Piemonte, VIII, A. Servadei leg.

CHOROTYPE. — Europeano-Mediterranean.

Dicyphus (Brachyceraea) globulifer (Fallén, 1829)

Capsus globulifer Fallén, 1829: 124.




FIG. 5. — Drawing of *Dimorphocoris ruffoi* Tamanini, 1971, from Tamanini (1976).

LITERATURE. — Tamanini (1956): Limone Piemonte, VIII, A. Servadei leg.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Dimorphocoris ruffoi Tamanini, 1971
(Fig. 5)

Dimorphocoris ruffoi Tamanini, 1971: 363.

LITERATURE. — Tamanini (1976), Carapezza & Faraci (2005, 2007): Entracque, Soria Retreat, loc. Prajet, northern slope of Entracque, 1840 m, 28.VII.1974, G. Osella leg., MSNV; Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1800 m, 1970, MSNV; Entracque, 1800 m, 1970, coll. AC.

CHOROTYPE. — Southwest Alpine endemic.

Europiella alpina (Reuter, 1875)

Plagiognatus alpinus Reuter, 1875b: 56.

NEW DATA. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-Europeano-Mediterranean.

Europiella artemisiae (Becker, 1864)

Capsus artemisiae Becker, 1864: 487.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Globiceps (Kelidocoris) flavomaculatus
(Fabricius, 1794)

Lygaeus flavomaculatus Fabricius, 1794: 182.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 28.VII.2015, LB & MN leg., 7 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, 2050 m, 44.20033°N, 7.47289°E (WGS84), 24.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 01.IX.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 01.IX.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Globiceps (Kelidocoris) fulvicollis Jakovlev, 1877

Globiceps fulvicollis Jakovlev, 1877: 293.

LITERATURE. — Mancini (1963): Colle di Tenda.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Globiceps (Globiceps) sphaegiformis (Rossi, 1790)

Cimex sphaegiformis Rossi, 1790: 250.

NEW DATA. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-European.

Hadrodemus m-flavum (Goeze, 1778)

Cimex m-flavum Goeze, 1778: 279.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 6 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 3 specimens, MZUF, MNHN. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 24.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 05.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1679 m, 44.2041°N, 7.48709°E (WGS84), 27.VII.2013,

MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1625 m, 44.19503°N, 7.49107°E (WGS84), 17.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 4 specimens, MZUF.

CHOROTYPE. — European.

Halticus apterus (Linnaeus, 1758)

Cicada aptera Linnaeus, 1758: 438.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1800 m, 1964, MSNV.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Halticus luteicollis (Panzer, 1804)

Lygaeus luteicollis Panzer, 1804: 18.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — West-Palaearctic.

Heterocapillus tigris (Mulsant & Rey, 1852)

Capsus tigris Mulsant & Rey, 1852: 129.

NEW DATA. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Hoplomachus thunbergii (Fallén, 1807)

Lygaeus thunbergii Fallén, 1807: 91.

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Horistus orientalis (Gmelin, 1790)

Cimex orientalis Gmelin, 1790: 2191.

NEW DATA. — Colle di Tenda, 1900 m, 07.VI.2001, A. Carapezza leg., 4 ♂, coll. AC. — Limone Piemonte, northern slope over the cableway, c. 1800 m, 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Europeano-Mediterranean.

Horwathia lineolata (A. Costa, 1862)

Capsus lineolatus A. Costa, 1862: 21.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.24899°N, 7.38166°E (WGS84), 22.VI.2014, MN, MD leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, MZUF, MNHN. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 20.VI.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 07-16.VIII.2013, Malaise trap, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 24.VII.2013, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Alpine endemic.

Lepidargyrus ancorifer (Fieber, 1858)

Apocremnus ancorifer Fieber, 1858: 336.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1978, coll. AS; Valdieri, Sant'Anna, 1978, coll. AS.

CHOROTYPE. — South-European.

Leptopterna dolabrata (Linnaeus, 1758)

Cimex dolabratus Linnaeus, 1758: 449.

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 5 specimens, MZUF.

CHOROTYPE. — Sibero-European.

Liocoris tripustulatus (Fabricius, 1781)

Cimex tripustulatus Fabricius, 1781: 370.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.24899°N, 7.38166°E (WGS84), 03.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Valasco, 1550 m, 31.VIII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Lopus decolor (Fallén, 1807)

Capsus decolor Fallén, 1807: 102.

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — European.

Lygocoris pabulinus (Linnaeus, 1761)

Cimex pabulinus Linnaeus, 1761: 253.

NEW DATA. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Holarctic.

Lygocoris rugicollis (Fallén, 1807)

Lygaeus rugicollis Fallén, 1807: 76.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, coll. PD, MZUF.

CHOROTYPE. — Holarctic.

Lygus pratensis (Linnaeus, 1758)

Cimex pratensis Linnaeus, 1758: 448.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 28.VII.2015, LB & MN leg., 7 specimens, MZUF, MNHN. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.19694°N, 7.4838°E (WGS84), 21.VI.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palearctic.

Lygus punctatus (Zetterstedt, 1838)

Phytocoris punctatus Zetterstedt, 1838: 273.

NEW DATA. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 15 specimens, coll. PD, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 28.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Gias della Casa, Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, FC, SR, FT, MD, PD, MN, LB leg., 4 specimens, MZUF. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 12 specimens, MZUF, MNHN.

CHOROTYPE. — Holarctic.

Lygus wagneri (Remane, 1955)

Lygus (Exolygus) wagneri Remane, 1955: 115.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 12 specimens, MZUF. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2003, MN, MD leg., 10 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, pebble ponds e Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 4 specimens, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 30.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Terme di Valdieri, La Valletta, 1552 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.1914, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré 1625 m, 44.19694°N, 7.4838°E (WGS84), 17.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 5 specimens, MZUF. — Vernante, loc. Palanfré, 1770 m, 44.198119°N, 7.478328°E (WGS84), 21.VIII.2014, MN, MD leg., 4 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1920 m, 44.199977°N, 7.474896°E (WGS84), 26.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2000 m, 44.20033°N, 7.47289°E (WGS84), 24.VIII.2014, MN, MD leg., 4 specimens, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 5 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 10 specimens, MZUF.

CHOROTYPE. — Asiatic-European.

Macrotylus (Macrotylus) herrichi (Reuter, 1873)

Hoplomachus herrichi Reuter, 1873: 24.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 2 specimens, MZUF. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Turano-European.

Macrotylus (Macrotylus) quadrilineatus (Schränk, 1785)

Cimex quadrilineatus Schrank, 1785: 339.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri,

Pian del Valasco, 1747 m, 44.19608°N, 7.23051°E (WGS84), 20.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 24.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, Norbiato, 1 specimen, MZUF.

CHOROTYPE. — European.

Mecomma (Globicellus) dispar (Boheman, 1852)

Cyllecoris dispar Boheman, 1852: 72.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 ♀, MZUF.

CHOROTYPE. — Sibero-European.

REMARK

In Northern Europe this species lives in coastal wetlands, while S of the Alps it can be found between 1000 and 1900 m a.s.l.

Megaloceroea recticornis (Geoffroy, 1785)

Cimex recticornis Geoffroy, 1785: 209.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 3 specimens, MZUF. — Entracque, Fus Carbonaro, 950 m, 44.250810°N, 7.377315°E (WGS84), 06.VII.2014, MN, MD leg., 2 specimens, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, small pools in a peat bog and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 05.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1625 m, 44.19694°N, 7.4838°E (WGS84), 17.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 9 specimens, MZUF. — Vernante, loc. Palanfré, 1730 m, 44.2041°N, 7.48709°E (WGS84), 06.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MNHN. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — West-Palaearctic.

REMARK

Introduced in North America and New Zealand; it is one of the more common Italian mirids (Carapezza 2007).

Miris striatus (Linnaeus, 1758)

Cimex striatus Linnaeus, 1758: 449.

NEW DATA. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 28.VI.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Notostira elongata (Geoffroy, 1785)

Cimex elongatus Geoffroy, 1785: 208.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 2 specimens, MZUF, coll. PD.

CHOROTYPE. — Palaearctic.

Notostira erratica (Linnaeus, 1758)

Cimex erraticus Linnaeus, 1758: 449.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 ♀, MZUF. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2003, MN, MD leg., 10 specimens, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 23.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 19.VI.2014, MN, MD leg., 2 ♀, MZUF.

CHOROTYPE. — Turano-European.

Orthocephalus coriaceus (Fabricius, 1777)

Acanthia coriacea Fabricius, 1777: 299.

NEW DATA. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 ♀, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 2 ♀, MZUF.

CHOROTYPE. — European.

Orthocephalus saltator (Hahn, 1835)

Capsus saltator Hahn, 1835: 11.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1964, MSNV.

CHOROTYPE. — Palaearctic.

Orthocephalus vittipennis (Herrich-Schäffer, 1835)

Orthocephalus vittipennis Herrich-Schäffer, 1835: 52, 88.

LITERATURE. — Carapezza & Faraci (2005, 2007) ["uncertain identification"]: Entracque, San Giacomo, 1600 m, 1966, MSNV.

CHOROTYPE. — Sibero-European.

Orthotylus marginalis Reuter, 1883

Orthotylus marginalis Reuter, 1883b: 350.

LITERATURE. — Mancini (1963): Colle di Tenda.

CHOROTYPE. — Sibero-European.

Orthonotus rufifrons (Fallén, 1807)

Capsus rufifrons Fallén, 1807: 105.

NEW DATA. — Entracque, S of Ponte Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 ♀, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 ♂, MZUF.

CHOROTYPE. — European.

Orthops campestris (Linnaeus, 1758)

Cimex campestris Linnaeus, 1758: 448.

NEW DATA. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — West-Palaearctic.

Orthops kalmii (Linnaeus, 1758)

Cimex kalmii Linnaeus, 1758: 448.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Orthops montanus (Schilling, 1837)

Phytocoris montanus Schilling, 1837: 83.

NEW DATA. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Europeano-Mediterranean.

Pachytomella passerinii (A. Costa, 1842)

Phytocoris passerinii A. Costa, 1842: 288.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, c. 2120 m, 44.19726°N, 7.47119°E (WGS84), 01.IX.2014, MN, MD leg., 1 ♀, MZUF.

CHOROTYPE. — Mediterranean.

REMARK

Despite the Mediterranean distribution range, this species has been found also at high altitudes (> 2000 m) and in cold habitats (e.g. Ehanno 1987).

Phytocoris austriacus Wagner, 1954

Phytocoris (*Ktenocoris*) *austriacus* Wagner, 1954: 9, 30.

LITERATURE. — Servadei (1957): Limone Piemonte.

CHOROTYPE. — South-European.

Phytocoris varipes Boheman, 1852

Phytocoris varipes Boheman, 1852: 107.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-European.

Pilophorus confusus (Kirschbaum, 1856)

Capsus confusus Kirschbaum, 1856: 232.

LITERATURE. — Mancini (1963): Colle di Tenda.

CHOROTYPE. — Palaearctic.

Pinalitus rubricatus (Fallén, 1807)

Lygaeus rubricatus Fallén, 1807: 91.

NEW DATA. — Valdieri, Valasco, 1550 m, 44.204067°N, 7.256410°E (WGS84), 31.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian del Valasco, 1747 m, 44.19608°N, 7.23051°E (WGS84), 20.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Pithanus maerkelii (Herrich-Schäffer, 1838)
(Fig. 6)

Capsus maerkelii Herrich-Schäffer, 1838: 78.

NEW DATA. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Placochilus seladonicus (Fallén, 1807)

Cimex seladonicus Fallén, 1807: 91.

NEW DATA. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 9 specimens, MZUF.

CHOROTYPE. — European.

Plagiognathus arbustorum (Fabricius, 1794)

Lygaeus arbustorum Fabricius, 1794: 175.

NEW DATA. — Colle di Tenda, 1.900 m, 07.VII.2001, A. Carapezza leg., 1 ♂, coll. AC. — Entracque, Fus Carbonaro, 1000 m, 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 950 m, 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 7 specimens, MZUF, MNHN. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Plagiognathus chrysanthemi (Wolff, 1804)

Miris chrysanthemi Wolff, 1804: 157.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1650 m, 44.17084°N, 7.61385°E (WGS84), 26.IX.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 11 specimenx, MZUF.

CHOROTYPE. — Asiatic-European.

Parapsallus vitellinus (Scholtz, 1847)

Capsus vitellinus Scholtz, 1847: 130.




FIG. 6. — Male (A) and female (B) of *Pithanus maerkelii* (Herrich-Schäffer, 1838). Scale bars: 1 mm. Photo: G. Strauss.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 20.VI.2014, MN, MD leg., 2 specimens, MZUF, coll. PD.

CHOROTYPE. — Sibero-European.

Polymerus (Poeciloscytus) cognatus (Fieber, 1858)

Poeciloscytus cognatus Fieber, 1858: 331.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Polymerus (Poeciloscytus) unifasciatus
(Fabricius, 1794)

Lygaeus unifasciatus Fabricius, 1794: 178.

NEW DATA. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1679 m, 44.2041°N, 7.48709°E (WGS84), 27.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Psallus (Mesopsallus) ambiguus (Fallén, 1807)

Lygaeus ambiguus Fallén, 1807: 89.

LITERATURE. — Mancini (1963 [sub *Psallus (Apocremonus) ambiguus*]): Colle di Tenda.

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Psallus (Psallus) varians (Herrich-Schäffer, 1841)

Capsus varians Herrich-Schäffer, 1841: 45.

NEW DATA. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Stenodema (Stenodema) sericans (Fieber, 1861)

Miris sericans Fieber, 1861: 240.

NEW DATA. — Valdieri, Pian del Valasco, 1747 m, 44.19608°N, 7.23051°E (WGS84), 20.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1679 m, 44.2041°N, 7.48709°E (WGS84), 27.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1625 m, 44.19503°N, 7.49107°E (WGS84), 17.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1730 m, 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1770 m, 21.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — South-European.

Stenodema (Stenodema) algoviense (Schmidt, 1934)

Stenodema algoviense Schmidt, 1934: 34.

NEW DATA. — Entracque, Vej del Bouc lake, 2054 m, 44.14209°N, 7.42284°E (WGS84), 23.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Alpine endemic.

Stenodema (Stenodema) holsata (Fabricius, 1787)

Cimex holsatus Fabricius, 1787: 306.

NEW DATA. — Entracque, Vej del Bouc lake, 2054 m, 44.14209°N, 7.42284°E (WGS84), 23.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 4 specimens, MZUF. — Limone Piemonte, Gias della Perla. — La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 5 specimens, MZUF, MNHN. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 11 specimenx, MZUF. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 4 specimens, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 2 specimens, MZUF. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2003, MN, MD leg., 10 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1636 m, 44.27444°N, 7.30426°E (WGS84), 22.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione,

1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Gias della Casa, small pools in a peat bog e Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 3 specimens, MZUF. — Valdieri, Pian del Valasco, 1747 m, 44.19608°N, 7.23051°E (WGS84), 20.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 30.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1740 m, 07.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Valdieri, Valasco, 1550 m, 44.204067°N, 7.256410°E (WGS84), 31.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1679 m, 44.2041°N, 7.48709°E (WGS84), 27.VII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 17.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 23.VI.2013, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1625 m, 44.19503°N, 7.49107°E (WGS84), 17.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Stenodema (Stenodema) laevigata (Linnaeus, 1758)

Cimex laevigatus Linnaeus, 1758: 449.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 3 specimens, MZUF. — Valdieri, Colle dell'Arpione 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, 1 juv., MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Stenodema (Stenodema) virens (Linnaeus, 1767)

Cimex virens Linnaeus, 1767: 730.

LITERATURE. — Servadei (1955 [sub *Stenodema virens* v. *fulva* (Fieber, 1837)]): Limone Piemonte.

CHOROTYPE. — Asiatic-European.

Stenotus binotatus (Fabricius, 1794)

Lygaeus binotatus Fabricius, 1794: 172.

NEW DATA. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 2 specimens, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 3 specimens, MZUF, MNHN. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 05.VIII.2013, MN, MD leg., 6 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1625 m, 44.19503°N, 7.49107°E (WGS84), 17.VIII.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Strongylocoris leucocephalus (Linnaeus, 1758)

Cimex leucocephalus Linnaeus, 1758: 446.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1250 m, 1964, MSNV.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 4 specimens, MZUF. — Valdieri, Colle dell'Arpione 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1636 m, 44.27444°N, 7.30426°E (WGS84), 26.VI.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1750 m, 44.19694°N, 7.4838°E (WGS84), 30.VII.2014, MN, MD leg., 1 specimen, MNHN. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Trigonotylus ruficornis (Geoffroy, 1785)

Cimex ruficornis Geoffroy, 1785: 209.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1250 m, 1964, MSNV.

NEW DATA. — Valdieri, Pian del Valasco, 1750 m, 44.19608°N, 7.23051°E (WGS84), 20.VIII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Family TINGIDAE (Laporte, 1832)

Acalypta finitima (Puton, 1884)

Orthostira finitima Puton, 1884: 313.

LITERATURE. — Péricart (1983), Carapezza & Faraci (2005, 2007): Entracque, Lago Brocan, 2020 m, 1930.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015 (litter sieving), LB & MN leg., 1 specimen, MZUF.

CHOROTYPE. — Central-European.

Acalypta samara (Puton, 1887)

Orthostira samara Puton, 1887: 98.

LITERATURE. — Roubal (1958 [sub *Acalypta* (*Orthostira*) *samara*]): Entracque, Lago Brocan.

CHOROTYPE. — European.

REMARK

This species has an Alpine-Balkan disjoint distribution.

Acalypta visolensis visolensis Péricart, 1976

Acalypta visolensis Péricart, 1976: 174.

LITERATURE. — Péricart (1983), Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1800 m, 1970.

CHOROTYPE. — Alpine endemic.

Catoplatus fabricii (Stål, 1868)

Tingis fabricii Stål, 1868: 93.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 3 specimens, MZUF.

CHOROTYPE. — European.

Copium clavicorne clavicorne (Linnaeus, 1758) (Fig. 7)

Cimex clavicornis Linnaeus, 1758: 442.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1250 m, 1964, MSNV.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 5 specimens,




FIG. 7. — Female (A) and male (B) of *Copium clavicorne clavicorne* (Linnaeus, 1758). Scale bars: 1 mm. Photo: G. Strauss.

MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Malaise trap, 1680 m, 44.2041°N, 7.48709°E (WGS84), 27.VI.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-European.

REMARK

This species has Turano-European range with extension to the Arabian Peninsula.

Dictyla echii (Schrank, 1782)

Cimex echii Schrank, 1782: 276.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1250 m, 1966, MSNV.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Tingis (Tingis) cardui (Linnaeus, 1758)

Cimex cardui Linnaeus, 1758: 443.

NEW DATA. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2013, MN, MD leg., 5 specimens, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 23.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Tingis (Tropidocheila) maculata (Herrich-Schäffer, 1838)

Monanthia maculata Herrich-Schäffer, 1838: 51, 56.

LITERATURE. — Carapezza & Faraci (2005, 2007): Entracque, San Giacomo, 1250 m, 1964, MSNV.

CHOROTYPE. — Turano-European.

Superfamily NABOIDEA A. Costa, 1853
Family NABIDAE A. Costa, 1853

Himacerus (Aptus) mirmicoides (O. Costa, 1834)

Nabis mirmicoides O. Costa, 1834: 71.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 3 specimens, MZUF.

CHOROTYPE. — Europeano-Mediterranean.

Himacerus (Himacerus) apterus (Fabricius, 1798)

Reduvius apterus Fabricius, 1798: 546.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 2 specimens, MZUF.

CHOROTYPE. — Asiatic-European.

Nabis (Nabicula) flavomarginatus
Scholtz, 1847

Nabis flavomarginatus Scholtz, 1847:114.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MNHN. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 5 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Nabis (Nabis) punctatus punctatus
A. Costa, 1847

Nabis punctatus A. Costa, 1847: 14.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 2 ♀, MZUF.

CHOROTYPE. — West-Palaearctic.

Nabis (Nabis) pseudoferus pseudoferus
Remane, 1949

Nabis pseudoferus pseudoferus Remane, 1949: 63.

LITERATURE. — Servadei (1955 [sub *Nabis pseudoferus*]): Limone Piemonte.

CHOROTYPE. — European.

Nabis (Nabis) rugosus (Linnaeus, 1758)

Cimex rugosus Linnaeus, 1758: 442.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 2 specimens, MZUF. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 28.VII.2015, LB & MN leg., 7 specimens, MZUF, MNHN. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 17.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 23.VI.2013, 1 specimen, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1770 m, 44.198119°N, 7.478328°E (WGS84), 21.VIII.2014, MN, MD leg., 1 ♂, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 3 specimens, MZUF.

CHOROTYPE. — Sibero-European.

Prostemma aeneicolle Stein, 1857

Prostemma aeneicolle Stein, 1857: 88.

LITERATURE. — Péricart (1987): « Piémont : [...] Alpes Maritimes, jusqu'à 2100 m d'altitude. » [Piedmont: Maritime Alps, up to 2100 m a.s.l.].

CHOROTYPE. — European.

Superfamily REDUVIOIDEA Börner, 1910
Family REDUVIIDAE Latreille, 1807

Rhynocoris annulatus (Linnaeus, 1758)

Cimex annulatus Linnaeus, 1758: 447.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Rhynocoris iracundus (Poda, 1761)

Cimex iracundus Poda, 1761: 58.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 3 specimens, coll. CR.

NEW DATA. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1630 m, 44.19694°N, 7.4838°E (WGS84), 12.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84),




FIG. 8. — Male of *Aneurus (Aneurodes) avenius* (Dufour, 1833). Scale bar: 1 mm. Photo: G. Strauss.

11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 3 specimens, MZUF.

CHOROTYPE. — Asiatic-European.

Infraorder PENTATOMOMORPHA
Leston, Pendegrast & Southwood, 1954
Superfamily ARADOIDEA Spinola, 1837
Family ARADIDAE Spinola, 1837

Aneurus (Aneurodes) avenius (Dufour, 1833)
(Fig. 8)

Aradus avenius Dufour, 1833: 182.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 06.VI.2014, MN, MD leg., 1 specimen [identified by photo].

CHOROTYPE. — Palaearctic.

Superfamily COREOIDEA Leach, 1815
Family ALYDIDAE Amyot & Serville, 1843

Alydus calcaratus (Linnaeus, 1758)

Cimex calcaratus Linnaeus, 1758: 450.

NEW DATA. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 3 specimens, MZUF.

CHOROTYPE. — Holarctic.

Camptopus lateralis (Germar, 1817)

Alydus lateralis Germar, 1817: 285.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 3 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Family COREIDAE Leach, 1815

Nemocoris fallenii R. F. Sahlberg, 1848

Nemocoris fallenii R. F. Sahlberg, 1848: 42.

LITERATURE. — Mancini (1960): Colle di Tenda.

CHOROTYPE. — Sibero-European.

Ceraleptus gracilicornis (Herrich-Schäffer, 1835)

Coreus gracilicornis Herrich-Schäffer, 1835: plate 5 with legend.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-European-Mediterranean.

Coreus marginatus (Linnaeus, 1758)

Cimex marginatus Linnaeus, 1758: 443.

LITERATURE. — EDIT 2010: Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 3 specimens, coll. CR.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MNHN. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2013, MN, MD leg., 6 specimens, MZUF. — Valdieri, Pian del Valasco, 1747 m, 44.19608°N, 7.23051°E




FIG. 9. — Male (A) and female (B) of *Spathocera lobata* (Herrich-Schäffer, 1840). Scale bar: 1 mm. Photo: G. Strauss.

(WGS84), 20.VIII.2013, MN, MD leg., 1 specimen juv., MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 4 specimens, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 1 specimen juv., MZUF. — Valdieri, Valasco, 1550 m, 44.204067°N, 7.256410°E (WGS84), 31.VIII.2013, MN, MD leg., 1 specimen juv., MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Coriomeris denticulatus (Scopoli, 1763)

Cimex denticulatus Scopoli, 1763: 125.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E

(WGS84), 27.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Spathocera lobata (Herrich-Schäffer, 1840) (Fig. 9)

Pseudophloeus lobatus Herrich-Schäffer, 1840: 3, 6.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, coll. PD, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Family RHOPALIDAE Amyot & Serville, 1843

Corizus hyosciامي hyosciامي (Linnaeus, 1758)

Cimex hyosciامي Linnaeus, 1758: 447.

LITERATURE. — Mancini (1963 [sub *Corizus hyosciami*]): Colle di Tenda.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1740 m, 44.17446°N, 7.2672°E (WGS84), 07.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1550 m, 44.204067°N, 7.256410°E (WGS84), 31.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MNHN. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014 MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Myrmus miriformis miriformis (Fallén, 1807)

Coreus miriformis Fallén, 1807: 60.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF.

CHOROTYPE. — Sibero-European.

Rhopalus (Rhopalus) parumpunctatus Schilling, 1829

Rhopalus parumpunctatus Schilling, 1829: 53.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2013, MN, MD leg., 2 specimens, MZUF. — Valdieri, Pian del Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 27.VII.2012, MN leg., 2 specimens, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Rhopalus (Rhopalus) subrufus (Gmelin, 1790)

Cimex subrufus Gmelin, 1790: 2168.

LITERATURE. — EDIT 2010: Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 28.VII.2015, LB & MN leg., 5 specimens, MZUF, MNHN. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, c. 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 4 specimens, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Rhopalus (Rhopalus) conspersus (Fieber, 1837)

Coreus conspersus Fieber, 1837: 350.

LITERATURE. — Mancini (1963): Colle di Tenda.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 5 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Stictopleurus abutilon (Rossi, 1790)

Cimex abutilon Rossi, 1790: 242.

LITERATURE. — Tamanini (1951): Colle di Tenda, "collez. [collection] Paolo Luigioni".

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 5 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Stictopleurus crassicornis (Linnaeus, 1758)

Cimex crassicornis Linnaeus, 1758: 448.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Valasco, 1500 m, 44.204067°N, 7.256410°E (WGS84), 09.IX.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E

(WGS84), 13.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1920 m, 27.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Stictopleurus punctatonervosus (Goeze, 1778)

Cimex punctatonervosus Goeze, 1778: 265.

NEW DATA. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 4 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Asiatic-European.

Superfamily LYGAEOIDEA Schilling, 1829
Family BLISSIDAE Stål, 1862

Ischnodemus quadratus Fieber, 1837

Ischnodemus quadratus Fieber, 1837: 338.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Family CYMIDAE Baeresprung, 1860

Cymus glandicolor Hahn, 1832

Cymus glandicolor Hahn, 1832: 79.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Asiatic-European.

Family GEOCORIDAE Baeresprung, 1860

Geocoris (Geocoris) lapponicus Zetterstedt, 1838
(Fig. 10)

Geocoris lapponicus Zetterstedt, 1838: 266.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.



FIG. 10. — Female of *Geocoris (Geocoris) lapponicus* Zetterstedt, 1838. Scale bar: 1 mm. Photo: G. Strauss.

Family HETEROGASTRIDAE Stål, 1872

Platyplax salviae (Schilling, 1829)

Heterogaster salviae Schilling, 1829: 85.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Heterogaster cathariae (Geoffroy, 1785)

Cimex cathariae Geoffroy, 1785: 203.

LITERATURE. — Péricart (1999a): Limone Piemonte, A. Servadei leg.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Heterogaster urticae (Fabricius, 1775)

Cimex urticae Fabricius, 1775: 723.

NEW DATA. — Limone Piemonte, VII.1915, G. Mantero leg., 2 specimens, MSNG.

CHOROTYPE. — Palaearctic.

Family ISCHNORYNCHYDAE Stål, 1872

Kleidocerys resedae (Panzer, 1797)

Lygaeus resedae Panzer, 1797: 20.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 7 specimens, MZUF, MNHN. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 4 specimens, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 20.VI.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 6 specimens, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 28.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1636 m, 44.27444°N, 7.30426°E (WGS84), 26.VI.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 24-30.VII.2013, Malaise trap, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1920 m, 27.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 30.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 2000 m, 24.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 01.IX.2014, MN, MD leg., 11 specimenx, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 01.IX.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 4 specimens, MZUF. — Vernante, loc. Palanfré, Malaise trap, 1680 m, 44.2041°N, 7.48709°E (WGS84), 27.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Family LYGAEIDAE Schilling, 1829

Lygaeus equestris (Linnaeus, 1758)

Cimex equestris Linnaeus, 1758: 447.

LITERATURE. — Ferrari (1884): "Alpi marittime e Piemonte (Ghiliani, Garbiglietti)"; Noelli (1897b [sub *Eulygaeus equestris*]): Limone Piemonte, Griffini leg.; Colle di Tenda, Nobili leg.

NEW DATA. — Alpi Marittime, V. Ghiliani leg., [sub *Spilostethus equestris*], 1 specimen, MSNG. — Alpi Marittime, VI.76, R. Oberthur leg., [sub *Aphanus phoeniceus*], 2 specimens, MSNG. — Limone Piemonte, VII.1938, G. Mantero leg., [sub *Spilostethus equestris*], 1 specimen, MSNG. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 30.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Lygaeus simulans Deckert, 1985

Lygaeus simulans Deckert, 1985: 277.

LITERATURE. — Dioli *et al.* (1994): Limone Piemonte, 28.VIII.1952, 3 specimens, coll. AS; San Giacomo di Entracque, 12.VII.1964, 3 specimens, MSNV; San Giacomo di Entracque, 1400-2000 m, VII.1966, 5 specimens, MSNV; San Giacomo di Entracque, 1500 m, 03.IX.1971, 1 specimen, MSNV; Colle di Tenda, 27.VII.1963, 1 specimen, MSNM; Limone Piemonte, 28.VII.1930, 1 specimen, MZR; Limone Piemonte, 20.VII.1963, 1 specimen, MSNM; Valle Gesso, Soria Reatreat, 1800 m, VII.1974, 4 specimens, MSNV.

NEW DATA. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-European.

Melanocoryphus albomaculatus (Goeze, 1778)

Cimex albomaculatus Goeze, 1778: 264.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Tropidothorax leucopterus (Goeze, 1778) (Fig. 11)

Cimex leucopterus Goeze, 1778: 264.

NEW DATA. — Alpi Marittime, VI.74, R. Oberthur leg., [sub *Lygaeus leucopterus*], 1 specimen, MSNG. — Alpi Marittime, VI.76, R. Oberthur leg., 1 specimen, MSNG.

CHOROTYPE. — Centralasiatic-Europeano-Mediterranean.




FIG. 11. — Male (A) and female (B) of *Tropidothorax leucopterus* (Goeze, 1778). Scale bar: 1 mm. Photo: G. Strauss.

Spilostethus saxatilis (Scopoli, 1763)

Cimex saxatilis Scopoli, 1763: 128.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, VIII.92, Doderò leg., 1 specimen, MSNG. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 21.VI.2013, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 23.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 2 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 27.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 3 specimens, MZUF.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Spilostethus pandurus (Scopoli, 1763)

Cimex pandurus Scopoli, 1763: 126.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Subcosmopolitan.

Family ORSILLIDAE Stål, 1872

Nithecus jacobaeae (Schilling, 1829)

Heterogaster jacobaeae Schilling, 1829: 87.

NEW DATA. — Colle di Tenda, 1900 m, 07.VI.2001, A. Carapezza leg., 2 ♂, 3 ♀, coll. AC. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 6 specimens, MZUF, MNHN. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 2020 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Colle dell'Arpione,

1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 05.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 08.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 30.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Nysius cymoides (Spinola, 1837)

Artheneis cymoides Spinola, 1837: 252.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Nysius graminicola (Kolenati, 1845)

Corizus ericae var. *graminicola* Kolenati, 1845: 21.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Nysus thymi thymi (Wolff, 1804)

Lygaeus thymi Wolff, 1804: 149.

NEW DATA. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 8 specimens, MZUF, MNHN. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Ortholomus punctipennis (Herrich-Schäffer, 1838)

Heterogaster punctipennis Herrich-Schäffer, 1838: 75.

NEW DATA. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Family OXYCARENIDAE Stål, 1862

Macroplax preysleri (Fieber, 1837)

Oxycarenus preysleri Fieber, 1837: 343.

NEW DATA. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — European.

Family RHYPAROCHROMIDAE Amyot & Serville, 1843

Acompus pallipes (Herrich-Schäffer, 1834)

Pachymerus pallipes Herrich-Schäffer, 1834: 2.

LITERATURE. — Péricart (1998 [sub *Acompus pallipes*]): Limone Piemonte, MSNV.

CHOROTYPE. — Turano-European-Mediterranean.

Drymus (Sylvadrymus) sylvaticus (Fabricius, 1775)

Cimex sylvaticus Fabricius, 1775: 722.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Emblethis verbasci (Fabricius, 1803)

Lygaeus verbasci Fabricius, 1803: 235.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

CHOROTYPE. — Turano-European-Mediterranean.

Gastrodes grossipes grossipes (De Geer, 1773)

Cimex grossipes De Geer, 1773: 308.

LITERATURE. — Mancini (1963 [sub *Gastrodes grossipes*]): Colle di Tenda.

CHOROTYPE. — Asiatic-European.

Ligyrocoris sylvestris (Linnaeus, 1758)

Cimex sylvestris Linnaeus, 1758: 449.

NEW DATA. — Vernante, loc. Palanfré, 1625 m, 44.19503°N, 7.49107°E (WGS84), 17.VIII.2013, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Holarctic.

Peritrechus geniculatus (Hahn, 1832)

Pachymerus geniculatus Hahn, 1832: 68.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 4 specimens, MZUF. — Vernante, loc. Palanfré, 1670 m, 44.2041°N, 7.48709°E (WGS84), 11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 23.VI.2013, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré 1635 m, 44.187716°N, 7.492824°E (WGS84), 11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF, MNHN.

CHOROTYPE. — Palaearctic.

Peritrechus gracilicornis Puton, 1877

Peritrechus gracilicornis Puton, 1877: 117.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-European-Mediterranean.

Pterotmetus staphyliniformis (Schilling, 1829)

Pachymerus staphyliniformis Schilling, 1829: 77.

LITERATURE. — EDIT 2010: Entracque, Torrente Bousset, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

NEW DATA. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque (CN), Fus Carbonaro, 1000 m, 22.VI.2014 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Rhyparochromus phoeniceus (Rossi, 1794)

Cimex phoeniceus Rossi, 1794: 54.

LITERATURE. — EDIT 2010: Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR; Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 2 specimens, coll. CR.

NEW DATA. — Alpi Marittime, VI.76, R. Oberthur leg., [sub *Aphanus phoeniceus*], 1 specimen, MSNG. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Valdieri, Pian del Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 27.VII.2012, MN leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-European-Mediterranean.

Rhyparochromus pini (Linnaeus, 1758)

Cimex pini Linnaeus, 1758: 448.

LITERATURE. — EDIT 2010: Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 28.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1650 m, 44.19694°N, 7.4838°E (WGS84), 26.IX.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 4 specimens, MZUF. — Vernante, loc. Palanfré, Vilazzo lake, 1840 m, 44.16689°N, 7.49876°E (WGS84), 15.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Raglius alboacuminatus (Goeze, 1778)

Cimex alboacuminatus Goeze, 1778: 266.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR; Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 4 specimens, coll. CR.

CHOROTYPE. — Holarctic.

Raglius confusus (Reuter, 1886)

Pachymerus confusus Reuter, 1886: 121.

NEW DATA. — Alpi Marittime, V. Ghiliani leg., [sub *Aphanus confusus*], 1 specimen, MSNG.

CHOROTYPE. — Turano-European-Mediterranean.

Scolopostethus thomsoni Reuter, 1875

Scolopostethus thomsoni Reuter, 1875a: 562.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Holarctic.

Trapezonotus (Trapezonotus) arenarius
(Linnaeus, 1758)

Cimex arenarius Linnaeus, 1758: 448.

LITERATURE. — EDIT 2010: Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 2 specimens, coll. CR.

CHOROTYPE. — Palaearctic.

Trapezonotus (Trapezonotus) desertus
Seidenstucker, 1951

Trapezonotus desertus Seidenstucker, 1951: 79.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 3 specimens, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, 3 juv., MZUF.

CHOROTYPE. — Holarctic.

Superfamily PYRRHOCOREOIDEA Southwood, 1956
Family PYRRHOCORIDAE Fieber, 1860

Pyrrhocoris apterus (Linnaeus, 1758)

Cimex apterus Linnaeus, 1758: 557.

NEW DATA. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Superfamily PENTATOMOIDEA Reuter, 1910
Family CYDNIDAE Billberg, 1820

Canthophorus impressus (Horváth, 1881)

Sehirus impressus Horváth, 1881: 184.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Legnotus limbosus (Geoffroy, 1785)

Cimex limbosus Geoffroy, 1785: 217.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

CHOROTYPE. — Turano-European-Mediterranean.

Legnotus picipes (Fallén, 1807)

Cydnus picipes Fallén, 1807: 54.

NEW DATA. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Tritomegas rotundipennis (Dohrn, 1862)

Sehirus rotundipennis Dohrn, 1862: 211.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — South-European.

Family PENTATOMIDAE Leach, 1815

Aelia acuminata (Linnaeus, 1758)

Cimex acuminatus Linnaeus, 1758: 446.

NEW DATA. — Colle di Tenda, 1900 m, 07.VI.2001, A. Carapezza leg., 1 ♀, coll. AC. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MNHN. — Entracque, loc. Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 21.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 23.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1650 m, 44.19694°N, 7.4838°E (WGS84), 11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1730 m, 06.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1780 m, 44.198119°N, 7.478328°E (WGS84), 21.VIII.2014, MN, MD leg., 2 specimens juv., MZUF.

CHOROTYPE. — Palaearctic.

Aelia klugii Hahn, 1833
(Fig. 12)

Aelia klugii Hahn, 1833: 122.

NEW DATA. — Colle di Tenda, 1900 m, 07.VI.2001, A. Carapezza leg., 5 ♀, coll. AC. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1920 m, 44.199977°N, 7.474896°E (WGS84), 26.VI.2014, MN, MD, 3 specimens, MZUF, coll. PD.

CHOROTYPE. — Palaearctic.

Aelia rostrata rostrata Boheman, 1852

Aelia rostrata rostrata Boheman, 1852: 50.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-European-Mediterranean.

Carpocoris melanocerus (Mulsant & Rey, 1852)

Pentatoma melanocerus Mulsant & Rey, 1852: 109.



FIG. 12. — Male (A) and female (B) of *Aelia klugi* Hahn, 1833. Scale bar: 1 mm. Photo: G. Strauss.

NEW DATA. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 7 specimens, MZUF, MNHN. — Valdieri, Sant'Anna di Valdieri, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 28.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1625 m, 44.19503°N, 7.49107°E (WGS84), 17.VIII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1650 m, 26.IX.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1730 m, 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1920 m, 44.199977°N, 7.474896°E (WGS84), 26.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Carpocoris purpureipennis (De Geer, 1773)

Cimex purpureipennis De Geer, 1773: 258.

NEW DATA. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Palaearctic.

Dolycoris baccarum (Linnaeus, 1758)

Cimex baccarum Linnaeus, 1758: 445.

NEW DATA. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Limone Piemonte, Gias della Perla, La Perla small lake, 2036 m, 44.1542°N, 7.6183°E (WGS84), 27.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 4 specimens, MZUF, MNHN. — Limone Piemonte, La Perla, 2000 m, 44.15528°N, 7.1753°E (WGS84), 11.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1636 m, 44.27444°N, 7.30426°E (WGS84), 22.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 09.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 24.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m,



Fig. 13. — *Eurydema fieberi* Fieber, 1837. Scale bar: 2 mm. Photo: S. Bambi.

44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 17.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1920 m, 44.199977°N, 7.474896°E (WGS84), 26.VI.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Eurydema fieberi Fieber, 1837
(Fig. 13)

Eurydema fieberi Fieber, 1837: 353.

NEW DATA. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Centralasiatic-Europeano-Mediterranean.

REMARK

This record is interesting because this species normally lives at lower altitudes. Dusoulrier (pers. com.) also found it once at 1750 m, in the Southern Alps. Usually, its twin species *E. rotundicollis* (Dohrn, 1860) can be found above 1500 m

in this part of the Alps. Indeed, in the Italian Central Alps (e.g.) it was found at 1425 m on Brassicaceae, near the Livrio stream, Sondrio Province (Dioli 1974), while the twin species *E. rotundicollis* was quoted in the high mountains of the same area, at about 2000 m: “Lombardy: Sondrio, Arigna, Ghiacciaio delle Fasciere” (Mancini 1959).

Eurydema oleracea (Linnaeus, 1758)

Cimex oleraceus Linnaeus, 1758: 446.

NEW DATA. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF. — Valdieri, Colle dell’Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 30.VII.2013, MN, MD leg., 1 specimen, MZUF. — Valdieri, Sant’Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 17.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 3 specimens, MZUF, MNHN.

CHOROTYPE. — Sibero-European.

Eysarcoris venustissimus (Schrank, 1776)

Cimex venustissimus Schrank, 1776: 80.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 7 specimens, MZUF, MNHN.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Graphosoma italicum italicum (O. F. Müller, 1766)

Cimex italicus O. F. Müller, 1766: 190.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell’Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Valdieri, Sant’Anna di Valdieri, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, MN leg., 2 specimens, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 21.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 1 specimen, MNHN.

CHOROTYPE. — Palaearctic.

Holcostethus (Holcostethus) sphacelatus
(Fabricius, 1794)

Cimex sphacelatus Fabricius, 1794: 120.

NEW DATA. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 2 specimens, MZUF.

CHOROTYPE. — Europeano-Mediterranean.

Neottiglossa pusilla (Gmelin, 1790)

Cimex pusillus Gmelin, 1790: 2160.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcra bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Sibero-European.

Palomena prasina (Linnaeus, 1761)

Cimex prasinus Linnaeus, 1761: 249.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 29.VII.2015, LB & MN leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII. 2014, MN, MD leg., 2 specimens, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Valdieri, Terme di Valdieri, La Valletta, 1550 m, 44.19083°N, 7.26778°E (WGS84), 17.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1600 m, 44.19503°N, 7.49107°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Pentatoma rufipes (Linnaeus, 1758)

Cimex rufipes Linnaeus, 1758: 443.

NEW DATA. — Entracque, Fus Carbonaro, 1360 m, 44.23738°N, 7.38143°E (WGS84), 21.VII.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 2 specimens, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 1 specimen, MNHN. — San Giacomo di Entracque, Prà del Rasur, 1500 m, 44.17261°N, 7.39636°E (WGS84), 05.IX.2003, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1770 m, 44.2694°N, 7.30712°E (WGS84), 22.VII.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian del Valasco, 1700 m, 44.19608°N, 7.23051°E (WGS84), 27.VII.2012, MN leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri,



FIG. 14. — Female of *Sternodontus obtusus* Mulsant & Rey, 1856. Scale bar: 1 mm. Photo: G. Strauss.

980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1679 m, 44.2041°N, 7.48709°E (WGS84), 27.VII.2013, MN, MD leg., 1 juv., MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Peribalus (Peribalus) strictus vernalis
(Wolff, 1804)

Cimex vernalis Wolff, 1804: 140.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 1 specimen, MZUF. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 28.VI.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84) 15.VII.2014, MN, MD leg., 2 specimens, MZUF. — Valdieri, Valscura inferior lake, 2260 m, 44.19861°N, 7.20833°E (WGS84), 27.VII.2012, MN leg., 1 specimen, MZUF. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Picromerus bidens (Linnaeus, 1758)

Cimex bidens Linnaeus, 1758: 443.

NEW DATA. — Valdieri, Valasco, 1550 m, 44.204067°N, 7.256410°E (WGS84), 31.VIII.2013, MN, MD leg., 2 specimens, MZUF.

CHOROTYPE. — Holarctic.

Sciocoris (Sciocoris) cursitans cursitans
(Fabricius, 1794)

Naucoris cursitans Fabricius, 1794: 67.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 2 specimens, coll. CR; Entracque, Valle delle Rovine, 1733 m, 30.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 2 specimens, coll. CR.

NEW DATA. — Valdieri, Pian della Casa, 1650 m, 44.17446°N, 7.2672°E (WGS84), 7-16.VIII.2013, Malaise trap, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 17.VI.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1635 m, 44.187716°N, 7.492824°E (WGS84), 11.VII.2013, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 3 specimens, MZUF. — Vernante, loc. Palanfré, 1680 m, 44.2041°N, 7.48709°E (WGS84), 25.VII.2014, MN, MD leg., 5 specimens, MZUF. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 13.VI.2014, MN, MD leg., 2 specimens, MZUF, MNHN. — Vernante, loc. Palanfré, Gias Garbella, 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.

Sciocoris (Aposciocoris) homalonotus Fieber, 1851

Sciocoris homalonotus Fieber, 1851: 448.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 22.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1580 m, 44.19503°N, 7.49107°E (WGS84), 03.VII.2014, MN, MD leg., 2 specimens, MZUF. — Entracque, Fus Carbonaro, 950 m, 44.24899°N, 7.38166°E (WGS84), 06.VII.2014, MN, MD leg., 1 specimen, MZUF. — Limone Piemonte, La Perla, 1650 m, 44.17084°N, 7.61385°E (WGS84), 15.VII.2014, MN, MD leg., 3 specimens, MZUF, coll. PD. — Vernante, loc. Palanfré (Garbella), 1650 m, 44.19694°N, 7.4838°E (WGS84), 19.VI.2014, MN, MD leg., 3 specimens, MZUF, coll. PD.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Sternodontus obtusus Mulsant & Rey, 1856
(Fig. 14)

Sternodontus obtusus Mulsant & Rey, 1856: 120.

NEW DATA. — Valdieri, Colle dell'Arpione, 1750 m, 44.27762°N, 7.30121°E (WGS84), 30.VI.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Colle dell'Arpione, 1636 m, 44.27444°N, 7.30426°E (WGS84), 22.VII.2013, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — South-European.

Troilus luridus (Fabricius, 1775)

Cimex luridus Fabricius, 1775: 701.

NEW DATA. — Valdieri, Gias della Casa, pebble ponds and Gesso della Valletta stream, c. 1675 m, 44.174°N, 7.267°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Asiatic-European.

Family SCUTELLERIDAE Leach, 1815

Eurygaster austriaca austriaca (Schränk, 1776)

Cimex austriacus Schränk, 1776: 78.

LITERATURE. — Griffini (1893 [sub *Eurygaster nigro-cucullatus* (Goeze, 1778)]): Valdieri, Ghiliani leg.

NEW DATA. — Vernante, loc. Palanfré, 1920 m, 44.199977°N, 7.474896°E (WGS84), 26.VI.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 2120 m, 44.19726°N, 7.47119°E (WGS84), 27.VIII.2014, MN, MD leg., 1 specimen, MZUF. — Vernante, loc. Palanfré, 1800 m, 44.198119°N, 7.478328°E (WGS84), 30.VIII.2014, MN, MD leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Eurygaster dilaticollis Dohrn, 1860

Eurygaster dilaticollis Dohrn, 1860: 100.

LITERATURE. — EDIT 2010: Entracque, Bousset stream, Porcera bridge, 1083 m, 29.V-3.VII.2008, K. Wolf-Schwenninger & H. Schwenninger leg., 1 specimen, coll. CR.

CHOROTYPE. — Sibero-European.

Eurygaster maura (Linnaeus, 1758)

Cimex maurus Linnaeus, 1758: 441.

LITERATURE. — Griffini (1893 [sub *Eurygaster maurus*]): Valdieri, Ghiliani leg.

NEW DATA. — Entracque, Fus Carbonaro, 1000 m, 44.250810°N, 7.377315°E (WGS84), 21.VI.2014, MN, MD leg., 2 specimens, MZUF. — Entracque, S of Suffiet bridge, Bousset stream and right tributary, 1185 m, 44.1925°N, 7.4466°E (WGS84), 28.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Turano-Europeano-Mediterranean.

Eurygaster testudinaria (Geoffroy, 1785)

Cimex testudinarius Geoffroy, 1785: 195.

NEW DATA. — Entracque, Fus Carbonaro, 900 m, 44.24899°N, 7.38166°E (WGS84), 02.IX.2014, MN, MD leg., 1 specimen, MZUF. — Valdieri, Pian della Casa, 1700 m, 44.17446°N, 7.2672°E (WGS84), 26.VII.2012, MN leg., 1 specimen, MZUF. — Valdieri, Sant'Anna di Valdieri, Gesso della Valletta stream, 980 m, 44.2433°N, 7.3228°E (WGS84), 26-29.VII.2015, LB, FC, PD, MD, MN, SR, FT leg., 1 specimen, MZUF.

CHOROTYPE. — Palaearctic.




FIG. 15. — Chorotypes related to the species of Heteroptera located in the study area.

CHOROTYPE ANALYSIS

Chorotype analysis of all the considered species is shown in Figure 15. It is evident that the Holarctic and European widely-distributed components are prevalent if compared to the Mediterranean one, which is very limited. In fact, the most represented chorotypes are Palearctic (41 spp.), European (26 spp.), Asian-European (24 spp.) and Holarctic (22 spp.); strictly Mediterranean components are only two: Mediterranean (2 spp.) and East-Mediterranean (1 sp.) chorotypes.

Chorotypes were assigned to the widely distributed groups suggested by Vigna Taglianti *et al.* (1993, 1999); endemic components and cosmopolitan/subcosmopolitan elements were considered separately and treated as two further groups. The categories are: 1) widely distributed chorotypes in the Holarctic region; 2) widely distributed chorotypes in Europe; 3) widely distributed chorotypes in the Mediterranean basin; 4) Italian endemic elements; and 5) cosmopolitan and subcosmopolitan elements. Regarding the widely distributed Holarctic chorotypes, the most represented is the Palearctic one, representing 25%. Among the widely distributed chorotypes in Europe, the European elements are the most represented (76%). Finally, the Mediterranean component, less represented but very significant, is composed by two species with Mediterranean distribution and one with Eastern-Mediterranean distribution. The endemic elements are represented by three species limited to the Alps (*Stenodema algoviense*, *Acalypta visolensis visolensis*, and *Horwathia lineolata*), one extending to the Apennines (*Velia currens*), and an element confined to the south-western sector of the Alps: the Maritime Alps (*Dimorphocoris ruffoi*).

GENERAL DISCUSSION

This paper provides the first annotated checklist of the Hemiptera Heteroptera of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (IT1160056). With the present data, 146 species are added to the preliminary list obtained from literature, bringing the total number of Heteroptera species from 62 to 208, belonging to 135 genera and 25 families. The increase was, therefore, significant (230%). The family with the highest number of species is Miridae, with 90 species, which exactly reflects the proportion within Heteroptera at the Italian level; then Pentatomidae (20 spp.), Rhyparochromidae (15 spp.) and Lygaeidae *sensu stricto* (20 spp.). The other families include a lower number of species (< 10 spp.).

Research on Hemiptera Heteroptera was also conducted in the adjacent Mercantour National Park and a database including 234 species exists (EDIT 2010). However, at the moment it is not possible to easily compare the two protected areas because of the difference of surface and habitats present on the two slopes of the Maritime Alps, and also for the different sampling efforts and techniques used.

Considerations on the heteropterian fauna of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” follow. Concerning the aquatic component *sensu lato*, which comprises the infraorders Gerromorpha, Nepomorpha and Leptopodomorpha (see Polhemus & Polhemus 2008), in the study area the diversity is rather low for Gerromorpha (2 spp.) and Nepomorpha (3 spp.), while it is significant for Leptopodomorpha, with 9 species of Saldidae.

The cryophilic component, typically alpine (*Arctocoris carinata*, *Macrosaldula scotica*, *Saldula orthochila*, *Salda henschii*) or less cryophilic, mainly distributed at high altitude (*Gerris*

costae, *Velia currens*, *Saldula c-album*) are prevalent. There are also some eurythermic species (*Sigara lateralis*, *Notonecta viridis*, *Saldula saltatoria*) and other euryecious species, more common at low altitudes (*Chartoscirta cocksi*, *Macrosaldula variabilis*, *Saldula arenicola*, *Saldula melanoscela*) are present too. On the contrary, more thermophilic elements, typically Mediterranean (cf. Cianferoni *et al.* 2013), are absent. To confirm these results it is important to mention the absence of two very common aquatic species, also easily detectable: *Hydrometra stagnorum* (Linnaeus, 1758) from Gerromorpha, and *Nepa cinerea* Linnaeus, 1758 from Nepomorpha, that are most likely limited by the average low temperatures of the studied area. The altitudinal limit of such species in northern Italy is close to the lowest elevations present in the study area (cf. Bacchi & Rizzotti Vlach 2005, 2007). It is still possible that further research, especially at these altitudes, could lead to their discovery in the studied area. Concerning *Gerris costae* two subspecies are reported in Italy: *G. costae costae* (Herrich-Schäffer, 1850) which tends to be more alpine, and *G. costae fieberi* Stichel, 1938 which lives at low altitude. Their taxonomic status and distribution, however, is not yet very clear (see Cianferoni 2011 for a bibliographic summary; cf. Damgaard *et al.* 2014). Studies were performed on mitochondrial DNA and small differences were detected between the two entities, but the issue has not yet been finally resolved (cf. Damgaard 2006; Damgaard *et al.* 2014). Therefore, we prefer to consider the species *sensu* Cianferoni (2011) without indicating the subspecies. *Arctocoris carinata* is another typical alpine species. This taxon has a Boreo-alpine distribution, and has a Sibero-European chorotype (cf. Vigna Taglianti *et al.* 1993, 1999); the species has a nearly continuous distribution in northern Europe. There are also three isolated populations in the southern alpine regions, defined as glacial relicts: in the Pyrenees, in the Alps and in the Balkans (Jansson 1986). The species can be found in Italy throughout the Alpine region: Liguria, Piedmont, Aosta Valley, Lombardy, Trentino-Alto Adige, and Veneto (Mancini 1963, Bacchi & Rizzotti Vlach 2005, 2007) and there are some populations in the glacial lakes of the northern Apennines (Cianferoni & Terzani 2013). Although its occurrence would have been highly probable, this species had not been found yet in the Maritime Alps Nature Park.

Salda henschii was recently recorded for the first time in Italy (Cianferoni *et al.* 2017). This taxon belongs to a group of Saldidae which are strictly boreo-montane. It is mainly restricted to the mountains of central Europe, with some populations in southern Scandinavia; in the Arctic region the species is replaced by the congeneric *Salda sahlbergi* Reuter, 1875, which lives further north. The southern distribution limit of *S. henschii* was until now in Austria and Switzerland, with some apparently relict stations in the Alps (Lindskog 1991); its discovery in the Maritime Alps was completely unexpected and therefore rather exceptional. A closely related species, *Salda littoralis* (Linnaeus, 1758), is also present in Italy. Péricart (1990) writes that this taxon is present “seulement dans l’arc alpin, mais non dans les Alpes Maritimes” [= only in the alpine arc, but not in the Maritime Alps]. To date this species was not detected in the area, where it could be replaced by the congener *S. henschii*.

Saldula orthochila is also an interesting species. The northern limit of its distribution is the Arctic Circle, while in the south it reaches the northern sector of the Mediterranean region where, however, is present only in the mountains (Péricart 1990). Therefore in Italy it can be considered as a typical alpine species.

An important endemic species found during this research was *Dimorphocoris ruffoi* Tamanini, 1971. It is an endemic Italian Miridae, confined to the Maritime Alps. The *locus typicus* (Tamanini 1971) is the Mount Mandolè (which falls outside the studied area), but it was found later, also near the Refuge Soria (Tamanini 1976), inside the protected area. It is a vicariant species with the similar *Dimorphocoris gallicus* Wagner, 1957, which lives in the near Mercantour National Park (EDIT 2010). Members of *Dimorphocoris* Reuter, 1890, because of wing polymorphism (winged males or sometimes wingless, and females always wingless) suffer of a remarkable isolation, especially in the Alps, where valleys and mountain ranges are often insuperable barriers.

Another Mirid that deserves to be remembered is *Orthocephalus coriaceus* (Fabricius, 1777). It was considered doubtful for Italy in the last published checklist (Faraci & Rizzotti Vlach 1995) and recently confirmed by Carapezza & Faraci (2005, 2007) and found also in the studied area after this entomological survey. This species was found in the investigated areas, with a few specimens, in an alpine meadow above the village of Palanfrè, at an altitude of about 1700 metres. *Orthocephalus coriaceus* is a widespread species throughout Europe, and it is also quoted for North Africa. It is typically found in open and xerophilous areas, where it feeds on several Asteraceae Martinov, 1820 and occasionally on plants belonging to families Lamiaceae Martinov, 1820 and Rubiaceae Jussieu, 1789. This insect attacks the leaves of the host plant, and also the stem, inside which it lays its eggs. It overwinters as egg, and adults begin to appear in mid-June, remaining active until late July–early August (Wagner 1970).

Dicyphus annulatus (Wolff, 1804) is another interesting species; it is a sporadic Mirid which lives in Italy in the northern regions: there are reliable data only for Piedmont, Lombardy, and Liguria (Dioli 1997), but its distribution needs to be confirmed after its separation from *Dicyphus ononidis* Wagner, 1951.

Among the remarkable taxa we can also count the Pentatomidae *Sternodontus obtusus* Mulsant & Rey, 1856. This species was until now considered rare in Europe, with a few specimens in reference collections, since for a long time its host plant has been unknown. Lupoli (1994) reports the discovery of its host plant, *Laserpitium gallicum* (Apiaceae), which hosts eggs, nymphs and adults. This finding allowed us to understand the difficulties previously encountered in finding this species. In particular: mimicry of nymphs and adults, which can be confused with the seeds of *Laserpitium* L., as both share the same brown coloration; feeding period of adults limited to only three weeks approximately, that is in the same time of the development of the host plant seeds, which represent their main source of food. Moreover, the production of seeds may vary between mid-June and mid-September depending on the

areas, altitude and climate. During this study, *Sternodontus obtusus* was found for the first time in the summer of 2013. The discovery was made at Colle dell'Arpione (Valdieri), at an altitude of about 1750 metres. During the entomological survey in 2014 the research went on in the same area in order to confirm the presence of the species and gather new data, and two specimens were found. However, *Laserpitium gallicum* L., the host plant of this shield bug, has not been found in the investigated area. In that area was instead very abundant the Pig-Nut, *Bunium bulbocastanum* L. (Apiaceae), an heliophilous and mesoxerophilous plant typical of pastures and dry meadows on limestone substrate. Therefore it is very likely that this could be another host species for this shield bug. This observation, however, has still to be verified through further investigations. *Sternodontus obtusus* was reported for the South of France in the 1990s, while in Italy a small number of locations are known: Cavriana in Lombardia (Péricart 2010) and San Giorio di Susa (P. Dioli, unpublished data); other findings cited in the literature (Servadei 1967; Péricart 2010) date back more than fifty years ago and several are very old records. For this reason, the data of Colle dell'Arpione (Valdieri) has to be considered of importance. Further research would be needed to check in detail the current presence of this species, in the studied area.

Among the species found, others deserve to be mentioned as they represent interesting entities of the alpine horizon: *Geocoris lapponicus* Zetterstedt, 1838 (Geocoridae) (Fig. 10), *Ligyrocorys sylvestris* (Linnaeus, 1758) and *Trapezonotus desertus* Seidenstucker, 1951 (Rhyparochromidae), *Calocoris affinis* (Herrich-Schäffer, 1835) and *Calocoris alpestris* (Meyer-Dür, 1843) (Miridae), *Carpocoris melanocerus* (Mulsant & Rey, 1852) (Pentatomidae), and *Canthophorus impressus* (Horváth, 1881) (Cydnidae).

Some general considerations after the chorotype analysis can be done. The Maritime Alps, due to their geographical position relatively close to the sea, show a very important Mediterranean vegetation component. This condition occurs because typically Mediterranean vegetation “rises up” to higher altitudes (Gallino & Pallavicini 2000). Concerning the entomofauna, in particular the heteropteran component of the studied area, and in general of the Maritime Alps, it can be noticed that the continental component is prevalent, with several species having a Holarctic and European distribution, but it is important to highlight that there are also Mediterranean species (3), even if their number is much lower than that of plants. All the Mediterranean species belong to the Miridae family: the strictly Mediterranean species are *Adelphocoris vandalicus* (Rossi, 1790) and *Pachytomella passerinii* (A. Costa, 1842), whereas *Closterotomus annulus* (Brullé, 1832) is confined in the eastern Mediterranean.

Among the species previously cited, it is important to remark the presence of some endemic elements. First of all, *Dimorphocoris ruffoi*, an endemic of the Maritime Alps. Moreover, three endemics of the Alps are present, *Stenodema algoviense* (Schmidt, 1934), *Acalypta visolensis visolensis* Péricart, 1976, and *Horwathia lineolata* (A. Costa, 1852). Another endemic species is *Velia currens*, whose distribution includes the Alps and the Apennines, up to the mountains of Calabria, in Italy (Cianferoni & Mazza 2012; Cianferoni & Santini 2012).

It is important also to remark the presence of three recently described mirid species – *Dimorphocoris poggii* Carapezza, 2002, *Cremnocephalus matocqi* Magnien, 2000, and *Plinththis heteroclitus* Matocq & Pluot-Sigwalt, 2013, which are considered endemic for the Ligurian Alps.

In the examined literature further two species, that we decided not to include in the checklist, are quoted. The first case is *Peirates hybridus* (Scopoli, 1763), belonging to the family Reduviidae. This species was reported by Noelli (1897a) for the locality “Colle delle Finestre”. It was not possible to verify whether it is a name which falls in the studied area, or if it refers to the same locality in the Susa Valley (Turin). Therefore, this species has been excluded from the list. The second doubtful case is related to *Himacerus (Stalia) boops* Schiödte, 1870, reported by Servadei (1957). There are no further records of this species for Italy, and we did not find it during the research. The records of the last century (Stichel 1959, 1960), then quoted by Servadei (1967), were not confirmed by the following reviews (Faraci & Rizzotti Vlach 1987, Péricart 1987). They can be therefore interpreted as misidentifications: the record probably refers to *Nabis flavomarginatus* Scholtz, 1847.

A good knowledge of the biodiversity of a protected area plays a key role in the conservation of nature. Moreover, the development of faunistic studies and the elaboration of species checklists may help stakeholders in the management of natural resources in order to protect them. In this paper we showed the high Heteroptera biodiversity of the Site of Community Importance and Special Area of Conservation “Alpi Marittime”, and we are confident that in this territory more species have still to be discovered.

Acknowledgements

We wish to thank our friend and colleague Marco Dogliotti, Naturalistic Guide of the Maritime Alps Nature Park, for helping us in the field collections, for the logistic support throughout the research period, and for the photos kindly provided. The work has benefited from the collaboration of the Maritime Alps Nature Park, under the aegis of the project “Piano Integrato Transfrontaliero Marittime-Mercantour”, funded by Alcotra 2007-2013 program; thanks to all the staff of the Maritime Alps Nature Park, and especially to Luca Giraud, Cati Caballo and Laura Martinelli, who gave us the authorisation for the research and for the logistic support. We also warmly thank Luca Bartolozzi, Saverio Rocchi, and Fabio Terzani (MZUF) for their help in the field; Luca Bartolozzi also revised the final version of the manuscript. Saulo Bambi (MZUF) kindly took the photo of *Eurydema fieberi*. Thanks are also due to the entomologists who provided us part of the data, mainly Attilio Carapezza (University of Palermo, Italy) and Roberto Poggi (MSNG). A special thank to Antoine Mantilleri (MNH) for the translation of the French abstract, and to the referees François Dusoulier (MNH) and Petr Kment (National Museum, Prague, Czech Republic) for their very useful comments. Finally, we are very grateful to Gerhard Strauss (Biberach, Germany), author of the electronic catalog “Corisa”, for granting us the use of his excellent photos.

REFERENCES

- AUKEMA B. & RIEGER C. 1995. — *Catalogue of the Heteroptera of the Palaearctic Region, Vol. 1, Enicocephalomorpha, Dipsocoromorpha, Nepomorpha, Gerromorpha and Leptodomorpha*. The Netherlands entomological Society, Amsterdam, XXVI, 222 p.
- AUKEMA B. & RIEGER C. 1996. — *Catalogue of the Heteroptera of the Palaearctic Region, Vol. 2, Cimicomorpha I*. The Netherlands entomological Society, Amsterdam, XIV, 361 p.
- AUKEMA B. & RIEGER C. 1999. — *Catalogue of the Heteroptera of the Palaearctic Region, Vol. 3, Cimicomorpha II*. The Netherlands entomological Society, Amsterdam, XIV, 577 p.
- AUKEMA B. & RIEGER C. 2001. — *Catalogue of the Heteroptera of the Palaearctic Region, Vol. 4, Pentatomorpha I*. The Netherlands entomological Society, Amsterdam, XIV, 346 p.
- AUKEMA B. & RIEGER C. 2006. — *Catalogue of the Heteroptera of the Palaearctic Region, Vol. 5, Pentatomorpha II*. The Netherlands entomological Society, Amsterdam, XIII, 550 p.
- AUKEMA B., RIEGER C. & RABITSCH W. 2013. — *Catalogue of the Heteroptera of the Palaearctic Region, Vol. 6, Supplement*. The Netherlands entomological Society, Amsterdam, XXIV, 629 p.
- BACCHI I. & RIZZOTTI VLACH M. 2005. — Insecta Heteroptera Nepomorpha e Gerromorpha (p. 147-149, CD-ROM), in RUFFO S. & STOCH F. (eds). *Checklist e distribuzione della fauna italiana*. Memorie del Museo Civico di Storia Naturale di Verona, 2. serie, Sezione Scienze della Vita 16, 307 p.
- BACCHI I. & RIZZOTTI VLACH M. 2007. — Insecta Heteroptera Nepomorpha e Gerromorpha (p. 147-149, CD-ROM), in RUFFO S. & STOCH F. (eds). *Checklist and Distribution of the Italian fauna*. Memorie del Museo Civico di Storia Naturale di Verona, 2. serie, Sezione Scienze della Vita 17 (2006), 303 p.
- BECKER A. 1864. — Naturhistorische Mitteilungen. *Bulletin de la Société des Naturalistes de Moscou* 37 (9): 477-493.
- BOHEMAN C. H. 1852. — Nya svenska Hemiptera. *Öfversigt af Kongliga Vetenskapsakademiens Förhandlingar* 9: 65-80.
- BONELLI S., BARBERO F., CASACCI L. P., CERRATO C. & BALLETO E. 2015. — The butterfly fauna of the Italian Maritime Alps: results of the EDIT project, in DAUGERON C., DEHARVENG L., ISAIA M., VILLEMANT C. & JUDSON M. (eds). *Mercantour/Alpi Marittime All Taxa Biodiversity Inventory*. *Zoosystema* 37 (1): 139-167. <https://doi.org/10.5252/z2015n1a6>
- BRULLÉ A. 1832. — Expédition scientifique de Morée. Section des Sciences Physiques. Tome III, 1^{re} Partie. Zoologie. Deuxième Section – Des animaux articulés. Levrault Imprimeur, Paris, Strasbourg, 402 p.
- CARAPEZZA A. 2007. — Gli Eterotteri (Heteroptera), in NARDI G. & VOMERO V. (eds). *Artropodi del Parco Nazionale del Vesuvio: Ricerche Preliminari. Conservazione Habitat Invertebrati* 4: 87-97.
- CARAPEZZA A. & FARACI F. 2005. — Insecta Heteroptera Leptopodidae, Saldidae, Miridae (partim), Tingidae (p. 151-153, CD-ROM), in RUFFO S. & STOCH F. (eds). *Checklist e distribuzione della fauna italiana*. Memorie del Museo Civico di Storia Naturale di Verona, 2. serie, Sezione Scienze della Vita 16, 307 p.
- CARAPEZZA A. & FARACI F. 2007. — Insecta Heteroptera Leptopodidae, Saldidae, Miridae (partim), Tingidae (p. 151-153, CD-ROM), in RUFFO S. & STOCH F. (eds). *Checklist and Distribution of the Italian fauna*. Memorie del Museo Civico di Storia Naturale di Verona, 2. serie, Sezione Scienze della Vita 17 (2006), 303 p.
- CIANFERONI F. 2011. — Notes on Gerromorpha, Nepomorpha and Leptopodomorpha from Sardinia (Hemiptera, Heteroptera), in NARDI G., WHITMORE D., BARDIANI M., BIRTELE D., MASON F., SPADA L. & CERRETTI P. (eds). *Biodiversity of Marganai and Montimannu (Sardinia). Research in the framework of the ICP Forests network*. *Conservazione Habitat Invertebrati*, 5. Cierre Edizioni, Sommacampagna, Verona: 255-268.
- CIANFERONI F. & MAZZA G. 2012. — The aquatic Heteroptera (Insecta: Hemiptera) of the “Foreste Casentinesi, Monte Falterona e Campigna” National Park (Central Italy). *Zootaxa* 3568: 36-52. <https://doi.org/10.11646/zootaxa.3568.1.2>
- CIANFERONI F. & SANTINI G. 2012. — Ecology and life histories of two Alpine-Apenninic species of *Velia* (Hemiptera: Heteroptera: Veliidae). *European Journal of Entomology* 109: 427-434. <https://doi.org/10.14411/eje.2012.055>
- CIANFERONI F. & TERZANI F. 2013. — Nuovi dati su Gerromorpha e Nepomorpha in Italia (Hemiptera Heteroptera). *Bollettino della Società Entomologica Italiana* 145 (2): 51-57. <https://doi.org/10.4081/BollettinoSEI.2013.51>
- CIANFERONI F., TERZANI F. & ROCCHI S. 2013. — Nepomorpha, Gerromorpha and Leptopodomorpha (Hemiptera: Heteroptera) of the Tuscan Archipelago (Italy). *Zootaxa* 3669 (3): 302-320. <https://doi.org/10.11646/zootaxa.3669.3.6>
- CIANFERONI F., NORBIATO M. & DOGLIOTTI M. 2017. — First records in Italy of the red-listed shore bug *Salda henschii* (Reuter, 1991) (Hemiptera: Heteroptera: Saldidae). *Zootaxa* 4238 (4): 597-598. <https://doi.org/10.11646/zootaxa.4238.4.9>
- COSTA A. 1842. — Mémoire pour servir à l'histoire des Hémiptères Héteroptyères de Deux-Siciles. *Annales de la Société entomologique de France* 10: 279-308.
- COSTA A. 1847. — Cimicum regni Neapolitani centuria secunda, decas 1-5. *Atti del Reale Istituto di Incoraggiamento delle Scienze Naturali di Napoli* 7: 239-279.
- COSTA A. 1862. — Additamenta ad Centurias Cimicum Regni Neapolitani. (Preprint from 1863, *Atti del Reale Istituto di Incoraggiamento delle Scienze Naturali di Napoli* 10: 329-367).
- COSTA O. G. 1834. — Enumeratio Insectorum rariorum quae per Apuliam, Agrum Puteolanum, Insulae Prochytae et Pithecusae ac in monte Vesubio perlegi, anno 1834. *Annuario Zoologico* 12: 69-81.
- CURTIS J. 1835. — *British Entomology: being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found. Vol. VII. Homoptera. Hemiptera. Aphaniptera*. London, printed by the author: pls 530-571. <https://doi.org/10.5962/bhl.title.8148>
- DAMGAARD J. 2006. — Phylogeny and mtDNA phylogeography of two widespread European pond skater species (Hemiptera-Heteroptera: Gerridae: *Gerris* Fabricius). *Insect Systematics and Evolution* 37: 335-350. <https://doi.org/10.1163/187631206788838554>
- DAMGAARD J., MOREIRA F. F. F., WEIR T. A. & ZETTEL H. 2014. — Molecular phylogeny of the pond skaters (Gerrinae), discussion of the fossil record and a checklist of species assigned to the subfamily (Hemiptera: Heteroptera: Gerridae). *Insect Systematics and Evolution* 45: 251-281. <https://doi.org/10.1163/1876312X-44042105>
- DE GEER C. 1773. — *Mémoires pour servir à l'histoire des Insectes*, 3. Pierre Hesselberg, Stockholm, 696 p. <https://catalogue.bnf.fr/ark:/12148/cb37256985v>
- DECKERT J. 1985 — Über *Lygaeus simulans* spec. nov. und *L. equestris* (Linnaeus, 1758), zwei nahe verwandte paläarktische Lygaeinae (Heteroptera, Lygaeidae). *Mitteilungen aus dem Zoologischen Museum in Berlin* 61 (2): 273-278. <https://doi.org/10.1002/mmnz.19850610203>
- DEHARVENG L., BEDOS A., DAUGERON C., VILLEMANT C. & JUDSON M. L. I. 2015. — Organization, usefulness and limitations of an ATBI (All Taxa Biodiversity Inventory): the inventory of terrestrial invertebrates in the Mercantour National Park, in DAUGERON C., DEHARVENG L., ISAIA M., VILLEMANT C. & JUDSON M. (eds), *Mercantour/Alpi Marittime All Taxa Biodiversity Inventory*. *Zoosystema* 37 (1): 9-30. <https://doi.org/10.5252/z2015n1a1>
- DEL COURT A. 1909. — Recherches sur la variabilité du genre «*Notonecta*». *Bulletin de la Société entomologique de Belgique* 43: 313-461.
- DIOLI P. 1974. — Emitteri Eterotteri nuovi o poco noti della Valtellina (Hemiptera, Heteroptera). *Memorie della Società Entomologica Italiana* 53: 30-38.
- DIOLI P. 1997. — Eterotteri (Heteroptera) del Monte Barro (Italia, Lombardia, Lecco). *Memorie della Società Italiana di Scienze Natu-*

- rali e Museo Civico Storia Naturale di Milano 27 (2): 159-174.
- DIOLI P., FARACI F. & RIZZOTTI VLACH M. 1994. — Osservazioni morfologiche e corologiche su *Lygaeus simulans* Deckert, 1985 in Italia (Heteroptera: Lygaeidae). *Atti XVII Congresso nazionale italiano di Entomologia*, Udine (13-18 giugno 1994): 195-198.
- DOHRN A. 1860. — Hemipterologische Miscellaneen. *Entomologische Zeitung Stettin* 21: 99-109.
- DOHRN A. 1862. — Drei neue europäische Hemiptera. *Stettiner Entomologische Zeitung* 23: 210-211.
- DONOVAN E. 1798. — *The natural history of British insects; explaining them in their several states, with the periods of their transformations, their food, oeconomy, &c. together with the history of such minute insects as require investigation by the microscope. The whole illustrated by coloured figures, designed and executed from living specimens. Vol. 7.* London, printed by the author, pls CCXLII-CCLII, 96 p.
- DOUGLAS J. W. & J. SCOTT. 1868. — List of captures of Hemiptera in Palestine and Syria; together with descriptions of several new species. *Entomologist's Monthly Magazine* 5: 114-118, 135-139.
- DUFOUR L. 1833. — Recherches anatomiques et physiologiques sur les Hémiptères, accompagnées de considérations relatives de l'histoire naturelle et de la classification de ces insectes. *Mémoires présentés par divers Savants de l'Académie des Sciences de l'Institut de France* 4: 129-462. <https://doi.org/10.5962/bhl.title.12418>
- EDIT 2010. — ATBI+M: All Taxa Biodiversity Inventories + Monitoring in the Mercantour/Alpi Marittime nature Parks. Available at: <http://www.atbi.eu/mercantour-marittime/> [last access: 3 January 2018].
- EHANNO B. 1987. — *Les Hétéroptères Mirides de France : Tome II A : Inventaire et synthèses écologiques*. Muséum national d'Histoire naturelle, Paris, 648 p. (Inventaires de faune et flore ; 40).
- FABRICIUS J. C. 1775. — *Systema entomologiae sistens insectorum classes, ordines, genera, species; adiectis synonymis, locis, descriptionibus et observationibus*. Flensburgi et Lipsiae, XXXII + 832 p. <https://doi.org/10.5962/bhl.title.36510>
- FABRICIUS J. C. 1777. — *Genera Insectorum eorumque characteres naturales secundum numerum figuram situm et proportionem omnium partium oris adiecta mantissa specierum nuper detectarum*. Chilonii - Litteris Nuch. Friedr. Bartschii. XIV + 310 p. <https://doi.org/10.5962/bhl.title.119827>
- FABRICIUS J. C. 1781. — *Species Insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosis adiectis observationibus, descriptionibus*. Vol. 2. Impensis Carol. Ernest. Bohnii, Hamburgi et Kilonii, 517 p. <https://doi.org/10.5962/bhl.title.119827>
- FABRICIUS J. C. 1787. — *Mantissa Insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus*. Vol. 2. Impensis Christ. Gottl. Proft, Hafniae, 382 p. <https://doi.org/10.5962/bhl.title.36471>
- FABRICIUS J. C. 1794. — *Entomologia systematica emendata et aucta, secundum classes, ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus*. 4: I-V. Impensis Christ. Gottl. Proft, Hafniae, 472 p. <https://doi.org/10.5962/bhl.title.125869>
- FABRICIUS J. C. 1798. — *Entomologia systematica emendata et aucta, secundum classes, ordines, genera, species, adiectis synonymis, locis, observationibus*. Supplementum. Proft et Storch, Hafniae, II + 572 p. <https://doi.org/10.5962/bhl.title.125869>
- FABRICIUS J. C. 1803. — *Systema Rhynogotorum secundum ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus*. C. Reichard, Brunsvigae, X + 335 p. <https://doi.org/10.5962/bhl.title.11644>
- FALLÉN C. F. 1807. — *Monographia Cimicum Sueciae*. C. G. Proft, Hafniae, 123 p.
- FALLÉN C. F. 1829. — *Hemiptera Sueciae. Sectio prior. Hemelytrata*. Gothorum, Londini, 188 p.
- FARACI F. & RIZZOTTI VLACH M. 1987. — I Nabidi italiani (Hemiptera, Heteroptera). *Bollettino del Museo di Storia naturale, Verona*, 13 (1866): 81-138.
- FARACI F. & RIZZOTTI VLACH M. 1992. — I Leptopodomorpha italiani. Catalogo topografico con considerazioni ecologiche, fenologiche e zoogeografiche (Heteroptera). *Memorie della Società entomologica italiana* 70 (2) (1991): 33-102.
- FARACI F. & RIZZOTTI VLACH M. 1995. — Fascicolo 41. Heteroptera, in MINELLI A., RUFFO S. & LA POSTA S. (eds). *Checklist delle specie della fauna italiana*. Calderini, Bologna: 1-56.
- FERRARI E. 1884. — Materiali per lo studio della fauna tunisina raccolti da G. e L. Doria. V. Rincoti. *Annali del Museo Civico di Storia Naturale di Genova. Serie 2a*, 1 (XXI): 439-522. <https://biodiversitylibrary.org/page/29846142>
- FIEBER F. X. 1837. — Beiträge zur Kenntniss der Schnabelkerfe (Rhynchota). *Beiträge zur Gesammten Natur- und Heilwissenschaft* 1: 97-111, 337-355.
- FIEBER F. X. 1851. — Rhynchotographien. Drei monographische Abhandlungen. *Abhandlungen der Böhmischen Gesellschaft der Wissenschaften* 5 (7): 424-488.
- FIEBER F. X. 1858. — Kriterien zur generischen Theilung der Phytocoriden (Capsini auct.). *Wiener entomologische Monatschrift* 2: 289-327, 329-347, 388, 1 pl.
- FIEBER F. X. 1859. — Die europäischen Arten der Gattung SaIda Fab. *Wiener Entomologische Monatschrift* 3: 230-241.
- FIEBER F. X. 1861. — Die europäischen Arten der Gattung Salda Fab. *Wiener Entomologische Monatschrift* 3: 230-241.
- GALLINO B. & PALLAVICINI G. 2000. — *La vegetazione delle Alpi Liguri e Marittime*. Blu edizioni, Torino, 223 p.
- GEOFFROY E. L. 1785. — [Description of new taxa], in FOURCROY A. F. DE (ed). *Entomologia Parisiensis; sive Catalogus Insectorum quae in Agro Parisiensi reperiuntur; secundum methodum Geoffroeanum in sectiones, genera et species distributus, cui addita sunt nomina trivialia et fere trecentae novae species. Pars prima*. Paris, 231 p.
- GERMAR E. F. 1817. — *Reise durch Österreich, Tyrol nach Dalmatien und in das Gebiet von Ragusa*. 2. Brockhaus, Leipzig, 323 p.
- GMELIN J. F. 1790. — *Hemiptera. Caroli a Linné Systema Naturae; Ed. 13 aucta, reformata. Tom. 1, pars IV*. Lipsiae: 2041-2224.
- GOEZE J. A. E. 1778. — *Entomologische Beyträge zu des Ritter Linné zwölften Ausgabe des Natursystems. Zweyter Theil*. Weidmanns Erben und Reich, Leipzig, 352 p. <https://doi.org/10.5962/bhl.title.45974>
- GRIFFINI A. 1893. — Pentatomidi e i Coreidi del Piemonte. *Annali della Reale Accademia di Agricoltura, Torino* 36: 61-104.
- HAHN C. W. 1832. — *Die Wanzenartigen Insekten. Getreu nach der Natur abgebildet und beschrieben*. C. H. Zeh'schen Buchhandlung, Nürnberg 1 (2): 37-80.
- HAHN C. W. 1833. — *Die Wanzenartigen Insekten. Getreu nach der Natur abgebildet und beschrieben*. C. H. Zeh'schen Buchhandlung, Nürnberg 1 (4): 119-158.
- HAHN C. W. 1834. — *Die Wanzenartigen Insekten. Getreu nach der Natur abgebildet und beschrieben*. C. H. Zeh'schen Buchhandlung, Nürnberg 2 (4): 81-100.
- HAHN C. W. 1835. — *Die Wanzenartigen Insekten. Getreu nach der Natur abgebildet und beschrieben*. C. H. Zeh'schen Buchhandlung, Nürnberg 3 (1): 1-16.
- HENRY T. J. 1997. — Phylogenetic analysis of family groups within the infraorder Pentatomomorpha (Hemiptera: Heteroptera), with emphasis on the Lygaeoidea. *Annals of the Entomological Society of America* 90: 275-301. <https://doi.org/10.1093/aesa/90.3.275>
- HERRICH-SCHÄFFER G. A. W. 1835. — *Nomenclator emomologicus. Verzeichniss der europäischen Insecten; zur Erleichterung des Tauschverkehrs mit Preisen versehen. Heft; Lepidoptera und Hemiptera, letztere synoptisch bearbeitet und mit vollständiger Synonymie. I-IV*. Pustet, Regensburg, 116 p.
- HERRICH-SCHÄFFER G. A. W. 1836-1853. — *Die wanzenartigen Insekten* 3 (1836b): 33bis, 34bis, 35-114; 4 (1837): 1-32, (1838): 33-92, (1839a): 93-108; 5 (1839b): 1-60, (1840a): 61-108; 6 (1840b): 1-36, (1841): 37-72, (1842): 73-118; 7 (1843): 17-40, (1844): 41-134; 8 (1845): 1-48, (1846): 49-100, (1847): 101-130; 9 (1849): 1-44, (1850): 45-256; (1851): 257-348; (1853a,

- Historische Uebersicht der einschlägigen Literatur): 1-31; (1853b, Index alphabeticum-synonymicus): 1-210. Zeh (vols 3-6) & Lotzbeck (vols 7-9), Nürnberg.
- HERRICH-SCHÄFFER G. A. W. 1850 — *Die Wanzenartigen Insecten*. 6. Lotzbeck, Nürnberg: 41-256. <https://doi.org/10.5962/bhl.title.11547>
- HORVÁTH G. 1881. — Hemipterologiai Közlemények. *Természetrájszi Füzetek* 4 (1880): 184-192.
- ISAIA M., PASCHETTA M. & CHIARLE A. 2015. — Annotated checklist of the spiders (Arachnida, Araneae) of the Site of Community Importance and Special Area of Conservation “Alpi Marittime” (NW Italy), in DAUGERON C., DEHARVENG L., ISAIA M., VILLEMANT C. & JUDSON M. (eds). Mercantour/Alpi Marittime All Taxa Biodiversity Inventory. *Zoosystema* 37 (1): 57-114. <https://doi.org/10.5252/z2015n1a4>
- JAKOVLEV B. E. 1877. — New bugs, Hemiptera Heteroptera of the Astrakhan fauna. Second Supplement. *Bulletin de la Société des Naturalistes de Moscou* 52 (2): 269-300.
- JANSSON A. 1986. — The Corixidae (Heteroptera) of Europe and some adjacent regions. *Acta Entomologica Fennica* 47: 1-94.
- KOLENATI F. A. 1845. — *Meletemata Entomologica II. Hemiptera Caucasi Tesseratomidae monographice dispositae*. Academia Scientiarum, Petropoli: 1-132.
- KIRSCHBAUM C. L. 1856. — Rhynchographische Beiträge. *Jahrbuch des Vereins für Naturkunde im Herzogthum Nassau* 10: 163-348.
- KMENT P. & BAÑAR P. 2012. — True bugs (Hemiptera: Heteroptera) of the Bílé Karpaty Protected Landscape Area and Biosphere Reserve (Czech Republic) (p. 323-628). *Acta Musei Moraviae, Scientiae Biologicae* 96 (2) (2011): 1-933.
- KOLENATI F. A. 1846. — *Meletemata Entomologica IV. Hemiptera Caucasi. Pentatomidae monographice dispositae*. Academia Scientiarum, Petropoli, II + 72 p.
- LEACH W. E. 1817 — On the classification of the natural tribe of insects Notonectides with descriptions of British species. *Transactions of the Linnean Society of London* 12: 10-18. <https://doi.org/10.1111/j.1095-8339.1817.tb00215.x>
- LINDSKOG P. 1991. — Taxonomy and distribution of the boreo-montane shore bugs *Salda sahlbergi* and *S. henschii* (Heteroptera, Saldidae). *Entomologisk tidskrift*, 112: 1-18.
- LINNAEUS C. 1758. — *Systema naturae per regna tria naturae. secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata: I-V. Salvii, Holmiae*, 824 p. <https://doi.org/10.5962/bhl.title.542>
- LINNAEUS C. 1761. — *Fauna Suecica sistens animalia Sueciae regni; quadrupedia, aves, amphibia, pisces, insecta, vermes distributa per classes, et ordines, genera et species, cum differentiis specierum, synonymis autorum, nominibus incolarum, locis habitationum, descriptionibus insectorum*. Stockholmiae, XLVI + 578 p. <https://doi.org/10.5962/bhl.title.63899>
- LINNAEUS C. 1767. — *Systema Naturae. Ed. 12. Vol. 1, pt. 2. L. Salvii, Homiae*: 533-1327. <https://doi.org/10.5962/bhl.title.559>
- LUPOLI R. 1994. — Les punaises des Ombellifères de France. *Insectes* 93: 10.
- MANCINI C. 1952. — *Miscellanea Emitterologica italiana. I. Bollettino della Società entomologica italiana* 82: 56-64.
- MANCINI C. 1959. — *Miscellanea emitterologica. Annali del Museo Civico di Storia Naturale, Genova* 71 (1960): 102-110.
- MANCINI C. 1960. — *Miscellanea Emitterologica. Annali del Museo Civico di Storia Naturale “Giacomo Doria”, Genova* 71: 102-110.
- MANCINI C. 1963. — *Res Ligusticae CXXXIII. Emitteri Eteroteri della Liguria. Annali del Museo Civico di Storia Naturale, Genova* 74: 30-121.
- MEYER-DÜR L. R. 1843. — *Verzeichniss der in der Schweiz einheimischen Rhynchoten (Hemiptera Linn.). Erstes Heft. Die Familie der Capsini*. Jent & Gassmann, Solothurn. X + 11-116 + IV p., 7 pls.
- MINELLI A., RUFFO S. & VIGNA TAGLIANTI A. 2006. — The Italian faunal provinces, in RUFFO S. & STOCH F. (eds). *Checklist and Distribution of the Italian Fauna*. Memorie del Museo Civico di Storia Naturale di Verona 17: 37-39.
- MÜLLER O. F. 1766. — [Description of new taxa], in ALLIONI C. (ed.). *Manipulus insectorum Taurinensium*. Mélanges de Philosophie et de Mathématique de la Société Royale de Turin 3: 185-198.
- MULSANT E. & REY C. 1852. — Description de quelques Hémiptères Hétéroptères nouveaux ou peu connus. *Annales de la Société Linnéenne de Lyon* 1850/1852: 76-141.
- MULSANT E. & REY C. 1856. — Description de quelques Hémiptères-Hétéroptères nouveaux ou peu connus. *Annales de la Société Linnéenne de Lyon, n. s. 3*: 118-131.
- NOELLI A. 1897a. — I Reduvidi del Piemonte. *Annali della Reale Accademia d'Agricoltura, Torino* 40: 33-48.
- NOELLI A. 1897b. — I Ligeidi del Piemonte. *Annali della Reale Accademia d'Agricoltura, Torino* 40: 163-192.
- PANZER G. W. F. 1797. — *Faunae insectorum Germanicae initia oder Deutschlands Insecten gesammelt und herausgegeben. Heft 39-48*. Felssecker Nürnberg.
- PANZER G. W. F. 1804. — *Faunae insectorum Germanicae initia oder Deutschlands Insecten gesammelt und herausgegeben. Heft 86-96*. Felssecker Nürnberg.
- PÉRICART J. 1972. — *Hémiptères Anthocoridae, Cimicidae et Microphysidae de l'Ouest-Paléarctique. Faune de l'Europe et du Bassin Méditerranéen*. 7. Masson et Cie, Paris, 402 p.
- PÉRICART J. 1976. — Une espèce nouvelle d'*Acalypta* des Alpes occidentales: *Acalypta visolensis* n. sp. *Bollettino della Società Entomologica Italiana* 108: 174-177.
- PÉRICART J. 1983. — *Hémiptères Tingidae euro-méditerranéens, Faune de France*. 69. Fédération française des Sociétés de Sciences naturelles, Paris, 618 p.
- PÉRICART J. 1987. — *Hémiptères Nabidae d'Europe occidentale et du Maghreb. Faune de France* 71. Fédération française des Sociétés de Sciences naturelles, Paris, 185 p.
- PÉRICART J. 1990. — *Hémiptères Saldidae et Leptopodidae d'Europe occidentale et du Maghreb. Faune de France*. 77. Fédération française des Sociétés des Sciences naturelles, Paris, 238 p.
- PÉRICART J. 1998. — *Hémiptères Lygaeidae euro-méditerranéens. Seconde partie: Systématique, II: Oxycaeninae, Bledionotinae, Rhyparochrominae I. Faune de France, Montpellier: vol. 84B, 1998, VI + 453p., 148 pl. de fig., 88 cartes*.
- PÉRICART J. 1999a. — *Hémiptères Lygaeidae euro-méditerranéens. Faune de France*. 84A. Fédération française des Sociétés de Sciences naturelles, Paris (1998), 468 p.
- PÉRICART J. 1999b. — *Hémiptères Lygaeidae euro-méditerranéens. Faune de France*. 84B. Fédération française des Sociétés de Sciences naturelles, Paris (1998), 453 p.
- PÉRICART J. 1999c. — *Hémiptères Lygaeidae euro-méditerranéens. Faune de France*. 84C. Fédération française des Sociétés de Sciences naturelles, Paris (1998), 487 p.
- PÉRICART J. 2010. — *Hémiptères Pentatomoidea euro-méditerranéens. Volume 3. Systématique: troisième partie. Sous-familles Podopinae et Anisopinae. (Faune de France 93)*. Fédération française des Sociétés de Sciences naturelles, Paris, 291 p., 24 pls.
- PODA N. 1761. — *Insecta Musei Graecensis, quae in ordines, genera et species juxta sistema naturae Caroli Linnaei*. Widmanstadius, Graz, 127 p.
- POLHEMUS J. T. & POLHEMUS D. A. 2008. — Global diversity of true bugs (Heteroptera; Insecta) in freshwater. *Hydrobiologia*, 595: 379-391. <https://doi.org/10.1007/s10750-007-9033-1>
- PUTON A. 1877. — Note sur le genre *Peritrechus* Fieb. *Petites nouvelles entomologiques* 2 (9), 168: 117.
- PUTON A. 1884. — Hémiptères nouveaux. *Revue d'Entomologie*, 3: 85-88, 142-149, 312-314.
- PUTON A. 1887. — Hémiptères nouveaux ou peu connus de la faune Paléarctique. *Revue d'Entomologie* 6: 96-105.
- REMANE R. 1949. — *Nabis (Reduviolus) pseudoferus* nova species, eine neue deutsche Nabis-Art. *Verhandlungen des Vereins für naturwissenschaftliche Heimatforschung zu Hamburg* 30: 65-67.
- REMANE R. 1955. — *Lygus (Exolygus) wagneri* nov. spec., eine weitere europäische Exolygus-Art. *Zoologischer Anzeiger* 155: 115-119.

- REUTER O. M. 1870. — Pargas sockens Heteroptera, förtecknade. *Notiser ur Sällskapet pro Fauna et Flora Fennica förhandlingar*, 11 (1871): 309-326.
- REUTER O. M. 1873. — Bidrag till nordiska Capsiders synonymi. *Notiser ur Sällskapet pro Fauna et Flora Fennica förhandlingar* 14: 1-25.
- REUTER O. M. 1875a. — Remarques synonymiques sur quelques Héteroptyères. *Annales de la Société entomologique de France* (5) 4 (1874): 559-566.
- REUTER O. M. 1875b. — Genera Cimicidarum Europae. *Bihang till Kongliga Svenska Vetenskapsakademiens Förhandlingar* 3 (1): 1-66.
- REUTER O. M. 1875c. — Hemiptera Heteroptera Austriaca, sum. Maji-Augusti 1870 a J. A. Palmén collecta. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, 25 (1876): 83-88.
- REUTER O. M. 1883a. — The British species of *Dicyphus*. *Entomologist's Monthly Magazine* 20: 49-53.
- REUTER O. M. 1883b. — Hemiptera Gymnocerata Europae. Hémiphtères Gymnocérates d'Europe, du bassin de la Méditerranée et de l'Asie Russe. III. *Acta Societatis Scientiarum Fennicae* 13 (1884): 313-496.
- REUTER O. M. 1884. — *Monographia Anthocoridarum orbis terrestris*. Societatis Litterariae Fannicae, Helsingfors, 204 p. <https://doi.org/10.5962/bhl.title.8576>
- REUTER O. M. 1886. — Notes synonymiques. *Revue d'Entomologie de Caen* 5: 120-122. <https://biodiversitylibrary.org/page/25446765>
- REUTER O. M. 1891. — Species novae generis *Acanthia* F., Latr. *Revue d'Entomologie* 10: 21-27.
- RIBAUT H. 1923. — Étude sur le genre *Triphleps* (Heteroptera-Anthocoridae). *Bulletin de la Société d'Histoire naturelle de Toulouse* 51: 522-538
- ROSSI P. 1790. — *Fauna Etrusca, sistens insecta quae in provinciis Florentina et Pisina praesertim collegit*. Vol. 2. Thomae Masi & Socio-rum, Liburni, 348 p. <https://doi.org/10.5962/bhl.title.128484>
- ROSSI P. 1794. — *Mantissa insectorum exhibens species nuper in Etruria collectas a Petro Rossio, adiectis faunae Etruscae illustrationibus, ac emendationibus*. Typographia Polloni, Pisis, 154 p., 8 pls.
- ROUBAL J. 1958 — Description d'une espèce nouvelle de Tingidae de France (Het.). *Bulletin de la Société entomologique de France* 63: 54-56.
- SAHLBERG C. R. 1819 — *Observationes quasdam historiam Notonectidum, imprimis Fennicarum, illustrantes: propositura*. Dissertatio Academica, Typis Frenckelliorum, Aboae, 15 p.
- SAHLBERG R. F. 1848. — *Monographia Geocorisarum Fenniae*. Officina Typographica Frenckelliana, Helsingforsiae, 154 p.
- SCHILLING P. S. 1829. — Hemiptera Heteroptera Silesiae, systematice disposuit (p. 34-96), in *Beiträge zur Entomologie, besonders in Bezug auf die Schlesische Fauna verfasst und herausgegeben von den Mitgliedern der entomologischen Section der Schlesischen Gesellschaft für Vaterländische Kultur*, 1 Heft, Breslau, 226 p. + pls.
- SCHILLING P. S. 1837. — Neue Arten der von Fallén gegründeten Gattung *Phytocoris*. Uebersicht der Arbeiten und Veränderungen der Schlesischen Gesellschaft für Vaterländische Kultur, in Jahr 1836: 83-84.
- SCHMIDT K. 1934. — Beiträge zur deutschen Wanzenfauna. I. (Hem. Het.). *Mitteilungen der Deutschen Entomologischen Gesellschaft* 5: 34.
- SCHOLTZ R. 1847 — Prodrömus zu einer Rhynchoten-Fauna von Schlesien. I. Abhandlungen der Schlesischen. Gesellschaft für Vaterländische Kultur, Naturwissenschaftlich-Medizinische Abteilung, 1846: 104-164.
- SCHRANK F. P. 1776. — Beiträge zur Naturgeschichte. Caspar Fritsch, Leipzig, 137 p. <https://doi.org/10.5962/bhl.title.36918>
- SCHRANK F. P. 1782. — Kritische Revision des österreichischen Insecten-Verzeichnisses. *Neues Magazin für die Liebhaber der Entomologie* 1: 135-168; 263-306.
- SCHRANK F. P. 1785. — Verzeichnis beobachteter Insecten im Fürstenthume Berchtesgaden. *Neues Magazin für die Liebhaber der Entomologie* 2: 313-345.
- SCHUH R. T. & SLATER J. A. 1995. — *True Bugs of the World (Hemiptera: Heteroptera). Classification and Natural History*. Cornell University Press, Ithaca, New York, XII + 336 p.
- SCOPOLI J. A. 1763. — *Entomologia Carniolica exhibens insecta Carniolae indigena et distributa in ordines, genera, species, varietates. Methodo Linneana*. Joannis Thomae Trattner, Vindobonae, XXXVI + 420 p. <https://doi.org/10.5962/bhl.title.119976>
- SCOTT J. 1871. — Neue europäische Hemiptera. *Stettiner Entomologische Zeitung* 31: 98-102.
- SEIDENSTÜCKER G. 1951. — *Trapezonotus desertus* n. sp., eine neue deutsche Lygaeiden-Art (Heteroptera). *Senckenbergiana* 32 (1-4): 79-85.
- SERVADEI A. 1955. — Primi reperti sulla emitterofauna del Terminillo. *Bollettino di Zoologia* 22 (2): 229-242. <https://doi.org/10.1080/11250005509439203>
- SERVADEI A. 1957. — Contributo alla conoscenza degli Emitteri Eterotteri del Gargano. *Memorie di Biogeografia Adriatica*, Padova 4: 1-26.
- SERVADEI A. 1967. — *Fauna d'Italia. IX. Rhyncota, Heteroptera e Homoptera Auchenorrhyncha. Catalogo topografico e sinonimico*. Calderini, Bologna, 852 p.
- SIMON H. 1992. — Vergleichende Untersuchungen zur Wanzenfauna (Heteroptera) von Streuobstwiesen in Nordpfälzer Bergland. *Beiträge zur Landespflege in Rheinland-Pfalz* 15: 189.
- SPINOLA M. 1837. — *Essai sur les genres d'insectes appartenants à l'ordre des Hémiptères, Lin. ou Rhynqotes, Fab. et à la section des Héteroptyères, Dufour*. Yves Gravier, Genova, 383 p.
- STÅL C. 1868-1869. — Hemiptera Fabriciana. Fabricianska Hemipterarter, efter de i Köpenhamn och Kiel Förvarade typexemplaren granskade och beskrifne. *Kongliga Svenska Vetenskaps-Akademiens Handlingar* 7 (2): 1-148. <https://biodiversitylibrary.org/page/33981809>
- STEIN J. P. E. F. 1857. — Die Gattung *Prostemma* Laporte. *Berliner Entomologische Zeitschrift* 1: 81-96. <https://doi.org/10.1002/mmnd.18570010113>
- STICHEL W. 1959 — *Illustrierte Bestimmungstabellen der Wanzen. II. Europa (Hemiptera-Heteroptera Europae). 3.6. Cimicomorpha, Reduviidae, Nabidae*. Berlin-Hermsdorf: 81-206.
- STICHEL W. 1960. — *Illustrierte Bestimmungstabellen der Wanzen. II. Europa (Hemiptera-Heteroptera Europae). 3.12. Liste der Paläarktischen Arten, Cimicomorpha*. Berlin-Hermsdorf: 364-381.
- STOCH F. 2003. — Checklist of the species of the Italian fauna. Online version 2.0. Available at: <http://www.faunaitalia.it/checklist/> [last access: 3 November 2017].
- TAMANINI L. 1951. — Gli *Stictopleurus* italiani (Heteroptera, Corizidae). *Memorie della Società Entomologica Italiana* 30: 77-91.
- TAMANINI L. 1956. — Alcune osservazioni sui *Dicyphus* italiani e loro distribuzione (Heteroptera Miridae). *Memorie della Società Entomologica Italiana* 35: 14-22.
- TAMANINI L. 1971. — Un nuovo miride delle Alpi Marittime italiane (Hemiptera, Heteroptera, Miridae). *Memorie del Museo civico di Storia naturale di Verona* 19: 363-369.
- TAMANINI L. 1976. — Un nuovo *Dimorphocoris* delle Alpi Cozie con ulteriori notizie sul *Dimorphocoris ruffoi* Tam., 1971 (Hemiptera, Heteroptera, Miridae). *Bollettino del Museo Civico di Storia Naturale di Verona* 2 (1975): 323-330.
- TAMANINI L. 1977. — Miridi nuovi o interessanti per la fauna italiana (Hemiptera, Heteroptera, Miridae). *Bollettino della Società Entomologica Italiana* 109: 35-41.
- VIGNA TAGLIANTI A., AUDISIO P. A., BELFIORE C., BIONDI M., BOLOGNA M. A., CARPANETO G. M., DE BIASE A., DE FELICI S., PIATTELLA E., RACHELI T., ZAPPAROLI M. & ZOIA S. 1993. — Riflessioni di gruppo sui corotipi fondamentali della fauna W-Palearctica ed in particolare italiana. *Biogeographia. Lavori della Società italiana di Biogeografia* (n. s.), 16 (1992): 159-179.
- VIGNA TAGLIANTI A., AUDISIO P. A., BIONDI M., BOLOGNA M. A., CARPANETO G. M., DE BIASE A., FATTORINI S., PIATTELLA E.,

- SINDACO R., VENCHI A. & ZAPPAROLI M. 1999. — A proposal for chorotype classification of the near east fauna, in the framework of the Western Palearctic region. *Biogeographia. Lavori della Società italiana di Biogeografia* (n. s.) 20: 31-59.
- VILLEMANT C., DAUGERON B., GARGOMINY O., ISAIA M., DEHARVENG L. & JUDSON M. L. I. 2015. — The Mercantour/Alpi Maritime All Taxa Biodiversity Inventory (ATBI): achievements and prospects, in DAUGERON C., DEHARVENG L., ISAIA M., VILLEMANT C. & JUDSON M. (eds), Mercantour/Alpi Maritime All Taxa Biodiversity Inventory. *Zoosystema* 37 (4): 667-679. <https://doi.org/10.5252/z2015n4a10>
- WAGNER E. 1943. — Einige neue Miriden aus dem Museum National d'Histoire Naturelle in Paris (Hem. Het.). *Mitteilungen der Deutschen Entomologischen Gesellschaft* 12: 8-15.
- WAGNER E. 1951. — Zur Systematik der Gattung *Dicyphus* Fieb. *Societatis Scientiarum Fennica, Commentationes Biologicae* 12: 1-36.
- WAGNER E. 1953. — Eine neue Miridenart aus Südfrankreich. *Revue française d'Entomologie* 20 (1): 84-86.
- WAGNER E. 1954. — Eine Beitrag zur Systematik der Gattung *Phytocoris* Fall. (Hem. Het. Miridae). *Nachrichten des Naturwissenschaftlichen Museums der Stadt Aschaffenburg* 42: 43.
- WAGNER E. 1970. — Die Miridae Hahn, 1831, des Mittelmeerraumes und Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 1. *Entomologische Abhandlungen Staatliches Museum für Tierkunde Dresden* 37 (Suppl.): 1-272.
- WOLFF J. F. 1804. — *Icones Cimicum descriptionibus illustratae*. J. J. Palm, Erlangen, Fasc. 4: 127-166, pls 13-16. <https://doi.org/10.5962/bhl.title.15807>
- WOLFF J. F. 1811. — *Abbildungen der Wanzen mit Beschreibungen*. Vol. 5. J. J. Palm, Erlangen, 167-208, pls 17-20. <https://doi.org/10.5962/bhl.title.12481>
- ZETTERSTEDT J. W. 1838. — *Insecta Lapponica descripta*. L. Voss, Lipsiae, 1: 1-868. <https://biodiversitylibrary.org/page/8316770>

*Submitted on 10 January 2018;
accepted on 28 September 2018;
published on 9 July 2019.*