


Lectotypification of two Turkish endemic taxa of *Centaurea* L. (Asteraceae, Cardueae, Centaureinae)

Kazem NEGARESH



TYPE

Phaeopappus paphlagonica Bornm.
Beih. Bot. Centralbl. 59B: 304. 1939
det. P.H. DAVIS, Flora of Turkey 5: 524. 1975
V. BOURGIGNON - 02-2003

DIRECTEUR DE LA PUBLICATION : Bruno David
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF* : Thierry Deroin

RÉDACTEURS / *EDITORS* : Porter P. Lowry II ; Zachary S. Rogers

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS* : Emmanuel Côté (adanson@mnhn.fr) ; Anne Mabilie

MISE EN PAGE / *PAGE LAYOUT* : Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD* :

P. Baas (Nationaal Herbarium Nederland, Wageningen)
F. Blasco (CNRS, Toulouse)
M. W. Callmender (Conservatoire et Jardin botaniques de la Ville de Genève)
J. A. Doyle (University of California, Davis)
P. K. Endress (Institute of Systematic Botany, Zürich)
P. Feldmann (Cirad, Montpellier)
L. Gautier (Conservatoire et Jardins botaniques de la Ville de Genève)
F. Ghahremaninejad (Kharazmi University, Téhéran)
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)
K. Kubitzki (Institut für Allgemeine Botanik, Hamburg)
J.-Y. Lesouef (Conservatoire botanique de Brest)
P. Morat (Muséum national d'Histoire naturelle, Paris)
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)
S. E. Rakotoarisoa (Millenium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)
É. A. Rakotobe (Centre d'Applications des Recherches pharmaceutiques, Antananarivo)
P. H. Raven (Missouri Botanical Garden, St. Louis)
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)
J. G. West (Australian National Herbarium, Canberra)
J. R. Wood (Oxford)

COUVERTURE / *COVER* :

Centaurea paphlagonica (Bornm.) Wagenitz (P00703840).

Adansonia est indexé dans / *Adansonia is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Adansonia est distribué en version électronique par / *Adansonia is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Adansonia est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Adansonia is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:
Geodiversitas, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)
Tél. : 33 (0)1 40 79 48 05 / Fax : 33 (0)1 40 79 38 40
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2018
ISSN (imprimé / *print*) : 1280-8571/ ISSN (électronique / *electronic*) : 1639-4798

Lectotypification of two Turkish endemic taxa of *Centaurea* L. (Asteraceae, Cardueae, Centaureinae)

Kazem NEGARESH

Department of Horticulture, Faculty of Agriculture,
Khuzestan Agricultural Sciences and Natural Resources University,
Mollasani, Ahvaz (Iran)
negaresh@ramin.ac.ir

Submitted on 13 February 2018 | accepted on 15 April 2018 | published on 27 August 2018

KEY WORDS

Asteraceae,
endemic,
Turkey,
lectotypifications.

MOTS CLÉS

Asteraceae,
endémiques,
Turquie,
lectotypifications.

Negaresh K. 2018. — Lectotypification of two Turkish endemic taxa of *Centaurea* L. (Asteraceae, Cardueae, Centaureinae). *Adansonia*, sér. 3, 40 (10): 135-139. <https://doi.org/10.5252/adansonia2018v40a10>. <http://adansonia.com/40/10>

ABSTRACT

Two Turkish endemic taxa of *Centaurea* L. (Asteraceae) are lectotypified, namely *Centaurea leptophylla* (K.Koch) Tchihat. and *C. paphlagonica* (Bornm.) Wagenitz.

RÉSUMÉ

Lectotypification de deux taxons endémiques turcs de *Centaurea* L. (Asteraceae, Cardueae, Centaureinae). Deux taxons endémiques turcs de *Centaurea* L. (Asteraceae) sont lectotypifiés, à savoir *Centaurea leptophylla* (K.Koch) Tchihat. et *C. paphlagonica* (Bornm.) Wagenitz.

INTRODUCTION

The genus *Centaurea* s.l. consists of about c. 600 species in 40 sections in world (Bremer 1994; Hellwig 2004; Negaresh & Rahiminejad 2014; Oreizi *et al.* 2017). This genus is considered to be taxonomically difficult due to the wide morphological and karyological variability and also nomenclatural confusions (Tchihat 1860; Bremer 1994; Wagenitz & Hellwig 1996, 2000; Hellwig 2004; Garcia-Jacas *et al.* 2006; Susanna & Garcia-Jacas 2007, 2009; Uysal *et al.* 2009; Negaresh & Rahiminejad 2014, 2015a).

Continuing my nomenclatural study of the names in *Centaurea* (Ranjbar & Negaresh 2013a, b, c, 2014a, b, c, d, e; Ranjbar *et al.* 2013; Negaresh & Rahiminejad 2014, 2015a, b; Negaresh *et al.* 2014, 2015a, b; Oreizi *et al.* 2017), I realized the necessity to lectotypify two species endemic to Turkey, i.e., *C. leptophylla* (K.Koch) Tchihat. and *C. paphlagonica* (Bornm.) Wagenitz, because no specific herbarium sheet was cited as holotype in their protologues and the taxa were also not lectotypified earlier (Wagenitz 1963, 1975). To clarify the application of these names, they are thus lectotypified here with specimens housed in the Muséum national d'Histoire naturelle, Herbier national de Paris (P).

LECTOTYPIFICATIONS

Centaurea leptophylla (K.Koch) Tchihat.
(Fig. 1)

In Asie Mineure. Troisième Partie. Botanique II: 309 (1860). — *Psephellus leptophyllus* (K.Koch) Boiss., *Flora Orientalis* 3: 611 (1875). — Basionym: *Centaureum leptophyllum* K.Koch, *Linnaea* 24: 419 (1851).

LECTOTYPE (designated here). — **Turkey**. [A8 Çoruh] im Gaue Pertakrek [Peterek] im Hochgebirge, c. 6000 ft (1829 m) hoch, auf Porphy, 1843, *K. H. E. Koch s.n.* (lecto-, P[P00646491]!; isolecto-, GOET[GOET001277]!).

REMARKS

Koch (1851) described *Centaureum leptophyllum* (*Centaurea leptophylla*) based on collections of him from Pertakrek, but did not designate an holotype nor did mention the name of the herbarium where the specimens were housed. Two specimens of *C. leptophylla* collected by Koch from Pertakrek were found. Based on art. 40, note 1 (McNeill *et al.* 2012) both must be treated as syntypes. The best one and better preserved sheet, P00646491, is designated here as the lectotype, because it is in better condition than another specimen and shows all relevant characters with the protologue.

Centaurea paphlagonica (Bornm.) Wagenitz
(Fig. 2)

In Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 82 (2): 206 (1963). — *Phaeopappus macrocephalus* Freyn & Sint. in Freyn, *Österreichische botanische Zeitschrift* 44: 219

(1894), nom. illeg. — *Plumosipappus macrocephalus* (Freyn & Sint.) Czerep., *Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Imeni V.L. Komarova Akademii Nauk SSSR* 20: 458 (1960). — Basionym: *Phaeopappus paphlagonicus* Bornm., *Beihefte zum Botanischen Centralblatt* 59B: 304 (1939).

LECTOTYPE (designated here). — **Turkey**. [A5 Kastamonu] Paphlagonia, Wilajet Kastamonu ad Tossia: Daltshadagh, in pratis subalpinis [inter Kalköi et Daltshadagh], 15.VII.1892, *P. E. E. Sintenis* 4626 (lecto-, P[P00703840]!; isolecto-, B[B100088272]!, BM[BM000883842]!, BR[BR0000005263776]!, FR[FR0030804]!, G[G00016449] (three sheets comprising one specimen), G00076555, G00076556 (two sheets comprising one specimen)!, GOET[GOET001299]!, GZU[GZU000271996]!, JE[JE00015892]!, L[L0001901]!, LD[LD1019555, LD1019619, LD1019683, LD1019747]!, M[M0031135, M0242176]!, P[P00703838, P00703839]!, S[S10-9572]!, W[W0005894]!).

REMARKS

Freyn & Sintenis in Freyn (1894) described *Phaeopappus macrocephalus* (synonym of *Centaurea paphlagonica*, not *C. macrocephala* Muss.Puschk. ex Willd., 1803) based on collections of *Sintenis* 4626 from Paphlagonia, but did not mention any specific sheet as the holotype nor the name of the herbarium where the specimens were housed. Twenty two herbarium specimens of collection number 4626 of *Sintenis* from Paphlagonia were found. Based on art. 40, note 1 (McNeill *et al.* 2012) these must all be treated as syntypes. Of these, the sheet P00703840 is better preserved and more complete than the other specimens and best represents the species description given in the protologue; hence it is designated here as the lectotype.

Acknowledgements

I am indebted to the personnel of the following herbaria: B, BM, BR, E, FR, G, GOET, GZU, JE, K, L, LD, M, P, S, and W, for their contribution during the revision of material and for providing the images of types. I would also like to thank to Khuzestan Agricultural Sciences and Natural Resources University for financial support. At last, the author thanks Zafer Kaya, Ernst Vitek and Thierry Deroin for having read a previous version of the manuscript.

REFERENCES

- BREMER K. 1994. — *Asteraceae: Cladistics and Classification*. Timber Press, Portland, Oregon, 752 p.
FREYN J. F. 1894. — *Plantae novae Orientales*, III. *Österreichische Botanische Zeitschrift* 44: 217-220.
GARCIA-JACAS N., UYSAL T., ROMASHCHENKO K., SUÁREZ-SANTIAGO V. N., ERTUĞRUL K. & SUSANNA A. 2006. — *Centaurea* revisited: a molecular survey of the *Jacea* group. *Annals of Botany* 98: 741-753. <https://doi.org/10.1093/aob/mcl157>
HELLWIG F. H. 2004. — *Centaureinae* (Asteraceae) in the Mediterranean – history of ecogeographical radiation. *Plant Systematics and Evolution* 246: 137-162. <https://doi.org/10.1007/s00606-004-0150-2>
KOCH K. H. E. 1851. — Beiträge zu einer Flora des Orientes. *Linnaea* 24: 305-480.
MCNEILL J., BARRIE F. R., BUCK W. R., DEMOULIN V., GREUTER W., HAWKSWORTH D. L., HERENDEN P. S., KNAPP S., MARHOLD K.,



FIG. 1. — Lectotype of *Centaurea leptophylla* (K. Koch) Tchiha., Koch s.n. (P00646491).

- PRADO J., PRUD'HOMME VAN REINE W. F., SMITH G. F., WIERSEMA J. H. & TURLAND N. J. (eds) 2012. — International Code of Nomenclature for algae, fungi and plants (Melbourne Code): Adopted by the Eighteenth International Botanical Congress, Melbourne, Australia. *Regnum Vegetabile* 154: 1-274.
- NEGARESH K. & RAHIMINEJAD M. R. 2014. — A contribution to the taxonomy of *Centaurea* sect. *Cynaroides* (Asteraceae, Cardueae–Centaureinae) in Iran. *Phytotaxa* 158: 229-244. <https://doi.org/10.11646/phytotaxa.158.3.3>
- NEGARESH K., RAHIMINEJAD M. R. & RANJBAR M. 2014. — A taxonomic study on *Centaurea* sect. *Rhizocalathium* and *C.* sect. *Pseudoseridia* (Asteraceae, Cardueae–Centaureinae) in Iran. *Phytotaxa* 174 (1): 43-60. <https://doi.org/10.11646/phytotaxa.174.1.3>
- NEGARESH K. & RAHIMINEJAD M. R. 2015a. — A taxonomic revision of *Centaurea* sect. *Microlophus* (Asteraceae, Cardueae–Centaureinae) and three new records for the flora of Iran. *Nordic Journal of Botany* 33: 335-353. <https://doi.org/10.1111/njb.00587>
- NEGARESH K. & RAHIMINEJAD M. R. 2015b. — The identity and revised circumscription of *Centaurea iranshabrii* (Asteraceae, Cardueae–Centaureinae) in Iran. *Feddes Repertorium* 126 (3-4): 55-66. <https://doi.org/10.1002/fedr.201500007>
- NEGARESH K., KHOSHROO S. M. R., KARAMIAN R. & JOHARCHI M. R. 2015a. — A revision of *Rhaponticoides* (Asteraceae, Cardueae–Centaureinae) from Iran. *Phytotaxa* 213 (2): 87-101. <https://doi.org/10.11646/phytotaxa.213.2.2>
- NEGARESH K., KAYA Z. & RAHIMINEJAD M. R. 2015b. — *Centaurea sennikoviana* (Asteraceae, Cardueae), a new species from central Anatolia, Turkey. *Annales Botanici Fennici* 52: 321-327. <https://doi.org/10.5735/085.052.0507>
- OREIZI E., NEGARESH K. & RAHIMINEJAD M. R. 2017. — A new name in *Centaurea* (Asteraceae, Cardueae) from Turkey. *Candollea* 72: 319-322. <https://doi.org/10.15553/c2017v722a7>
- RANJBAR M. & NEGARESH K. 2013a. — A revision of *Centaurea* sect. *Phaeopappus* (Asteraceae, Cardueae). *Phytotaxa* 123 (1): 1-40. <https://doi.org/10.11646/phytotaxa.123.1.1>
- RANJBAR M. & NEGARESH K. 2013b. — Taxonomic notes on *Centaurea* sect. *Acrocentron* (Asteraceae) in Iran, with the description of a new species. *Annales Botanici Fennici* 50: 239-248. <https://doi.org/10.5735/086.050.0406>
- RANJBAR M. & NEGARESH K. 2013c. — A contribution to *Centaurea* sect. *Acrocentron* (Asteraceae) from Iran. *Phytotaxa* 149 (1): 50-60. <https://doi.org/10.11646/phytotaxa.149.1.6>
- RANJBAR M., NEGARESH K. & JOHARCHI M. R. 2013. — Novelty in *Centaurea* sect. *Acrocentron* (Asteraceae, Cardueae) from Iran. *Feddes Repertorium* 124: 116-121. <https://doi.org/10.1002/fedr.201400030>
- RANJBAR M. & NEGARESH K. 2014a. — On the identity of *Centaurea paradoxa* (Asteraceae, Cardueae, Centaureinae) from Iran. *Willdenowia* 44: 13-20. <https://doi.org/10.3372/wi.44.44102>
- RANJBAR M. & NEGARESH K. 2014b. — A revision of *Centaurea* sect. *Centaurea* (Asteraceae) from Iran. *Turkish Journal of Botany* 38: 969-987. <https://doi.org/10.3906/bot-1304-26>
- RANJBAR M. & NEGARESH K. 2014c. — The genus *Centaurea* (Asteraceae, Cardueae) in Iran: two new species and a lectotypification. *Edinburgh Journal of Botany* 71 (1): 51-56. <https://doi.org/10.1017/S0960428613000322>
- RANJBAR M. & NEGARESH K. 2014d. — Taxonomic notes and two new species in *Centaurea* sect. *Cynaroides* (Compositae, Cardueae) from Iran. *Novon* 23 (3): 321-327. <https://doi.org/10.3417/2012056>
- RANJBAR M. & NEGARESH K. 2014e. — New combinations of *Rhaponticoides* (Asteraceae, Cardueae) from Afghanistan. *Global Journal of Science Frontier Research* 14 (3): 13-17.
- SUSANNA A. & GARCIA-JACAS N. 2007. — Tribe Cardueae, in KADEREIT J. W. & JEFFREY C. (eds), *The Families and Genera of Vascular Plants*. Vol. 8. Springer Verlag, Berlin, Heidelberg & New York: 123-146.
- SUSANNA A. & GARCIA-JACAS N. 2009. — Cardueae (Carduoideae), in FUNK V. A., SUSANNA A., STUESSY T. F. & BAYER R. J. (eds), *Systematics, Evolution, and Biogeography of Compositae*. Vienna, International Association for Plant Taxonomy: 293-313.
- TCHIHATCHEFF P. DE 1860. — *Asie Mineure. Description physique, statistique et archéologique de cette contrée*. Troisième partie. Botanique II. Gide, Libraire-Éditeur, Paris, 676 p.
- UYSAL T., ERTUĞRUL K., SUSANNA A. & GARCIA-JACAS N. 2009. — New chromosome counts in the genus *Centaurea* (Asteraceae) from Turkey. *Botanical Journal of the Linnean Society* 159: 280-286. <https://doi.org/10.1111/j.1095-8339.2008.00939.x>
- WAGENITZ G. 1963. — Die Eingliederung der 'Phaeopappus'-Arten in das System von *Centaurea*. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 82 (2): 137-215.
- WAGENITZ G. 1975. — *Centaurea*, in DAVIS P. H. (ed.), *Flora of Turkey and the East Aegean Islands*. Vol. 5. Edinburgh, UK: Edinburgh University Press: 465-585.
- WAGENITZ G. & HELLWIG F. H. 1996. — Evolution of characters and phylogeny of the Centaureinae, in HIND D. J. N. & BEENTJE H. J. (eds), *Compositae: Systematics*. Proceedings of the International Compositae Conference, Kew, 1994. Royal Botanic Gardens, Kew: 491-510.
- WAGENITZ G. & HELLWIG F. H. 2000. — *Psephellus* Cass. (Compositae, Cardueae) revisited with a broadened concept. *Willdenowia* 30: 29-44. <http://www.jstor.org/stable/3997072>

Submitted on 13 February 2018;
accepted on 15 April 2018;
published on 27 August 2018.



FIG. 2. — Lectotype of *Centaurea paphlagonica* (Bornm.) Wagenitz, Sintenis 4626 (P00703840).