

Lectotypification of *Hedychium thyrsiforme*
Buch.-Ham. ex Sm. (Zingiberaceae)

Ajith ASHOKAN & Vinita GOWDA



DIRECTEUR DE LA PUBLICATION / *PUBLICATION DIRECTOR*: Gilles Bloch
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Thierry Deroin

RÉDACTEURS / *EDITORS*: Porter P. Lowry II; Zachary S. Rogers; Mathieu Gardère

ÉDITEUR TECHNIQUE (SUIVI ÉDITORIAL) / *DESK EDITOR (EDITORIAL PROCESS)*: Emmanuel Côté (adanson@mnhn.fr)

ÉDITEUR TECHNIQUE (PRODUCTION) / *DESK EDITOR (PRODUCTION)*: Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

F. Blasco (CNRS, Toulouse)
M. W. Callmender (Conservatoire et Jardin botaniques de Genève)
J. A. Doyle (University of California, Davis)
P. K. Endress (Institute of Systematic Botany, Zürich)
P. Feldmann (Cirad, Montpellier)
L. Gautier (Conservatoire et Jardin botaniques de Genève)
F. Ghahremaninejad (Kharazmi University, Téhéran)
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)
A. A. Khapugin (Tyumen State University, Russia)
J.-Y. Lesouef (Conservatoire botanique de Brest)
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)
S. E. Rakotoarisoa (Millenium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)
P. H. Raven (Missouri Botanical Garden, St. Louis)
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)
J. G. West (Australian National Herbarium, Canberra)
J. R. Wood (Oxford)

COUVERTURE / *COVER*:

Réalisée à partir des Figures de l'article/*Made from the Figures of the article.*

Adansonia est indexé dans / *Adansonia is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Adansonia est distribué en version électronique par / *Adansonia is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Adansonia est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Adansonia is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*: *Geodiversitas*, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Cryptogamie* sous-sections *Algologie*, *Bryologie*, *Mycologie*, *Comptes Rendus Palevol*

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)
Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

Les articles publiés dans *Adansonia* sont distribués sous [licence CC-BY 4.0](#)/Articles published in *Adansonia* are distributed under a [CC-BY 4.0 license](#).
ISSN (imprimé / *print*): 1280-8571/ ISSN (électronique / *electronic*): 1639-4798

Lectotypification of *Hedychium thyrsoforme* Buch.-Ham. ex Sm. (Zingiberaceae)

Ajith ASHOKAN

Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences,
Indian Institute of Science Education and Research (IISER) Bhopal, Madhya Pradesh 462066 (India)
and Department of Biology, University of Naples Federico II,
Complesso Universitario di Monte S. Angelo, Via Cintia, 80126, Napoli (Italy)
ajithashoksridhar@gmail.com (corresponding author)

Vinita GOWDA

Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences,
Indian Institute of Science Education and Research (IISER) Bhopal, Madhya Pradesh 462066 (India)
gowdav@iiserb.ac.in

Submitted on 23 March 2025 | accepted on 25 May 2025 | published on 19 January 2026

Ashokan A. & Gowda V. 2026. — Lectotypification of *Hedychium thyrsoforme* Buch.-Ham. ex Sm. (Zingiberaceae). *Adansonia*, sér. 3, 48 (1): 1-5. <https://doi.org/10.5252/adansonia2026v48a1>. <http://adansonia.com/48/1>

ABSTRACT

The tropical and subtropical Asian genus *Hedychium* J.Koenig (Zingiberaceae) comprises over 90 recognized species, with more than twice as many published names. It is renowned for its enduring taxonomic uncertainties. Here, we assign the lectotype for *H. thyrsoforme* Buch.-Ham. ex Sm. and provide comprehensive morphological description for this species.

RÉSUMÉ

Lectotypification de Hedychium thyrsoforme Buch.-Ham. ex Sm. (Zingiberaceae).

Le genre tropical et subtropical asiatique *Hedychium* J.Koenig (Zingiberaceae) comprend plus de 90 espèces reconnues, avec plus de deux fois autant de noms publiés. Il est réputé pour ses incertitudes taxonomiques persistantes. Nous désignons ici un lectotype pour *H. thyrsoforme* Buch.-Ham. ex Sm. et fournissons une description morphologique complète de cette espèce.

KEY WORDS
Zingiberaceae,
India,
lectotypification.

MOTS CLÉS
Zingiberaceae,
Inde,
lectotypification.

INTRODUCTION

Johann Gerhard Koenig established the genus *Hedychium* in 1783 (Koenig 1783), based on an illustration of *Gandasilum* by Georg Eberhard Rumphius (Rumphius 1747; Sanoj 2011; Ashokan & Gowda 2017; Ashokan & Gowda 2019). Following the initial description of *H. coronarium* by J. G. Koenig, James Edward Smith expanded the genus by adding four more species (*H. coccineum*, *H. ellipticum*, *H. spicatum* and *H. thyriforme*; Smith 1811). During the nineteenth and twentieth centuries, numerous *Hedychium* species were described (Smith 1811; Roscoe 1824-1825; Wallich 1853; Horaninow 1862; Baker 1892; Schumann 1904), often with insufficient type information, leading to persistent taxonomic challenges within the genus (Sanoj 2011; Ashokan & Gowda 2023). In this study, we aim to lectotypify *H. thyriforme* that was originally described by J. E. Smith (Smith 1811).

MATERIAL AND METHODS

We carried out field collection trips between 2015 and 2019 across Northeast (NE) India. Metadata documentation included morphological, phenological, and ecological characteristics. Morphological measurements were taken using both a ruler and digital calipers. Collected specimens comprised herbarium vouchers and spirit samples, and leaf tissues preserved in silica for molecular studies (Ashokan *et al.* 2022).

Protologues of all *Hedychium* names were examined in monographs and taxonomic revisions (Koenig 1783; Smith 1811; Roscoe 1824-1825; Wallich 1853; Horaninow 1862; Baker 1892; Schumann 1904; Naik & Panigrahi 1961; Rao & Verma 1969; Srivastava 1984; Smith 1994; Sanoj 2011). Herbarium specimens, including type collections, were studied at the listed herbaria. Additionally, digital resources were consulted, including Global Plants (<https://plants.jstor.org/>), the Kew Herbarium Catalogue (<http://apps.kew.org/herbcat/>), Muséum national d'Histoire naturelle (<https://science.mnhn.fr/>), the Smithsonian Institution (<https://www.si.edu/>), The Linnean Collections (<http://linnean-online.org/>), and the Zingiberaceae Resource Centre (<http://padme.rbge.org.uk/ZRC/>).

ABBREVIATIONS

Collection acronyms

ARUN	Herbarium, Botanical Survey of India, Arunachal Pradesh Regional Centre;
ASSAM	Herbarium, Botanical Survey of India, Eastern Regional Centre;
BHPL	Herbarium, Indian Institute of Science Education and Research Bhopal;
BM	Herbarium, The Natural History Museum;
BO	Herbarium Bogoriense, The National Research and Innovation Agency;
BSA	Herbarium, Botanical Survey of India, Central Regional Centre;
BSHC	Herbarium, Botanical Survey of India, Sikkim Himalayan Circle;
BSI	Herbarium, Botanical Survey of India, Western Circle;

CAL	Herbarium, Botanical Survey of India, Howrah;
E	Herbarium, Royal Botanic Garden Edinburgh;
K	Herbarium, Royal Botanic Gardens, Kew;
LINN	Herbarium, Linnean Society of London;
LIV	Herbarium, World Museum, National Museums Liverpool;
MH	Herbarium, Botanical Survey of India, Southern Circle;
QBG	Herbarium, Queen Sirikit Botanic Garden, Chiang Mai;
SING	Herbarium, Singapore Botanic Gardens;
TBGT	Herbarium, Tropical Botanic Garden and Research Institute, Kerala.

TAXONOMIC TREATMENT

Family ZINGIBERACEAE Martinov, *nom. cons.*
Genus *Hedychium* J.Koenig.

Hedychium thyriforme Buch.-Ham. ex Sm.
(Fig. 1)

Hedychium no. 5, Cycl. 17 (Smith in Rees 1811).

TYPE MATERIAL. — Nepal • Buchanan-Hamilton *s.n.*; lectotype: LINN [8.34!], here designated (Fig. 1A).

PHENOLOGY AND ECOLOGY. — Flowering from August to September. Fruiting September to October. Flowers are not remarkably fragrant.

GLOBAL DISTRIBUTION. — India (Arunachal Pradesh, Sikkim, Manipur, Mizoram, Nagaland), Bhutan, Myanmar, Nepal; this taxon is distributed at an elevation of more than 1400 m.

IUCN STATUS. — We categorize it as data deficient (DD) following the IUCN guidelines (IUCN Standards and Petitions Subcommittee 2024).

ETYMOLOGY. — The specific epithet, “*thyriforme*”, refers to the inflorescence: panicle-like or thyrse (thyrsi-); resembling or shaped (-forme).

VERNACULAR NAME. — Pincushion Ginger Lily.

MORPHOLOGICAL AFFINITIES. — *Hedychium thyriforme* closely resembles *H. larsenii* in its pyramidal inflorescence, white flowers, and the size of the labellum, lateral staminodes, and filament. However, the two species differ in the lengths of the bracteole, calyx, floral tube, anther, and lamina (Table 1).

PHYLOGENETIC PLACEMENT. — *Hedychium thyriforme* is consistently recovered in the clade IV in phylogenetic reconstructions of the genus (Wood *et al.* 2000; Ashokan *et al.* 2022).

DESCRIPTION

Terrestrial, rhizomatous herb up to 1.2 m tall; rhizome branching parallel to leaf distichy, pale brown externally, creamy white internally; leafy shoot slanting with an erect inflorescence; basal sheath greenish. Leaves sessile; sheath pale green; ligule 3.2-4.8 cm wide, ovate, densely sericeous, pale green; lamina 31-44 × 8-14 cm, elliptic-lanceolate, glabrous and dark green (adaxial) and densely sericeous and pale green (abaxial); acumen 2.5-3.0 cm long, ends in a small caducous appendage; base cuneate. Inflorescence 8-13 × 6-10 cm, dense, pyramidal; rachis hidden; peduncle pubescent,




FIG. 1. — *Hedychium thyrsiforme* Buch.-Ham. ex Sm.: **A**, lectotype of *H. thyrsiforme* available at LINN; **B**, inflorescence; **C**, floral parts: **a**, bract; **b**, bracteole; **c**, old flower; **d**, young flower buds; **e**, calyx; **f**, floral tube; **g**, dorsal corolla lobe ($\times 1$); **h**, lateral corolla lobes ($\times 2$); **i**, lateral staminodes ($\times 2$); **j**, central labellum; **k**, filament; **l**, anther; **m**, ovary; **n**, epigynous nectaries ($\times 2$); **o**, style; **p**, stigma; **D**, flower. Scale bar: C, D, 2.0 cm. Photo credit: A, © Linnean Museum; B, C, D, Ajith Ashokan.

green; bracts 2.5(3.2 × 1.1-1.2 cm, elliptic, puberulent, green, folded, involute, 2-6 flowered; bracteoles 2.0-2.5 cm long, tubular, pubescent, pale green. Flowers 11.5-13.5 cm long, ascending, white (when fresh), dull white the second day, many flowers (10-15) open simultaneously, faintly fragrant; calyx 2.4-3.0 cm long, tubular, translucent, 3-toothed at tip, slightly hairy at tip, pale green; floral tube 3.4-4.2 × 0.2-0.3 cm, glabrous, erect, white; corolla lobes linear, curled, orientated downwards, glabrous, pale green; dorsal lobe 5.2-5.9 cm long, linear, curled (more curled as the flower ages), glabrous, beaked at tip, embracing the filament and anther in the bud stage, pale green; lateral lobes 5.1-5.7 cm long, linear, curled (more curled as the flower ages), non-beaked at tip, glabrous, white; lateral staminodes 3.0-3.6 cm long, widest part 0.4 cm, narrowly oblanceolate, one edge linear, obtuse, unguiculate, upper halves reflexed backwards, positioned acute (< 80°) with respect to labellum, white, pale yellow towards base; central labellum 2.2-3.0 × 2.0-2.5 cm, orbicular, glabrous, white with light yellow blotch, bilobed, canaliculate, distinctly clawed; lobes acute, upper halves slightly reflexed backwards; sinus 0.9-1.3 cm deep; claw 1.0-1.3 × 0.2-0.3 cm; filament 5.5-6.5 cm long, slightly arching, deeply grooved on one side, white; anther 0.9-1.1 cm long, linear, basifixed, cream white; thecae 2, split longitudinally; connective 0.1-0.2 cm long, cream white; stigma 0.1 cm wide, cup-shaped, hairy at tip, exerted from the anther by at least 0.1 cm, greenish yellow; style 11.4-13.4 cm long, filiform, white; epigynous nectaries 0.3-0.6 cm long, glabrous, pale yellow; ovary 0.3-0.4 × 0.2-0.3 cm, barrel-shaped, pubescent, yellowish green; placentation axile; ovules spherical, white. Fruit not seen.

TIPIFICATION REMARKS

Francis Buchanan-Hamilton (1762-1829), a Scottish surgeon with the British East India Company, participated in a diplomatic mission to Nepal from 1802 to 1803, during which he collected numerous plants, including *Hedychium* (Watson & Noltie, 2016). On his return to Britain, he gave his collections, notes and drawings to J. E. Smith, the founding President of the Linnean Society of London, and the person to whom the Court of Directors of the British East India Company turned when they needed botanical advice (Watson & Noltie, 2016). A partial set of Buchanan-Hamilton's botanical collections also went to Aylmer Bourke Lambert; this material is now at the Natural History Museum, London while the material given to J.E. Smith is still held at the Linnean Society of London. Using Buchanan-Hamilton's collections and notes, Smith described four species of *Hedychium* including *H. thyrsiforme*. The entry in Rees's Cyclopaedia for *H. thyrsiforme* should be quoted, 'Gathered by Dr Buchanan, in August 1802, at Narainhetty, in Upper Nepal, with the last [*H. spicatum*], from which by name the natives do not distinguish it'. During our visit to the Linnean Society of London Herbarium (LINN), we discovered a specimen of *H. thyrsiforme* in the Smith Herbarium at LINN. This specimen carries the date "August 1802" and the location "Narainhetty, Upper Nepal" as recorded in Rees's Cyclopaedia (Smith, 1811). Based on this finding, we designate it as the lectotype.

Acknowledgements

We thank the respective Indian state forest departments for field research permits- Arunachal Pradesh, Kerala, Manipur, Meghalaya, Mizoram, and Nagaland. We also thank the following herbaria and associated botanic gardens- ARUN, ASSAM, BHPL, BM, BO, BSA, BSHC, BSI, CAL, E, HLA, K, LINN, LIV, MH, QBG, SING, TBGT, US, and W for letting us access all *Hedychium* specimens in their care. AA acknowledges the Council of Scientific and Industrial Research for the Senior Research Fellowship, Heliconia Society International for the Field Research Grant, Science and Engineering Research Board for International Travel Scheme Grant, and Royal Botanic Garden Edinburgh for the Sibbald Trust Grant. VG acknowledges the Ministry of Human Resource Development, Science and Engineering Research Board, and National Geographic Society for research funds, and IISER Bhopal for the infrastructure. Further, we thank Biodiversity Heritage Library (BHL), International Plant Names Index (IPNI), Padme (Zingiberaceae Resource Centre, RBGE), RBGE Herbarium Catalogue, Herbarium Catalogue of Royal Botanic Gardens, Kew for generously hosting information on historical collections of *Hedychium* including protologues and type specimens. At last, the authors thank Thierry Deroin and an anonymous referee for their comments on a previous version of the article.

REFERENCES

- ASHOKAN A. & GOWDA V. 2017. — Diversity of the genus *Hedychium* J. Koenig (Zingiberaceae) in the Northeast of India-I. *HSI Bulletin* 23: 1-11 [retrieved from https://www.heliconia.org/_files/ugd/ef1aa2_b31c53686b7b43c4b92d717ad9f4d6f2.pdf].
- ASHOKAN A. & GOWDA V. 2019. — *Hedychium ziroense* (Zingiberaceae), a new species of ginger lily from Northeast India. *PhytoKeys* 117: 73-84. <https://doi.org/10.3897/phytokeys.117.24951>
- ASHOKAN A. & GOWDA V. 2023. — Etymological histories of all names in *Hedychium* (Zingiberaceae), Part I. *HSI Bulletin* 29: 9-14.
- ASHOKAN A., XAVIER A., SUKSATHAN P., ARDIYANI M., LEONG-ŠKORNIČKOVÁ J., NEWMAN M., KRESS W. J. & GOWDA V. 2022. — Himalayan orogeny and monsoon intensification explain species diversification in an endemic ginger (*Hedychium*: Zingiberaceae) from the Indo-Malayan Realm. *Molecular Phylogenetics and Evolution* 170: 107440. <https://doi.org/10.1016/j.ympev.2022.107440>
- ASHOKAN A., NEWMAN M. F., SANOJ E., SABU M. & GOWDA V. 2025. — Synonymizing two species names, designating lectotypes, and cataloging intermediates in two hybrid zones of *Hedychium*. *Annals of the Missouri Botanical Garden* 110: 128-137. <https://doi.org/10.3417/2025940>
- BAKER J. G. 1892. — *Hedychium* Koenig, in HOOKER J. D. (ed.), *Flora of British India*. Vol. 6. Reeve & Co., London: 225-233. <https://www.biodiversitylibrary.org/page/37634311>
- HORANINOW P. F. 1862. — *Prodromus Monographiae Scitaminarum*. St. Petersburg. <https://www.biodiversitylibrary.org/page/30822561>
- IUCN STANDARDS AND PETITIONS SUBCOMMITTEE 2024. — *Guidelines for Using the IUCN Red List Categories and Criteria*. Version 16. <https://www.iucnredlist.org/resources/redlistguidelines>.
- KOENIG J. G. 1783. — Descriptions Monandrarum pro annis, in RETZIUS A. J. (ed.), *Observationes Botanicae*. Vol. 3. Lipsiae: 45-76. <https://www.biodiversitylibrary.org/page/12127387>
- NAIK V. N. & PANIGRAHI G. 1961. — Genus *Hedychium* in Eastern India. *Bulletin of Botanical Survey of India* 3: 67-73. <https://doi.org/10.20324/nelumbo/v3/1961/76577>

- RAO A. S. & VERMA D. M. 1969. — Notes on *Hedychium* Koenig, including four new species from Khasi & Jaintia Hills, Assam. *Bulletin of the Botanical Survey of India* 11: 120-128.
- ROSCOE W. 1824-1825. — *Monandrian Plants of the Order Scitamineae*. Liverpool. <https://www.biodiversitylibrary.org/page/12127386>
- RUMPHIUS G. E. 1747. — *Herbarium amboinense*. Vol. 5. <https://bibdigital.rjb.csic.es/idurl/1/14104>
- SANOJ E. 2011. — Taxonomic revision of the genus *Hedychium* J. Koenig (Zingiberaceae) in India. University of Calicut, Kerala, 300 p. <http://hdl.handle.net/10603/80898>
- SCHUMANN K. 1904. — Zingiberaceae, in ENGLER A. (ed.), *Das Pflanzenreich* IV. 46 (Heft 20). Engelmann, Leipzig: 1-458. <https://www.biodiversitylibrary.org/page/16815036>
- SMITH J. E. 1811. — *Hedychium*, in REES A. (ed.), *Cyclopaedia* 17, *Hedychium* no. 5. London. <https://www.biodiversitylibrary.org/page/39486751>
- SMITH R. M. 1994. — Zingiberaceae, in NOLTIE H. J. (ed.), *Flora of Bhutan: including a record of plants from Sikkim and Darjeeling* 3 (1). Royal Botanic Garden Edinburgh: 182-209.
- SRIVASTAVA S. C. 1984. — *A Taxonomic Study of Genus Hedychium Koen. (Zingiberaceae) in India and its Vicinity*. University of Calcutta, West Bengal, 250 p.
- WALLICH N. 1853. — Initiatory attempt to define the species of *Hedychium*, and settle their synonymy. *Hooker's Journal of Botany and Kew Garden Miscellany* 5: 321-329, 367-377. <https://www.biodiversitylibrary.org/page/783982>
- WATSON M. F. & NOLTIE H. J. 2016. — Career, collections, reports and publications of Dr. Francis Buchanan (later Hamilton), 1762-1829: natural history studies in Nepal, Burma (Myanmar), Bangladesh and India. Part 1. *Annals of Science* 73: 392-424. <https://doi.org/10.1080/00033790.2016.1195446>
- WOOD T. H., WHITTEN W. M. & WILLIAMS N. H. 2000. — Phylogeny of *Hedychium* and related genera (Zingiberaceae) based on ITS sequence data. *Edinburgh Journal of Botany* 57: 261-270. <https://doi.org/10.1017/S0960428600000196>

*Submitted on 23 March 2025;
accepted on 25 May 2025;
published on 19 January 2026.*