



General Palaeontology, Systematics, and Evolution (Vertebrate Palaeontology)

A new Late Cretaceous snake from Patagonia: Phylogeny and trends in body size evolution of madtsoiid snakes



Un nouveau serpent du Crétacé supérieur de Patagonie : phylogénie et tendances de l'évolution de la taille des serpents madtsoiidés

Raúl O. Gómez^{a,b,*}, Fernando F. Garberoglio^c, Guillermo W. Rougier^d

^a CONICET–Departamento de Ciencias Geológicas, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad de Buenos Aires, Pabellón 2, Ciudad Universitaria, 1428 Buenos Aires, Argentina

^b Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad de Buenos Aires, Pabellón 2, Ciudad Universitaria, 1428 Buenos Aires, Argentina

^c CONICET–Área de Paleontología, Fundación de Historia Natural Félix de Azara. CEBBAD, Universidad Maimónides, Hidalgo 775, 1405 Buenos Aires, Argentina

^d Department of Anatomical Sciences and Neurobiology, University of Louisville, 511 S. Floyd Street, 40292 Louisville, KY, USA

ARTICLE INFO

Article history:

Received 20 April 2019

Accepted after revision 30 September 2019

Available online 19 November 2019

Handled by Annelise Folie

Keywords:

Ophidia
Serpentes
Madtsoiidae
Phylogenetics
La Colonia Formation

Mots clés :

Ophidia
Serpentes
Madtsoiidae
Approche phylogénétique
Formation La Colonia

ABSTRACT

Madtsoiids constitute a successful group of extinct snakes widely distributed across Gondwana and the European archipelago during Late Cretaceous times, surviving in reduced numbers to the Pleistocene. They are renowned for including some of the largest snakes that have ever crawled on earth, yet diverse small madtsoiids are also known. Uncovering the evolutionary trends that led these snakes into disparate body sizes has been hampered mainly by the lack of phylogenetic consensus and the paucity of taxa with novel combinations of features. Here we describe a new large madtsoiid snake based on isolated vertebrae from the La Colonia Formation (Maastrichtian–Danian) of Patagonia, Argentina. A comprehensive phylogenetic analysis recovers Madtsoiidae as a basal ophidian lineage and the new snake as sister to a clade of mostly big-to-gigantic taxa, providing insights into early stages and evolutionary trends towards madtsoiid gigantism.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Les madtsoiidés constituent un groupe réussi de serpents éteints, largement répartis dans le Gondwana et l'archipel européen au cours du Crétacé supérieur, survivant en nombre réduit jusqu'au Pléistocène. Ils sont réputés pour inclure certains des plus grands serpents qui aient jamais rampé sur Terre, mais divers petits madtsoiidés sont également connus. La découverte des tendances évolutives qui ont conduit ces serpents à des tailles de corps disparates a été entravée principalement par le manque de consensus phylogénétique et le manque de taxons dotés de nouvelles combinaisons de caractéristiques. Nous décrivons ici un nouveau grand serpent madtsoiidé sur la base de vertèbres isolées de la formation

* Corresponding author at: Departamento de Ciencias Geológicas, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad de Buenos Aires, Pabellón 2, Ciudad Universitaria, 1428 Buenos Aires, Argentina.

E-mail address: raulorenciogomez@gmail.com (R.O. Gómez).

t La Colonia (Maastrichtien–Danien) de Patagonie, Argentine. Une analyse phylogénétique complète récupère les Madtsoiidae en tant que lignée basale ophidienne et le nouveau serpent en tant que frère d'un clade de taxons pour la plupart énormes à gigantesques, fournissant un aperçu des débuts et des tendances évolutives vers le gigantisme des madtsoïdés.

© 2019 Académie des sciences. Publié par Elsevier Masson SAS. Tous droits réservés.

1. Introduction

Madtsoiids compose a successful group of extinct snakes with a relatively rich fossil record spanning 100 million years and a widespread distribution across Gondwana and the European archipelago in the Late Cretaceous (reviewed by Rage et al., 2014; Rio and Mannion, 2017; Vasile et al., 2013). Their monophyly and position within snakes have been intensely debated (McDowell, 1987; Rage, 1984, 1998; Rieppel et al., 2002; Scanlon, 2006; Scanlon and Lee, 2000), and even their internal relationships remain uncertain (Rio and Mannion, 2017; Vasile et al., 2013). However, most recent studies support madtsoiids as a clade of basal snakes (Caldwell et al., 2015; Garberoglio et al., 2019a, 2019b; Harrington and Reeder, 2017; Martill et al., 2015; Vasile et al., 2013). Two basal snakes from the Upper Cretaceous of Patagonia might also be related to madtsoiids (Garberoglio et al., 2019a, 2019b): *Dinilysia patagonica* Smith-Woodward, 1901 and *Najash rionegrina* Apesteguía and Zaher, 2006, both well known from crania and postcrania (Caldwell and Albino, 2002; Garberoglio et al., 2019a, 2019b; Scanferla and Bhullar, 2014).

Madtsoiids are renowned for including some of the largest snakes that have ever crawled on earth; large forms are already present in the Latest Cretaceous of India, Madagascar, and Europe (LaDuke et al., 2010; Mohabey et al., 2011; Wilson et al., 2010), becoming truly gigantic worldwide in the Paleogene (Rage et al., 2014; Rio and Mannion, 2017; Simpson, 1933). Diverse small-to-medium-sized madtsoiids are also known over most of the stratigraphic range of the group (Pritchard et al., 2014; Rage, 1996; Scanlon, 1997, 2006; Vasile et al., 2013), which are the only size-class of unambiguous madtsoiids represented to date in the Upper Cretaceous of Patagonia (Albino, 1986, 1994, 2000, 2007; Gómez, 2006, 2011; Gómez and Báez, 2005, 2006; Martinelli and Forasiepi, 2014). Understanding how and when these disparate body sizes have evolved among madtsoiids has been hampered mainly by the lack of resolution and consensus on their phylogeny (Rio and Mannion, 2017; Vasile et al., 2013), but also by the paucity of madtsoiid taxa in general, and particularly, new taxa showing novel character combinations.

Here we describe a large new Late Cretaceous madtsoiid snake from the La Colonia Formation of Patagonia based on isolated vertebrae. We assess the evolutionary relationships of the new and previously known madtsoiids by means of a comprehensive phylogenetic analysis and discuss major trends of body size evolution in Madtsoiidae.

2. Geological and paleontological framework

The fossils described herein come from middle levels of the La Colonia Formation (Pesce, 1979) cropping out on the southeastern slopes of the Somún Curá Massif, Chubut Province, Argentina (Fig. 1A). Particularly, they were exhumed from a small fossil-rich lens at El Uruguayo site (Harper et al., 2018; Rougier et al., 2009a) located on the north-facing slopes of the Mirasol Chico Creek Valley (Fig. 1B). These levels correspond to the second facies association of Pascual et al. (2000), which is the most representative of the unit and consists mostly of massive and laminated claystones and siltstones with scarce intercalated beds of massive, laminated, or cross-bedded sandstones (see stratigraphic sections north of Cerro Bayo in Gasparini et al., 2015: fig. 2). Their depositional environments have alternatively been interpreted as marginal marine, estuaries, tidal flats, lagoons or coastal plains or as more restricted littoral settings including muddy floodplains, marshes or ponds, both influenced by continental freshwater flows and tidal currents from the sea (Ardolino and Delpino, 1987; Cúneo et al., 2014; Gasparini et al., 2015; Page et al., 1999; Pascual et al., 2000). Deposition of these levels occurred during the Late Cretaceous–Paleocene Atlantic transgression (Gasparini et al., 2015; Malumián and Náñez, 2011) that extensively flooded Patagonia with the epeiric Kawas Sea (Casamiquela, 1978; Hugo and Leanza, 2001).

The La Colonia Formation has often been regarded as encompassing, at least, the Campanian–Maastrichtian interval based on microfossils (Ardolino and Franchi, 1996), stratigraphic correlations, and faunal comparisons (Gasparini et al., 2015; Pascual et al., 2000). Noteworthy, Pascual et al. (2000) considered the lower facies association of the La Colonia Formation as equivalent to the Puntudo Chico Formation (Pesce, 1979), a proposal that was, either implicitly or explicitly, followed by many subsequent studies (Gandolfo et al., 2014; Gasparini et al., 2015; O'Gorman et al., 2013; Rougier et al., 2009a; Sterli and de la Fuente, 2011). However, the deposits of the Puntudo Chico Formation (middle-late Campanian–early Maastrichtian) could not be attributed to the La Colonia Formation based on available data (Ardolino *vide* Vera et al., 2019) and should instead be considered as a separate unit underlying the latter in the area (Anselmi et al., 2004; Cúneo et al., 2014; Vera et al., 2019). Furthermore, the uppermost levels of the La Colonia Formation in the Telsen area were regarded as Danian based on palynomorphs (Guler et al., 2014), whereas the mammals from its middle levels at El Uruguayo have been considered slightly younger than those from the Campanian–Maastrichtian Los Alamitos and




Fig. 1. Locality. **A.** Map of Chubut province, Patagonia, Argentina, showing the area of interest on the southeastern slopes of the Somún Curá Massif. **B.** Detail of the study area and location of El Uruguayo fossil site (white star).

Fig. 1. Localité. **A.** Carte de la province de Chubut, Patagonie, Argentine, montrant la zone d'intérêt sur les versants sud-est du massif du Somún Curá. **B.** Détail de la zone d'étude et de l'emplacement du site fossilifère d'El Uruguayo (étoile blanche).

Allen formations (Rougier et al., 2009a, 2009b). Based on all available evidence, the age of the La Colonia Formation might be considered Maastrichtian–Danian (Anselmi et al., 2004; Page et al., 1999; Vera et al., 2019) and based on the geological framework and associated fauna of the El Uruguayo lens it is most likely that the snake-bearing levels are therefore late Maastrichtian in age (Harper et al., 2018; Rougier et al., 2009a, 2009b).

Abundant fossils of algae, plants, crustaceans, and mostly terrestrial, freshwater or brackish vertebrates indicative of tropical or subtropical wetlands have been recovered to date from the La Colonia Formation from different stratigraphic levels and localities (Cúneo et al., 2014; Gandolfo et al., 2014; Gasparini et al., 2015), although most vertebrate remains come from the middle levels of the unit that crop on the area of Sierra de La Colonia and Sierra de Los Tehuelches (Albino, 2000; Gasparini et al., 2015; Harper et al., 2018; O'Gorman et al., 2013; Rougier et al., 2009a; Sterli and de la Fuente, 2011). At El Uruguayo site, these levels have yielded a rich assemblage of vertebrates including fishes, amphibians, mammals, plesiosaurs, crocodiles, non-avian dinosaurs, enantiornithine birds, turtles, and snakes (Harper et al., 2018; Lawver et al., 2011; Rougier et al., 2009a). Only a few fragmentary snake vertebrae from middle levels of the La Colonia Formation between El Buitre and Buitre Chico hills have been described so far and were ascribed to the small madtsoiid *Alamitophis argentinus*, ?Madtsoiidae, ?Boidae or Madtsoiidae, and a 'Serpentes incertae sedis' (Albino, 2000). *Alamitophis argentinus* as well as other small-to-mid sized madtsoiids have also been reported based on isolated vertebrae from the Campanian–Maastrichtian Los Alamos and Allen formations of northern Patagonia (Albino, 1986, 1994, 2007; Gómez, 2006; Gómez and Báez, 2005, 2006; Martinelli and Forasiepi, 2014).

3. Materials and methods

3.1. Materials, measurements, and nomenclature

The fossil material object of this contribution has been collected along successive fieldtrips led by one of us (GWR) in collaboration with the Museo Paleontológico Egidio Feruglio to Mesozoic outcrops in northern central Chubut Province (Fig. 1). Fossils are relatively well preserved but show weathering and polished breakage surfaces, particularly in articulation facets, likely due to hydraulic transport prior to deposition (Harper et al., 2018; Varela and Parras, 2013). Measurements were mainly those defined by LaDuke (1991) and Rage (2001) and subsequently used by several authors (Rio and Mannion, 2017; Vasile et al., 2013) and were either taken digitally from photographs or directly on vertebrae with a manual digital caliper (0.01 mm error). Drawings were made with the aid of a Zeiss Stemi SV11 binocular microscope equipped with a camera lucida and photographs were taken with a Nikon D3200 digital camera equipped with a macro lens. The osteological terminology follows that of Rage (1984, 1998, 2001) and LaDuke (1991), with a few additional terms from Scanferla and Canale (2007) and Garberoglio et al. (2019a). The collected fossils are housed at the Museo Paleontológico Egidio Feruglio.

3.2. Phylogenetic analyses

The position of the new taxon from La Colonia Formation as well as the monophyly and internal relationships of Madtsoiidae were tested through a quantitative phylogenetic analysis. For this we used a modified version of the data set of Garberoglio et al. (2019a), which in turn stemmed from that of Caldwell et al. (2015). For

the purpose of this study, we added several additional madtsoiid taxa known mostly or only from vertebrae and vertebral characters based on the data set of Vasile et al. (2013), as later modified by Rio and Mannion (2017). Also, we included the presumably basal snake *Seismophis* from the Upper Cretaceous of Brazil (Hsiou et al., 2014), which is known from *Najash*-like vertebrae (Garberoglio et al., 2019a), and the Cretaceous marine snake *Simoliophis*, also known only from isolated vertebrae (Rage et al., 2016). In total, 18 extant and 36 extinct taxa were scored for 264 osteological characters, of which 52 are of the axial skeleton (Supplementary data).

A maximum parsimony analysis of the complete data set was conducted in TNT Version 1.5-beta (Goloboff et al., 2008). Heuristic searches under equal weights consist of 500 tree replicates obtained by random addition sequence (RAS), searching for new tree topologies with tree bisection and reconnection (TBR), and saving 100 trees per replication and collapsing branches of zero length after tree search. The resulting trees were subjected to a final round of TBR branch swapping and optimal topologies were rooted with *Varanus*. During tree searches, multi-state characters based on continuous underlying data or depicting clear morphoclines (Wiens, 2001) were treated as ordered (Supplementary data). Node support was estimated by calculation of the Bremer support (BS) and absolute frequencies under jackknifing, performing 10,000 pseudo-replicates of 10 random sequence additions each followed by TBR swapping, keeping up to 10 trees, with a probability of alteration $P=0.36$.

3.3. Body size evolution

We explored trends in body size evolution among madtsoiids by mapping a proxy of absolute size onto a stratocladogram. As a proxy of size we measured the maximum width of cotyle of middle precloacal vertebrae (Supplementary data). Measurements were either taken digitally from photographs or directly on vertebrae with a manual digital caliper (0.01 mm error). The stratocladogram was derived from one of the optimal trees obtained in the parsimony analysis, including madtsoiids plus a few outgroup taxa. Numerical ages of each extinct taxon were derived from the youngest possible age of the oldest fossil-bearing unit following the current chronostratigraphic framework (Cohen et al., 2013, updated) and internodes were set to a minimum time interval (1 Myr herein), following previous approaches (Gómez and Pérez-Ben, 2019; Marjanović and Laurin, 2008, 2014). Stratocladogram assemblage and character optimization under squared change parsimony were performed in Mesquite Version 3.31 (Maddison and Maddison, 2011).

3.4. Institutional abbreviations

MPEF-PV: Vertebrate Paleontological collection, Museo Paleontológico Egidio Feruglio, Trelew, Chubut Province, Argentina; MML-PV: Vertebrate Paleontological collection, Museo Municipal de Lamarque, Lamarque, Río Negro Province, Argentina.

3.5. Anatomical abbreviations

ar, Arqual ridge; cd, condyle; cot, cotyle; hk, haemal keel; izc, interzygapophyseal constriction; izr, interzygapophyseal ridge; lf, lateral foramen; lr, lateral ridge; na, neural arch; naf, neural arch foramen; nc, neural canal; ns, neural spine; pcf, paracotylar foramen; pe, posterior embayment; prz, prezygapophysis; pzf, parazygantral foramen; scf, subcentral fossa; scr, subcentral ridge; sy, synapophyses; zp, zygosphenes; zt, zygantrum.

4. Systematic Paleontology

Squamata **Oppel, 1811**

Ophidia **Brongniart, 1800** (*sensu* Lee, 1998)

Madtsoiidae (**Hoffstetter, 1961**) **McDowell, 1987**

Eomadtsoia gen. nov.

Etymology. From *eos*, Greek for dawn, for the early divergence of a large madtsoiid at the base of a clade of gigantic forms, and *Madtsoia* **Simpson, 1933**, type genus of Madtsoiidae.

Diagnosis. As for the type and only known species.

Eomadtsoia ragei sp. nov.

Figs. 2 and 3

Etymology. *ragei*, in honor of Jean-Claude Rage for his outstanding work on fossil snakes.

Holotype. MPEF-PV 2378, isolated middle precloacal vertebra (Fig. 2A–E).

Type locality and horizon. The holotype comes from El Uruguayo fossil site (**Rougier et al., 2009a**), southeastern slopes of the Somún Curá Massif, Chubut Province, Patagonia, Argentina. Second facies association (*sensu* Pascual et al., 2000) of the La Colonia Formation (Maastrichtian–Danian).

Referred specimens. MPEF-PV 641, 2379–2387, several isolated precloacal vertebrae.

Tentatively referred specimens. MPEF-PV 644–649, 2388–2389, fragmentary precloacal vertebrae and fragments of vertebrae.

Geographic and stratigraphic provenance. The referred materials come from El Uruguayo fossil site (**Rougier et al., 2009a**) or from a nearby locality north of El Buitre hill, southeastern slopes of the Somún Curá Massif, Chubut Province, Patagonia, Argentina. Second facies association (*sensu* Pascual et al., 2000) of the La Colonia Formation (Maastrichtian–Danian).

4.1. Diagnosis

Large madtsoiid with wide and short, massively built mid-precloacal vertebrae showing a single parazygantral foramen located in an elliptical shallow depression on each side of the zygantrum and lacking prezygapophyseal processes. *Eomadtsoia* is distinct from all other madtsoiids by the following combination of features (autapomorphies indicated by character and character state numbers): moderately vaulted neural arch shallowly indented posteriorly (215:0) lacking dorsolateral (= parasagittal) ridges




Fig. 2. *Eomadtsoia ragei* gen. et sp. nov. MPEF-PV 2378 (Holotype) in: anterior (A), left lateral (B), posterior (C), dorsal (D), and ventral views (E). See anatomical abbreviations in Section 3.5.

Fig. 2. *Eomadtsoia ragei* gen. et sp. nov. MPEF-PV 2378 (Holotype) en vues : antérieure (A), latérale gauche (B), postérieure (C), dorsale (D) et ventrale (E). Voir les abréviations anatomiques en Section 3.5.




Fig. 3. *Eomadtsoia ragei* gen. et sp. nov. Referred specimens. MPEF-PV 2380 in anterior (A), right lateral (B), and ventral views (C). MPEF-PV 2381 in anterior view (D). MPEF-PV 2382 in lateral view (E). MPEF-PV 2383 in dorsal view. All to the same scale. See anatomical abbreviations in Section 3.5.

Fig. 3. *Eomadtsoia ragei* gen. et sp. nov. Spécimens référés. MPEF-PV 2380 en vues : antérieure (A), latérale droite (B) et ventrale (C). MPEF-PV 2381 en vue antérieure (D). MPEF-PV 2382 en vue latérale (E). MPEF-PV 2383 en vue dorsale. Tous à la même échelle. Voir les abréviations anatomiques en Section 3.5.

(226:0), pierced by paired small foramina, and bearing a low neural spine (249:2); thick zygosphene nearly as wide as the cotyle; very short and wide triangular centrum in ventral view having clearly laterally convex subcentral ridges; narrow haemal keel clearly demarcated from the centrum by subcentral fossae pierced by small subcentral

foramina; large lateral foramina associated with short lateral ridges; at least a large paracotylar foramen on each side of the cotyle; and finally, synapophyses broad and medial to prezygapophyseal lateral edge (221:1).

Eomadtsoia further differs from most other madtsoiids from Patagonia by its markedly larger size. Additionally,

it differs from the mid-sized *Rionegrophis* (Albino, 1986, 2007; ROG pers. observ. based on MML-PV91–115) by the lower neural spine and the less laterally extended synapophyses. It is easily distinguished from *Madtsoia* by the well-demarcated, narrow and sharp haemal keel and the lack of parasagittal ridges.

4.2. Description

All the vertebrae identified as belonging to *Eomadtsoia* are from the precloacal region. Among the well-preserved vertebrae, MPEF-PV 2383 is one of the largest with a cotylar width of 10.7 mm, but a specimen tentatively referred to *E. ragei* that consists of an isolated zygosphenes and portion of the neural arch (MPEF-PV 2388) is noticeably larger than the remaining material, indicating an even larger maximum size for this species.

Anterior view. The cotyle is big and subcircular, slightly wider than high (Fig. 2A). The robust zygosphenes is as wide as the cotyle and relatively thick (low width/height ratio). The zygosphenes has moderately inclined articular facets (angle between 25–35 degrees from vertical), and an almost straight, horizontal, dorsal margin. The neural canal is relatively small, about half the area of the cotyle, and subtrifoliate in cross section. The prezygapophyses are stout, well projected laterally and moderately inclined, up to 15–20° from horizontal. There is no prezygapophyseal process, though a subtle dorsoventral crest on the prezygapophyseal buttress is present, which extends to the anterodorsal edge of the synapophyses. The synapophyses are heavily weathered and, thus, diapophysis and parapophysis do not differentiate well. It is clear though that the lateral edge of the diapophyses remain clearly medial to the lateral limit of the prezygapophyses, whereas the parapophyses surpass the cotylar margin ventrally. Deep paracotylar foramina, at least one on each side of the cotyle, are located in shallow depressions that are ventrally delimited by blunt short buttresses (Figs. 2A and 3A).

Lateral view. The neural spine is nearly as long as the neural arch, very low along its length, and only slightly slanted posteriorly (Fig. 3B and E). The zygosphenal articular facets are oval in shape. The condyle is eroded in all specimens, but it is clearly posterodorsally oriented and well demarcated from the centrum by a slight precondylar constriction. The synapophyses are broad and dorsoventrally elongated. A thick and oblique lateral ridge is more or less evident among specimens (Figs. 2B, 3B, 3E). The subcentral ridge is almost straight all along its length. The interzygapophyseal ridge is distinct and sigmoid-shaped (Fig. 3E). There is a big lateral foramen of central position, below the above-mentioned lateral ridge. A small lateral ridge or bulge can be present on the lateral wall, below the lateral foramen.

Posterior view. The neural arch is only moderately vaulted (Fig. 2C). The condyle is subcircular. The parazygantral foramina are wide, with a distinct elliptical shape and obliquely oriented, one on each side of the zygantrum, located in small fossae. The zygantral roof is slightly convex ventrally. A pair of zygantral foramina is located on separate deep rounded fossae. The synapophyses are

posterolaterally oriented, showing nearly all their articular surface in this view.

Dorsal view. The vertebrae are wider than long with a neural arch only slightly constricted (Fig. 2D). The interzygapophyseal width (minimal neural arch width) is almost the same as the prezygapophyseal width. The prezygapophyseal facets are anterolaterally oriented (angle between 60–80° from the anteroposterior axis) and have a nearly oval outline, although slight posterolateral and anteromedial corners exist. The zygosphenal anterior margin is almost straight, with only a very shallow medial notch. The neural spine is narrow, extending from the back of the anterior margin of the zygosphenes, and widens as it extends posteriorly. The neural spine likely has a posterior tubercle based on its terminal widening, but its surface is not well preserved enough to confirm this feature. To both sides of the neural spine, posteriorly to the zygosphenal base, several foramina are present on the neural arch (at least one to each side in all available vertebrae). Posteriorly, the neural arch shows a shallow embayment. The parasagittal ridges are not developed on the neural arch, which in turn is not markedly faceted. Arqual ridges are present on the posterior dorsal margin of the neural arch, anteromedial to the postzygapophyseal lappet (Fig. 3F).

Ventral view. The centrum is very short and subtriangular (Fig. 2E). The subcentral ridges are straight to slightly convex laterally. Much of the synapophyseal articular surface is visible in ventral view. The haemal keel is well developed and demarcated, low and narrow, laterally is delimited by shallow depressions, where small subcentral foramina are located in anterior position. The haemal keel contacts the condylar constriction posteriorly, not reaching the condyle. Posterior processes on the haemal keel are absent.

5. Results and discussion

5.1. Taxonomic comparisons

Among the four snakes that have been previously reported from La Colonia Formation, only one of these, namely the small madtsoiid *Alamitophis argentinus*, was unambiguously referred to family level (Albino, 2000). A single vertebra (MPEF-PV 642) originally reported as a ‘*Serpentes incertae sedis*’ (Albino, 2000, 2007) is more likely to represent a dolichosaur rather than a snake (Albino, 2011; Scanlon and Hocknull, 2008). The remaining snakes from this unit, formerly reported as a possible madtsoiid and tentatively as a boid or madtsoiid (Albino, 2000), were based on poorly preserved material that precluded a better taxonomic assignment at the time. Now, additional and better preserved snake vertebrae from the La Colonia Formation that show a unique combination of characters justifying the erection of a new taxon, *Eomadtsoia ragei* gen et sp. nov., allow referral of previously known material of uncertain madtsoiids or boids to this same taxon. Therefore, the snake assemblage from La Colonia Formation is restricted to madtsoiids, with *E. ragei* increasing the known diversity of this basal snake radiation.

Among snakes, the overall vertebral configuration of *E. ragei* is only present in the Madtsoiidae and boids, but

it can be set apart from extinct and extant boids by the presence of distinct parazygantral foramina along with paracotylar foramina, arqual ridges on the neural arch, and the absence of prezygapophyseal processes (Gómez and Báez, 2005; Rage, 1998; Scanlon, 2005). It is readily distinguishable from small-to-mid-sized madtsoiids from the Late Cretaceous of Patagonia and Europe and Paleogene of Australia, namely *Alamitophis*, *Herensugea*, *Nanowana*, *Nidophis*, *Patagoniophis*, *Rionegrophis*, and *Wonambi barrii* (Albino, 1986, 1994, 2007; Rage, 1996; Scanlon, 1997, 2005; Scanlon and Lee, 2000; Vasile et al., 2013), by its conspicuous larger size, along with the wide and short triangular centrum and the distinctly medial position of the synapophyses relative to the lateral edge of the prezygapophyses. *Eomadtsoia* clearly contrasts with the much older *Dinilysia* (Caldwell and Albino, 2002; Scanferla and Canale, 2007), which is the only other Late Cretaceous large snake from Patagonia, in having parazygantral foramina, lower neural spine, and poorly constricted neural arch, among other features.

Noteworthy, the narrow and sharp haemal keel of *Eomadtsoia* contrasts with that of large-to-gigantic madtsoiids, such as species of *Gigantophis*, *Madtsoia*, *Menarana*, and *Platyspondylophis*, in which the haemal keel is a somewhat broad and flat process more or less demarcated from the rest of the centrum (Andrews, 1901; LaDuke et al., 2010; Rage, 1996, 1998; Rage et al., 2014; Rio and Mannion, 2017; Smith et al., 2016). In addition, an interzygapophyseal constriction as shallow as that of *E. ragei* is rarely found among large madtsoiids, which generally show a more marked constriction. The lack of parasagittal ridges of *Eomadtsoia* is also uncommon, only resembling *Herensugea* (Rage, 1996) and *Adinophis* (Pritchard et al., 2014) among the surveyed madtsoiids; although some intracolumnar variation cannot be ruled out taken into account what is seen in *Gigantophis* (Rio and Mannion, 2017).

5.2. Phylogenetic analysis

The parsimony analysis yielded 42 most parsimonious trees (MPTs) of 803 steps (consistency index [CI]=0.400; retention index [RI]=0.697), the strict consensus of which is relatively well resolved and depicts all madtsoiids in a clade also including the basal snakes *Najash*, *Dinilysia*, and *Seismophis* (Fig. 4). In agreement with recent hypotheses, these snakes lie outside crown-group Serpentes (Caldwell et al., 2015; Garberoglio et al., 2019a, 2019b; Harrington and Reeder, 2017; Martill et al., 2015; Vasile et al., 2013). The monophyly of madtsoiids is here rigorously tested, since we use as outgroup taxa all major groups of extant snakes plus several Mesozoic snakes and the lizard *Varanus* for rooting (Fig. S1), instead of solely the basal snakes *Najash* and *Dinilysia* as in previous similar studies (Rio and Mannion, 2017).

Our results depict *Najash* and *Dinilysia* forming a clade crown-ward to *Seismophis* and all nested within taxa usually considered part of Madtsoiidae, since the poorly known *Herensugea* from the Maastrichtian of Iberia (Rage, 1996) emerges as an early diverging taxon sister to all other snakes in this clade; a position that contrasts with that recovered in previous studies (Rio and Mannion, 2017;

Vasile et al., 2013). The isolated vertebrae of *Seismophis* recall in many aspects the vertebrae of *Najash* (Garberoglio et al., 2019a; Hsiou et al., 2014), both from the Cenomanian of South America and, hence, the oldest members of this clade. Interestingly, the vertebral morphology of *Herensugea* is quite unique among madtsoiids, having depressed neural arch with low parasagittal keels, and prezygapophyses and main axis of zygosphenal facets close to horizontal (Rage, 1996), and it would not be surprising if it represented an end-Cretaceous relict of a much earlier snake radiation along with a still unnamed putative madtsoiid from the Cenomanian of North Africa (Rage and Dutheil, 2008).

All these basal snakes from Gondwana and neighboring terrains share the presence of conspicuous parazygantral foramina, which has been considered the clearest synapomorphy of Madtsoiidae by most authors (Albino, 2000; Rage, 1984, 1998; Rio and Mannion, 2017; Scanlon, 2005; Vasile et al., 2013), although they have also been reported as irregularly present in a few other snakes (Holman and Case, 1992; Rage, 1998; Rage et al., 2004). In view of our results depicting *Seismophis*, *Najash*, and *Dinilysia* basal to all madtsoiids other than *Herensugea*, and that the anatomy and relationships of the enigmatic *Herensugea* needs further scrutiny, we opt to not strictly consider the former taxa as part of Madtsoiidae and instead regard the parazygantral foramen as synapomorphic of a more inclusive clade than traditionally acknowledged.

Our results recover all large-bodied madtsoiids as a clade sister to mostly small-sized madtsoiids (Fig. 4), mirroring to some extent previous hypotheses derived from comprehensive analyses of Madtsoiidae (Rio and Mannion, 2017; Vasile et al., 2013). However, our tree topology showing two clades reflecting overall size has not been recovered previously (Vasile et al., 2013: fig. 5A). Large madtsoiids are not members of a single clade either in the analysis of Rio and Mannion (2017: fig. 10B). These size-related groups are here recovered despite the fact that ‘body size’, in contrast with the previous studies, was not included as a discrete character in our analysis.

Eomadtsoia ragei is here recovered as an early diverging lineage of the clade of large madtsoiids, crown-ward to *Sanajeh* and *Rionegrophis* and sister to a poorly resolved clade encompassing snakes in the genera *Gigantophis*, *Madtsoia*, *Menarana*, and *Platyspondylophis* from the Maastrichtian–Eocene interval, as well as the younger *Wonambi* and *Yurlunggur* from Australia (Fig. 4). Interestingly, deeply nested within this clade also appears the small *Adinophis fisaka* (Pritchard et al., 2014), which was previously recovered as more closely related to other small madtsoiids (Rio and Mannion, 2017). None of the madtsoiid genera represented by more than one species have been recovered as monophyletic neither here nor in previous analyses (Rio and Mannion, 2017; Vasile et al., 2013), calling for further revision of the taxonomy of these snakes.

5.3. Body size evolution

Gigantic Madtsoiidae are some of the largest snakes that have ever existed, reaching in some cases 10 m of total length (Head et al., 2009; Rio and Mannion, 2017), but according to our results they evolved from small-to-mid-




Fig. 4. Phylogenetic relationships of *Eomadtsoia ragei* gen. et sp. nov. (bold) and other madtsoiids. Strict consensus of 42 most parsimonious trees of 803 steps. For simplification, crown-group Serpentes is depicted by a solid triangle (Supplementary data).

Fig. 4. Relations phylogénétiques de *Eomadtsoia ragei* gen. et sp. nov. (gras) et autres madtsoïdés. Consensus strict des 42 arbres les plus parcimonieux de 803 pas. Pour simplifier, le groupe couronne Serpentes est représenté par un triangle plein (voir les données supplémentaires).

sized forms at some point during the Cretaceous (Fig. 5), likely in the Late Cretaceous of Gondwana. Among the closest relatives of true madtsoiids are the relatively small snakes *Seismophis* and *Najash*, but also the much larger *Dinilysia* that might have exceeded 1.8 m in length (Yi and Norell, 2015), suggesting that this early snake radiation was not constrained in size, which is also echoed by the initial split of madtsoiids into sister clades of disparate body size.

Different body-size evolutionary trends characterize these madtsoiid clades. That of small forms, which appear to have radiated to reach a widespread geographic distribution during the Maastrichtian (Vasile et al., 2013), shows relative stasis in body size over time. Into the Cenozoic they

remained diverse apparently only in Australia (Scanlon, 1997, 2005), although Paleogene putative madtsoiids scattered across Gondwanan landmasses (Rage, 1991; Rage et al., 2008) might also belong to this group. Conversely, its sister clade shows a general trend of increased body size since the Latest Cretaceous with several independent transitions towards gigantism prior to the K/Pg boundary (Fig. 5). In addition, this subclade exhibits a higher diversity of body sizes with some reversal shifts to a smaller size in *Menarana nosymena* and, more markedly, *Adinophis fisaka*, both from Madagascar (LaDuke et al., 2010; Pritchard et al., 2014).

In the current phylogenetic framework, *Eomadtsoia*, together with *Sanajeh* from the Maastrichtian of India and




Fig. 5. Evolution of body size in madtsoiid snakes. Squared-change-parsimony optimization of cotyle width on a most parsimonious tree adjusted in time, depicting the ancestral reconstructions for small and large madtsoiids. Diagrams of vertebrae are at natural size.

Fig. 5. Évolution de la taille des serpents madtsoïdés. Optimisation en *squared-change-parsimony* de la largeur du cotyle sur un arbre très parcimonieux ajusté dans le temps, illustrant les reconstructions ancestrales de madtsoïdés de petite et grande taille. Les diagrammes des vertèbres sont à la taille naturelle.

Rionegrophis from the slightly older Los Alamos and Allen formations of Patagonia (Albino, 1986, 2007; Gómez, 2006), emerge as early diverging mid-to-large madtsoiids with vertebral morphologies and body sizes that could be interpreted as foretelling of the size increase characteristic of the more nested clades. *Eomadtsoia* fits in a morphocline, linking the ancestral madtsoiid conditions with those of large-to-gigantic taxa such as *Madtsoia bai* from the Eocene of Patagonia (Simpson, 1933).

6. Conclusions

Eomadtsoia ragei gen. et sp. nov. from the La Colonia Formation (Maastrichtian–Danian) represents the largest madtsoiid snake from the Upper Cretaceous of Patagonia. Given its phylogenetic position sister to a clade that includes the largest forms, the new snake suggests that early stages and evolutionary trends towards madtsoiid gigantism occurred prior to the K/Pg boundary. Further increase of body size following the Cretaceous extinction mirrors trends seen in other groups (but see Apesteguía et al., 2014) such as amphibians (Agnolin, 2012; Gómez et al., 2011), turtles (Bona and de La Fuente, 2005), and archaic mammals (Páez Arango, 2008; Rougier et al., 2009a).

Acknowledgments

We would like to thank M. Laurin (MNHN) for inviting us to contribute to this volume honoring Jean-Claude Rage, who had inspired two of us early in our careers in the study of fossil snakes. We also thank R. Cúneo, L. Canessa, E. Ruigómez, and other personnel of the Museo Paleontológico Egidio Feruglio for years of support. Thanks are extended to J. Rio (Imperial College London), who kindly provided data for the analysis and the Willi Hennig Society for the free internet access to the TNT software package. We are also grateful to the University of Buenos Aires and the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) of Argentina for continuous support. This research was funded by NSF via the DEB 0946430 and DEB 1068089 grants (to GWR), USA, and by the RAICES program (PICT-2016-3682), Agencia Nacional de Promoción Científica y Tecnológica, Argentina.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.crpv.2019.09.003>.

References

- Agnolin, F., 2012. A new Calyptocephalellidae (Anura, Neobatrachia) from the Upper Cretaceous of Patagonia, Argentina, with comments on its systematic position. *Stud. Geol. Salmanticensis* 48, 129–178.
- Albino, A.M., 1986. Nuevos Boidae Madtsoiinae en el Cretácico tardío de Patagonia (Formación Los Alamitos, Río Negro, Argentina). *Actas IV Congreso Argentino de Paleontología y Bioestratigrafía* 2, 15–21.
- Albino, A.M., 1994. Una Nueva Serpiente (Reptilia) en el Cretácico Superior de Patagonia, Argentina. *Pesquisas em Geociências* 21, 58–63, <http://dx.doi.org/10.22456/1807-9806.21251>.
- Albino, A.M., 2000. New records of snakes from the Cretaceous of Patagonia, Argentina. *Geodiversitas* 22, 247–253.
- Albino, A.M., 2007. Lepidosauromorpha. In: Gasparini, Z., Salgado, L., Coria, R.A. (Eds.), *Patagonian Mesozoic Reptiles*. Indiana University Press, Bloomington, pp. 87–115.
- Albino, A.M., 2011. Evolution of Squamata Reptiles in Patagonia based on the fossil record. *Biol. J. Linn. Soc.* 103, 441–457, <http://dx.doi.org/10.1111/j.1095-8312.2011.01691.x>.
- Andrews, C.W., 1901. Preliminary notes on some recently discovered extinct vertebrates from Egypt (Part II). *Geol. Mag.* 8, 434–444, <http://dx.doi.org/10.1017/S0016756800179750>.
- Anselmi, G., Gamba, M.T., Panza, J.L., 2004. Hoja geológica 4369–IV, Los Altares. Provincia del Chubut. Instituto de Geología y Recursos Minerales. Servicio Geológico Minero Argentino. Boletín 313, 98.
- Apesteguía, S., Zaher, H., 2006. A Cretaceous terrestrial snake with robust hindlimbs and a sacrum. *Nature* 440, 1037–1040, <http://dx.doi.org/10.1038/nature04413>.
- Apesteguía, S., Gómez, R.O., Rougier, G.W., 2014. The youngest South American rhynchocephalian, a survivor of the K/Pg extinction. *Proc. R. Soc. B* 281, 20140811, <http://dx.doi.org/10.1098/rspb.2014.0811>.
- Ardolino, A., Delpino, D., 1987. Senoniano (continental-marino). Comarca Nordpatagónica, Provincia del Chubut, Argentina. X Congreso Geológico Argentino (Tucumán). *Actas* 3, 193–196.
- Ardolino, A., Franchi, M., 1996. Hoja geológica 4366–I Telsen. Provincia del Chubut. Programa Nacional de Cartas Geológicas de la República Argentina, escala 1:250.000. Dirección Nacional del Servicio Geológico, Buenos Aires. Boletín 215, 110.
- Bona, P., de La Fuente, M.S., 2005. Phylogenetic and paleobiogeographic implications of *Yaminuechelys maior* (Staesch, 1929) new comb., a large long-necked chelid turtle from the early Paleocene of Patagonia, Argentina. *J. Vert. Paleontol.* 25, 569–582, [http://dx.doi.org/10.1671/0272-4634\(2005\)025\[0569:PAPIOY\]2.0.CO;2](http://dx.doi.org/10.1671/0272-4634(2005)025[0569:PAPIOY]2.0.CO;2).
- Brongniart, A., 1800. Essai d'une classification naturelle des reptiles. 1^{re} partie. Etablissement des Orders. *Bull. Sci. Soc. Philom. Paris* 35, 81–82.
- Caldwell, M.W., Albino, A., 2002. Exceptionally preserved skeletons of the Cretaceous snake *Dinilysia patagonica* Woodward, 1901. *J. Vert. Paleontol.* 22, 861–866, [http://dx.doi.org/10.1671/0272-4634\(2002\)022\[0861:EPSCOT\]2.0.CO;2](http://dx.doi.org/10.1671/0272-4634(2002)022[0861:EPSCOT]2.0.CO;2).
- Caldwell, M.W., Nydam, R.L., Palci, A., Apesteguía, S., 2015. The oldest known snakes from the Middle Jurassic–Lower Cretaceous provide insights on snake evolution. *Nat. Commun.* 6, 5996, <http://dx.doi.org/10.1038/ncomms6996>.
- Casamiquela, R.M., 1978. La zona litoral de la transgresión mastrichtense en el norte de la Patagonia. Aspectos ecológicos. *Ameghiniana* 15, 137–148.
- Cohen, K.M., Finney, S.C., Gibbard, P.L., Fan, J.-X., 2013. The ICS International Chronostratigraphic Chart. Episodes 36, 199–204 (updated).
- Cúneo, N.R., Gandolfo, M.A., Zamalao, M.C., Hermesen, E., 2014. Late Cretaceous Aquatic Plant World in Patagonia, Argentina. *PLoS One* 9, e104749, <http://dx.doi.org/10.1371/journal.pone.0104749>.
- Gandolfo, M.A., Cúneo, N.R., Hermesen, E.J., 2014. La Paleoflora de la Formación La Colonia. Reporte preliminar sobre la paleoflora de la Formación La Colonia (Campaniano Maastrichtiano, Cretácico tardío), Chubut, Patagonia, Argentina. *Bol. Soc. Geol. Mex.* 66, 11–23.
- Garberoglio, F.F., Gómez, R.O., Apesteguía, S., Caldwell, M.W., Sánchez, M.L., Veiga, G., 2019a. A new specimen with skull and vertebrae of *Najash rionegrina* (Lepidosauria: Ophidia) from the early Late Cretaceous of Patagonia. *J. Syst. Palaeontol.* 17, 1313–1330.
- Garberoglio, F.F., Gómez, R.O., Simões, T.R., Caldwell, M.W., Apesteguía, S., 2019b. The evolution of the axial intercentrum system in snakes revealed by new data from the Cretaceous snakes *Dinilysia* and *Najash*. *Sci. Rep.* 9, 1276, <http://dx.doi.org/10.1038/s41598-018-36979-9>.
- Gasparini, Z., Sterli, J., Parras, A., O'Gorman, J.P., Salgado, L., Varela, J., Pol, D., 2015. Late Cretaceous reptilian biota of the La Colonia Formation, central Patagonia, Argentina: Occurrences, preservation and paleoenvironments. *Cretaceous Res.* 54, 154–168, <http://dx.doi.org/10.1016/j.cretres.2014.11.010>.
- Goloboff, P., Farris, J., Nixon, J.K., 2008. TNT, a free program for phylogenetic analysis. *Cladistics* 24, 774–786, <http://dx.doi.org/10.1111/j.1096-0031.2008.00217.x>.
- Gómez, R.O., (Unpublished Undergraduate thesis) 2006. Serpientes de la Formación Allen (Campaniano–Maastrichtiano), en los bajos de Trapalcó y Santa Rosa, Provincia de Río Negro, Argentina. Facultad de Ciencias Exactas y Naturales Universidad de Buenos Aires, Buenos Aires (81 p.).
- Gómez, R.O., 2011. A Snake Dentary from the Upper Cretaceous of Patagonia. *J. Herpetol.* 45, 230–233, <http://dx.doi.org/10.1670/10-048.1>.
- Gómez, R.O., Báez, A.M., 2005. Intracolumnar variation and morphological remarks on madtsoiid snakes (Squamata, Serpentes) from the Upper Cretaceous of Patagonia. *Ameghiniana* 42 (30R–31R).
- Gómez, R.O., Báez, A.M., 2006. A new madtsoiid snake (Squamata, Ophidia) from the Upper Cretaceous of Patagonia. *Ameghiniana* 43, 21R.
- Gómez, R.O., Pérez-Ben, C.M., 2019. Fossils reveal long-term continuous and parallel innovation in the sacro-caudo-pelvic complex of the highly aquatic pipid frogs. *Front. Earth Sci.* 7, 56, <http://dx.doi.org/10.3389/feart.2019.00056>.
- Gómez, R.O., Báez, A.M., Muzzopappa, P., 2011. A new helmeted frog (Anura: Calyptocephalellidae) from an Eocene subtropical lake in northwestern Patagonia, Argentina. *J. Vert. Paleontol.* 31, 50–59, <http://dx.doi.org/10.1080/02724634.2011.539654>.
- Guler, M.V., Borel, C.M., Brinkhuis, H., Navarro, E., Astini, E., Alfredo, R., 2014. Brackish to freshwater dinoflagellate cyst assemblages from the La Colonia Formation (Paleocene?), Northeastern Patagonia, Argentina. *Ameghiniana* 51, 141–153, <http://dx.doi.org/10.5710/AMGH.15.02.2014.949>.
- Harper, T., Parras, A., Rougier, G.W., 2018. *Reigitherium* (Meridiolestida, Mesungulatoidea) an enigmatic Late Cretaceous mammal from Patagonia, Argentina: Morphology, affinities, and dental evolution. *J. Mammal. Evol.*, <http://dx.doi.org/10.1007/s10914-018-9437-x>.
- Harrington, S.M., Reeder, T.W., 2017. Phylogenetic inference and divergence dating of snakes using molecules, morphology and fossils: new insights into convergent evolution of feeding morphology and limb reduction. *Biol. J. Linn. Soc.* 121, 379–394, <http://dx.doi.org/10.1093/biolinnean/blw039>.
- Head, J.J., Bloch, J.L., Hastings, A.K., Bourque, J.R., Cadena, E.A., Herrera, F.A., Polly, P.D., Jaramillo, C.A., 2009. Giant boid snake from the Palaeocene neotropics reveals hotter past equatorial temperatures. *Nature* 457, 715–718, <http://dx.doi.org/10.1038/nature07671>.
- Hoffstetter, R., 1961. Nouveaux restes d'un serpent Boidé (*Madtsoia madagascariensis* nov. sp.) dans le Crétacé supérieur de Madagascar. *Bull. Mus. Hist. Nat., Paris* 33, 152–160.
- Holman, J.A., Case, G.R., 1992. A puzzling new snake (Reptilia: Serpentes) from the late Paleocene of Mississippi. *Ann. Carnegie Mus.* 61, 197–205.
- Hsiou, A.S., Albino, A.M., Medeiros, M.A., Santos, R.A.B., 2014. The oldest Brazilian snakes from the Cenomanian (early Late Cretaceous). *Acta Paleontol. Pol.* 59, 635–642, <http://dx.doi.org/10.4202/app.2012.0091>.
- Hugo, C.A., Leanza, H.A., 2001. Hoja Geológica 3966–III, Villa Regina, provincia Río Negro. Instituto de Geología y Recursos Naturales, SEGEMAR 309, 1–53.
- LaDuke, T.C., 1991. The fossil snakes of Pit 91, Rancho La Brea, California. *Nat. Hist. Mus. LA County Contrib. Sci.* 424, 1–28.
- LaDuke, T.C., Krause, D.W., Scanlon, J.D., Kley, N.J., 2010. A Late Cretaceous (Maastrichtian) snake assemblage from the Maevran Formation, Mahajanga Basin, Madagascar. *J. Vert. Paleontol.* 30, 109–138, <http://dx.doi.org/10.1080/02724630903409188>.
- Lawver, D.R., Debee, A.J.M., Clarke, J.A., Rougier, G.W., 2011. A New Enantiornithine Bird from the Upper Cretaceous La Colonia Formation of Patagonia, Argentina. *Ann. Carnegie Mus.* 80, 35–42, <http://dx.doi.org/10.2992/007.080.0104>.
- Lee, M.S.Y., 1998. Convergent evolution and character correlation in burrowing reptiles: towards a resolution of squamate relationships. *Biol. J. Linn. Soc.* 65, 369–453, <http://dx.doi.org/10.1111/j.1095-8312.1998.tb01148.x>.
- Maddison, W.P., Maddison, D.R., 2011. Mesquite: a modular system for evolutionary analysis, Version 3. 31. <http://www.mesquiteproject.org>.
- Malumián, N., Nánuez, C., 2011. The Late Cretaceous–Cenozoic transgressions in Patagonia and the Fuegian Andes: foraminifera, palaeoecology, and palaeogeography. *Biol. J. Linn. Soc.* 103, 269–288, <http://dx.doi.org/10.1111/j.1095-8312.2011.01649.x>.
- Marjanović, D., Laurin, M., 2008. Assessing Confidence Intervals for Stratigraphic Ranges of Higher Taxa: The Case of Lissamphibia. *Acta Paleontol. Pol.* 53, 413–432, <http://dx.doi.org/10.4202/app.2008.0305>.

- Marjanović, D., Laurin, M., 2014. An updated paleontological timetable of lissamphibians, with comments on the anatomy of Jurassic crown-group salamanders (Urodela). *Hist. Biol.* 26, 535–550, <http://dx.doi.org/10.4202/app.2008.0305>.
- Martill, D.M., Tischlinger, H., Longrich, N.R., 2015. A four-legged snake from the Early Cretaceous of Gondwana. *Science* 349, 416–419, <http://dx.doi.org/10.1126/science.aaa9208>.
- Martinelli, A.G., Forasiepi, A.M., 2014. Late Cretaceous vertebrates from Bajo de Santa Rosa (Allen Formation), Río Negro Province, Argentina, with description of a new sauropod dinosaur (Titanosauridae). *Rev. Mus. Argentino Cienc. Nat., n.s.* 6, 257–305.
- McDowell, S.B., 1987. Systematics. In: Collins, J.T., Novak, S.S., Seigel, R.A. (Eds.), *Snakes: Ecology and evolutionary biology*. McGraw-Hill Publ. Co, New York, pp. 3–50.
- Mohabey, D.N., Head, J.J., Wilson, J.A., 2011. A new species of the snake *Madtsoia* from the Upper Cretaceous of India and its paleobiogeographic implications. *J. Vert. Paleontol.* 31, 588–595, <http://dx.doi.org/10.1080/02724634.2011.560220>.
- O'Gorman, J.P., Salgado, L., Cerda, I.A., Gasparini, Z., 2013. First record of gastroliths associated with elasmosaur remains from La Colonia Formation (Campanian–Maastrichtian), Chubut, Patagonia Argentina, with comments on the probable depositional palaeoenvironment of the source of the gastroliths. *Cretaceous Res.* 40, 212–217, <http://dx.doi.org/10.1016/j.cretres.2012.07.004>.
- Oppel, M., 1811. *Die Ordnungen, Familien und Gattungen der Reptilien, als Prodrom einer Natugeschichte derselben*. Joseph Lindauer, Munich.
- Páez Arango, N., (Unpublished MS thesis) 2008. Dental and craneomandibular anatomy of *Pelagotherium tropicalis*: the evolutionary radiation of South American dryolestoid mammals. University of Louisville, Louisville, USA (105 p.).
- Page, R., Ardolino, A., de Barrio, R.E., Franchi, M., Lizuain, A., Page, S., Silva Nieto, D., 1999. Estratigrafía del Jurásico y Cretácico del Macizo de Somún Curá, provincias de Río Negro y Chubut. In: Caminos, R. (Ed.), *Geología Argentina*. Servicio Geológico Minero Argentino SEGEMAR, Buenos Aires, pp. 460–488.
- Pascual, R., Goin, F.J., González, P., Ardolino, A., Puerta, P.F., 2000. A highly derived docodont from the Patagonian Late Cretaceous: evolutionary implications for Gondwanan mammals. *Geodiversitas* 22, 395–414.
- Pesce, A.H., 1979. Estratigrafía del arroyo Perdido en su tramo medio e inferior provincia del Chubut. VII Congreso Geológico Argentino (Neuquén, 1978). *Actas* 1, 315–333.
- Pritchard, A.C., McCartney, J.A., Krause, D.W., Kley, N.J., 2014. New snakes from the Upper Cretaceous (Maastrichtian) Maevarano Formation, Mahajanga Basin, Madagascar. *J. Vert. Paleontol.* 34, 1080–1093, <http://dx.doi.org/10.1080/02724634.2014.841706>.
- Rage, J.-C., 1984. *Serpentes*. Fischer, Stuttgart.
- Rage, J.-C., 1991. Squamate reptiles from the early Paleocene of Tiupampa area (Santa Lucía Formation), Bolivia. In: Suarez-Soruco, R. (Ed.), *Fósiles y Facies de Bolivia*, vol. 1. Vertebrados. Rev. Técnica YPF 12, pp. 503–508.
- Rage, J.-C., 1996. Les Madtsoiidae (Reptilia, Serpentes) du Crétacé supérieur d'Europe : témoins gondwaniens d'une dispersion transthéthysienne. *C. R. Acad. Sci. Paris, Ser. IIa* 322, 603–608.
- Rage, J.-C., 1998. Fossil snakes from the Paleocene of São José de Itaboraí, Brazil. Part I. Madtsoiidae, Aniliidae. *Palaeovertebrata* 27, 109–144.
- Rage, J.-C., 2001. Fossil snakes from the Paleocene of São José de Itaboraí, Brazil. Part II. Boidae. *Palaeovertebrata* 30, 111–150.
- Rage, J.-C., Dutheil, D.B., 2008. Amphibians and squamates from the Cretaceous (Cenomanian) of Morocco—A preliminary study, with description of a new genus of pipid frog. *Palaeontographica Abt. A Band* 285 Lieferung 1–3, 1–22, <http://dx.doi.org/10.1127/pala/285/2008/1>.
- Rage, J.-C., Folie, A., Rana, R.S., Singh, H., Rose, K.D., Smith, T., 2008. A Diverse Snake Fauna from the Early Eocene of Vastan Lignite Mine, Gujarat, India. *Acta Palaeontol. Pol.* 53, 391–403, <http://dx.doi.org/10.4202/app.2008.0303>.
- Rage, J.-C., Métais, G., Bartolini, A., Brohi, I.A., Lashari, R.A., Mariavaux, L., Merle, D., Solangi, S.H., 2014. First report of the giant snake *Gigantophis* (Madtsoiidae) from the Paleocene of Pakistan: Paleobiogeographic implications. *Geobios* 47, 147–153, <http://dx.doi.org/10.1016/j.geobios.2014.03.004>.
- Rage, J.-C., Prasad, G.V.R., Bajpai, S., 2004. Additional snakes from the uppermost Cretaceous (Maastrichtian) of India. *Cretaceous Res.* 25, 425–434, <http://dx.doi.org/10.1016/j.cretres.2004.02.003>.
- Rage, J.-C., Vullo, R., Néraudeau, D., 2016. The mid-Cretaceous snake *Simoliophis rochebrunei* Sauvage, 1880 (Squamata: Ophidia) from its type area (Charentes, southwestern France): Redescription, distribution, and palaeoecology. *Cretaceous Res.* 58, 234–253, <http://dx.doi.org/10.1016/j.cretres.2015.10.010>.
- Rieppel, O., Kluge, A.G., Zaher, H., 2002. Testing the phylogenetic relationships of the Pleistocene snake *Wonambi naracoortensis* Smith. *J. Vert. Paleontol.* 22, 812–829, [http://dx.doi.org/10.1671/0272-4634\(2002\)022\[0812:TTPROT.2.0.CO;2\]](http://dx.doi.org/10.1671/0272-4634(2002)022[0812:TTPROT.2.0.CO;2]).
- Rio, J.P., Mannion, P.D., 2017. The osteology of the giant snake *Gigantophis garstini* from the upper Eocene of North Africa and its bearing on the phylogenetic relationships and biogeography of Madtsoiidae. *J. Vert. Paleontol.* 37, e1347179, <http://dx.doi.org/10.1080/02724634.2017.1347179>.
- Rougier, G.W., Forasiepi, A.M., Hill, R.V., Novacek, M., 2009a. New mammalian remains from the Late Cretaceous La Colonia Formation, Patagonia, Argentina. *Acta Palaeontol. Pol.* 54, 195–212, <http://dx.doi.org/10.4202/app.2006.0026>.
- Rougier, G.W., Chornogubsky, L., Casadio, S., Páez Arango, N., Giallombardo, A., 2009b. Mammals from the Allen Formation, Late Cretaceous, Argentina. *Cretaceous Res.* 30, 223–238, <http://dx.doi.org/10.1016/j.cretres.2008.07.006>.
- Scanferla, C.A., Bhullar, B.-A.S., 2014. Postnatal development of the skull of *Dinilysia patagonica* (Squamata–Stem Serpentes). *Anat. Rec.* 297, 560–573, <http://dx.doi.org/10.1002/ar.22862>.
- Scanferla, C.A., Canale, J.I., 2007. The youngest record of the Cretaceous snake genus *Dinilysia* (Squamata, Serpentes). *S. Am. J. Herpetol.* 2, 76–81, [http://dx.doi.org/10.2994/1808-9798\(2007\)2\[76:TYROTC.2.0.CO;2\]](http://dx.doi.org/10.2994/1808-9798(2007)2[76:TYROTC.2.0.CO;2]).
- Scanlon, J.D., 1997. *Nanowana* gen. nov., small madtsoiid snakes from the Miocene of Riversleigh: Sympatric species with divergently specialised dentition. *Mem. Queensl. Mus.* 41, 393–412.
- Scanlon, J.D., 2005. Australia's oldest known snakes: *Patagoniophis*, *Alamitophis*, and cf. *Madtsoia* (Squamata: Madtsoiidae) from the Eocene of Queensland. *Mem. Queensl. Mus.* 51, 215–235.
- Scanlon, J.D., 2006. Skull of the large non-macrostomatan snake *Yurlungur* from the Australian Oligo-Miocene. *Nature* 439, 839–842, <http://dx.doi.org/10.1038/nature04137>.
- Scanlon, J.D., Hocknull, S.A., 2008. A dolichosaurid lizard from the latest Albian (mid-Cretaceous) Winton Formation, Queensland, Australia. *Trans. Kad. Acad. Sci., Proc. 2nd Mosasaur Meeting*, 131–136.
- Scanlon, J.D., Lee, M.S.Y., 2000. The Pleistocene serpent *Wonambi* and the early evolution of snakes. *Nature* 403, 416–420, <http://dx.doi.org/10.1038/35000188>.
- Simpson, G.G., 1933. A new fossil snake from the *Notostylops* beds of Patagonia. *Bull. Am. Mus. Nat. Hist.* 67, 1–22.
- Smith, T., Kumar, K., Rana, R.S., Folie, A., Solé, F., Noiret, C., Steeman, T., Sahni, A., Rose, K.D., 2016. New early Eocene vertebrate assemblage from western India reveals a mixed fauna of European and Gondwana affinities. *Geoscience Frontiers* 7, 969–1001, <http://dx.doi.org/10.1016/j.gsf.2016.05.001>.
- Smith-Woodward, A., 1901. On some extinct reptiles from Patagonia, of the genera *Miolania*, *Dinilysia*, and *Genyodectes*. *Proc. Zool. Soc. London*, 169–184, <http://dx.doi.org/10.1111/j.1469-7998.1901.tb08537.x>.
- Sterli, J., de la Fuente, M.S., 2011. A new turtle from La Colonia Formation (Campanian–Maastrichtian), Patagonia, Argentina, with remarks on the evolution of the vertebral column in turtles. *Palaeontology* 54, 63–78, <http://dx.doi.org/10.1111/j.1475-4983.2010.01002.x>.
- Varela, J.A., Parras, A., 2013. Análisis tafonómico de una concentración de vertebrados en la Formación La Colonia (Cretácico Tardío), Chubut, Argentina. *Ameghiniana* 50 (Suplemento) (74R–75R).
- Vasile, S., Csiki-Sava, Z., Venczel, M., 2013. A new madtsoiid snake from the Upper Cretaceous of the Hătege Basin, western Romania. *J. Vert. Paleontol.* 33, 1100–1119, <http://dx.doi.org/10.1080/02724634.2013.764882>.
- Vera, E.I., Pérez Loinaze, V.S., Llorens, M., Páez, M., Passalia, M.G., 2019. Fossil woods with coniferalean affinities from the Upper Cretaceous (Campanian–Maastrichtian) Puntudo Chico Formation, Chubut Province, Argentina. *Cretaceous Res.* 99, 321–333, <http://dx.doi.org/10.1016/j.cretres.2019.01.022>.
- Yi, H., Norell, M.A., 2015. The burrowing origin of modern snakes. *Sci. Adv.* 1, e1500743, <http://dx.doi.org/10.1126/sciadv.1500743>.
- Wiens, J.J., 2001. Character analysis in morphological phylogenetics: problems and solutions. *Syst. Biol.* 50, 689–699, <http://dx.doi.org/10.1080/106351501753328811>.
- Wilson, J.A., Mohabey, D.N., Peters, S.E., Head, J.J., 2010. Predation upon hatchling dinosaurs by a new snake from the Late Cretaceous of India. *Plos Biol.* 8, e1000322, <http://dx.doi.org/10.1371/journal.pbio.1000322>.