

A new species of possible
archipolypodan millipede
from the Carboniferous of the Netherlands
with unusually long tergites

Joachim T. HAUG, René H. B. FRAAIJE &
Carolyn HAUG

art. 22 (29) — Published on 20 October 2023
www.cr-palevol.fr



PUBLICATIONS
SCIENTIFIQUES



DIRECTEURS DE LA PUBLICATION / PUBLICATION DIRECTORS:

Bruno David, Président du Muséum national d'Histoire naturelle
Étienne Ghys, Secrétaire perpétuel de l'Académie des sciences

RÉDACTEURS EN CHEF / EDITORS-IN-CHIEF: Michel Laurin (CNRS), Philippe Taquet (Académie des sciences)

ASSISTANTE DE RÉDACTION / ASSISTANT EDITOR: Adenise Lopes (Académie des sciences; cr-palevol@academie-sciences.fr)

MISE EN PAGE / PAGE LAYOUT: Emmanuelle Rocklin (Muséum national d'Histoire naturelle; emmanuelle.rocklin@mnhn.fr)

RÉVISIONS LINGUISTIQUES DES TEXTES ANGLAIS / ENGLISH LANGUAGE REVISIONS: Kevin Padian (University of California at Berkeley)

RÉDACTEURS ASSOCIÉS / ASSOCIATE EDITORS (*, took charge of the editorial process of the article/a pris en charge le suivi éditorial de l'article):

Micropaléontologie/Micropalaeontology

Lorenzo Consorti (Institute of Marine Sciences, Italian National Research Council, Trieste)

Paléobotanique/Palaeobotany

Cyrille Prestianni (Royal Belgian Institute of Natural Sciences, Brussels)

Métazoaires/Metazoa

Annalisa Ferretti* (Università di Modena e Reggio Emilia, Modena)

Paléoichthyologie/Palaeoichthyology

Philippe Janvier (Muséum national d'Histoire naturelle, Académie des sciences, Paris)

Amniotes du Mésozoïque/Mesozoic amniotes

Hans-Dieter Sues (Smithsonian National Museum of Natural History, Washington)

Tortues/Turtles

Walter Joyce (Universität Freiburg, Switzerland)

Lépidosauromorphes/Lepidosauromorphs

Hussam Zaher (Universidade de São Paulo)

Oiseaux/Birds

Eric Buffetaut (CNRS, École Normale Supérieure, Paris)

Paléomammalogie (mammifères de moyenne et grande taille)/Palaeomammalogy (large and mid-sized mammals)

Lorenzo Rook (Università degli Studi di Firenze, Firenze)

Paléomammalogie (petits mammifères sauf Euarchontoglires)/Palaeomammalogy (small mammals except for Euarchontoglires)

Robert Asher (Cambridge University, Cambridge)

Paléomammalogie (Euarchontoglires)/Palaeomammalogy (Euarchontoglires)

K. Christopher Beard (University of Kansas, Lawrence)

Paléoanthropologie/Palaeoanthropology

Aurélien Mounier (CNRS/Muséum national d'Histoire naturelle, Paris)

Archéologie préhistorique (Paléolithique et Mésolithique)/Prehistoric archaeology (Palaeolithic and Mesolithic)

Nicolas Teyssandier (CNRS/Université de Toulouse, Toulouse)

Archéologie préhistorique (Néolithique et âge du bronze)/Prehistoric archaeology (Neolithic and Bronze Age)

Marc Vander Linden (Bournemouth University, Bournemouth)

RÉFÉRÉS / REVIEWERS: <https://sciencepress.mnhn.fr/fr/periodiques/comptes-rendus-palevol/referes-du-journal>

COUVERTURE / COVER:

New millipede *Lauravolsella willemeni* n. gen., n. sp., MAB13859A/B.

Comptes Rendus Palevol est indexé dans / *Comptes Rendus Palevol* is indexed by:

- Cambridge Scientific Abstracts
- Current Contents® Physical
- Chemical, and Earth Sciences®
- ISI Alerting Services®
- Geoabstracts, Geobase, Georef, Inspec, Pascal
- Science Citation Index®, Science Citation Index Expanded®
- Scopus®.

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Comptes Rendus Palevol* sont référencés par /
Articles and nomenclatural novelties published in Comptes Rendus Palevol are registered on:

- ZooBank® (<http://zoobank.org>)

Comptes Rendus Palevol est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris et l'Académie des sciences, Paris
Comptes Rendus Palevol is a fast track journal published by the Museum Science Press, Paris and the Académie des sciences, Paris

Les Publications scientifiques du Muséum publient aussi / The Museum Science Press also publish:

Adansonia, *Geodiversitas*, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Cryptogamie* sous-sections *Algologie*, *Bryologie*, *Mycologie*.

L'Académie des sciences publie aussi / The Académie des sciences also publishes:

Comptes Rendus Mathématique, *Comptes Rendus Physique*, *Comptes Rendus Mécanique*, *Comptes Rendus Chimie*, *Comptes Rendus Géoscience*, *Comptes Rendus Biologies*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <https://sciencepress.mnhn.fr>

Académie des sciences, Institut de France, 23 quai de Conti, 75006 Paris.

© This article is licensed under the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>)
ISSN (imprimé / print): 1631-0683/ ISSN (électronique / electronic): 1777-571X

A new species of possible archipolypodan millipede from the Carboniferous of the Netherlands with unusually long tergites

Joachim T. HAUG

Ludwig-Maximilians-Universität München (LMU), Faculty of Biology,
Großhaderner Straße 2, 82152 Planegg-Martinsried (Germany)
and GeoBio-Center at Ludwig-Maximilians-Universität München (LMU),
Richard-Wagner Straße 10, 80333 München (Germany)
joachim.haug@palaeo-evo-devo.info (corresponding author)

René H. B. FRAAIJE

Oertijdmuseum, Bosscheweg 80, 5283WB, Boxtel (The Netherlands)
directie@oertijdmuseum.nl

Carolyn HAUG

Ludwig-Maximilians-Universität München (LMU), Faculty of Biology,
Großhaderner Straße 2, 82152 Planegg-Martinsried (Germany)
and GeoBio-Center at Ludwig-Maximilians-Universität München (LMU),
Richard-Wagner Straße 10, 80333 München (Germany)
carolin.haug@palaeo-evo-devo.info

Submitted on 24 February 2022 | Accepted on 13 March 2023 | Published on 20 October 2023

[urn:lsid:zoobank.org:pub:0A87443C-65B6-496F-9E24-AB85BC06E4F3](https://zoobank.org/pub:0A87443C-65B6-496F-9E24-AB85BC06E4F3)

Haug J. T., Fraaije R. H. B. & Haug C. 2023. — A new species of possible archipolypodan millipede from the Carboniferous of the Netherlands with unusually long tergites. *Comptes Rendus Palevol* 22 (29): 595-604. <https://doi.org/10.5852/cr-palevol2023v22a29>

ABSTRACT

Millipedes have a long evolutionary history, with the oldest presumed fossils of Diplopoda de Blainville in Gervais, 1844 being from the Silurian and the first definite fossil record originating from Devonian deposits. The phylogeny of Diplopoda is not fully resolved yet, especially not concerning fossil representatives. At the same time, already in the Palaeozoic millipedes showed quite a morphological and presumably also ecological variety. We describe here a new species of a Carboniferous millipede from the Westphalian A of the Netherlands, *Lauravolsella willemeni* n. gen., n. sp., a possible representative of Archipolypoda (†Archipolypoda Scudder, 1882). The species is based on a single specimen, preserved with part and counterpart, which both show a three-dimensional preservation. The specimen has unusually long tergites, in normal life position covering most of the following segment. These long tergites might have been beneficial when performing defensive enrolling. In extant millipedes, enrolling is usually facilitated by softer areas between the sternites, allowing for a certain degree of ventral compression. In the new fossil, the sclerotic sternites occupy the entire length of the ventral side of the segment, not allowing for any type of compression. The new fossil therefore demonstrates another solution for the mechanical challenges during enrolment and increases the morphological diversity of Carboniferous millipedes.

KEY WORDS

Myriapoda,
Diplopoda,
Archipolypoda,
stereo imaging,
new genus,
new species.

RÉSUMÉ

Une nouvelle espèce de millipède archipolypode du Carbonifère des Pays-Bas avec des tergites inhabituellement longs.

Les millipèdes ont une longue histoire évolutive, les plus anciens fossiles présumés de Diplopoda de Blainville *in* Gervais, 1844 datant du Silurien et le premier fossile défini provenant de dépôts dévonien. La phylogénie des diplopodes n'est pas encore totalement résolue, en particulier en ce qui concerne les représentants fossiles. En même temps, dès le Paléozoïque, les mille-pattes présentaient une grande variété morphologique et probablement aussi écologique. Nous décrivons ici une nouvelle espèce de mille-pattes carbonifères du Westphalien A des Pays-Bas, *Lauravolsella willemeni* n. gen., n. sp., un représentant possible d'Archipolypoda (†Archipolypoda Scudder, 1882). L'espèce est basée sur un seul spécimen, préservé avec une partie et une contrepartie, qui montrent toutes deux une préservation tridimensionnelle. Le spécimen a des tergites inhabituellement longs, en position de vie normale couvrant la majeure partie du segment suivant. Ces longs tergites auraient pu être utiles lors de l'enroulement défensif. Chez les millipèdes existants, l'enroulement est généralement facilité par des zones plus souples entre les sternites, permettant un certain degré de compression ventrale. Dans le nouveau fossile, les sternites sclérosés occupent toute la longueur de la face ventrale du segment, ne permettant aucun type de compression. Le nouveau fossile démontre donc une autre solution pour les défis mécaniques lors de l'enroulement et augmente la diversité morphologique des millipèdes du Carbonifère.

MOTS CLÉS

Myriapoda,
Diplopoda,
Archipolypoda,
imagerie stéréo,
genre nouveau,
espèce nouvelle.

INTRODUCTION

Diplopoda de Blainville *in* Gervais, 1844, the group of millipedes, may not be as species-rich as beetles, flies or moths, but with more than 10 000 formally described species in the extant fauna, it is still one of the more diverse lineages in Euarthropoda. Furthermore, millipedes represent important components of the soil and leaf-litter fauna, by this fulfilling important eco-system functions.

The group Diplopoda has a long evolutionary history; the oldest fossils supposed to be millipedes are known from the Silurian (Almond & Lawson 1985; Wilson & Anderson 2004; Wilson 2005a; Shear & Edgecombe 2010: 183); definite representatives of the group are known from the Devonian (Shear 1998: 123). All older fossils of millipede-type appearance (e.g. Budd *et al.* 2001) could not be identified as representatives of Diplopoda more reliably. This also applies to some fossils that are contemporary with true millipedes (Tesakov & Alekseev 1992; Edgecombe 2015a).

The phylogeny of Diplopoda has not been well reconstructed so far, many conflicting hypotheses have been put forward (Shear & Edgecombe 2010: 179, 180; Edgecombe 2015b). Furthermore, most of the phylogenetic reconstructions include only extant representatives. Interpreting fossils in this frame remains still very challenging.

This challenge is especially true for Palaeozoic representatives. Most of the relatively numerous Palaeozoic specimens have been interpreted as representatives of the group Archipolypoda (†Archipolypoda Scudder, 1882). Representatives of Archipolypoda, together with their contemporary relatives, seem to have played an important role in the past, being central components of the Palaeozoic terrestrial fauna. We know quite some details about the biology of some of these animals based on rather complete specimens (e.g. Wilson 2005a, b,

2006a, b; Wilson & Hannibal 2005; Wilson *et al.* 2005); others remain known only from rather fragmentary pieces (e.g. Shear 1993; Hannibal & Krzemiński 2005; Hannibal & May 2021).

Some of the more complete Palaeozoic millipedes, for example, are known to have been able to fully enrol into a tightly closed ball armed with spines, supposedly for defensive purposes (Hannibal & Feldmann 1981; Racheboeuf *et al.* 2004). While such a morphology and behaviour are well known in the extant fauna (e.g. pill bugs, false sand crab larvae, mantis shrimp larvae), this behaviour has likely evolved independently in these Palaeozoic forms, like in other lineages of Euarthropoda (Haug & Haug 2014; Rudolf *et al.* 2016).

While it is well known that many extant millipedes have specialised appendages, most Palaeozoic ones have rather uniform appendages. Still some seem to have possessed already differentiated appendages on the trunk (Hannibal 1995).

Even some interaction between individuals is known for Palaeozoic millipedes. Several small-sized specimens have been preserved in close proximity, indicating an aggregation behaviour of these animals (Wilson 2006a). As the different finds demonstrate, millipedes in the later Palaeozoic already show quite a variety, from a number of deposits from all over the world.

Studies on fossils of Euarthropoda from the Carboniferous of the Netherlands are so far scarce. Pruvost (1927) published an isolated Namurian insect wing from a borehole near Gulpen, southern Limburg, the Netherlands. Laurentiaux (1950) described seven more winged insects from diverse coal mines of southern Limburg. Another new insect wing from the Westphalian A of the “Laura” mine was documented by Brauckmann & Gröning (1996). Laurentiaux-Vieira & Laurentiaux (1961) described a new whip scorpion (Thelyphonida Latreille, 1804; Arachnida Cuvier, 1812), whereas Steur &

Van Houtem (2012) reported on a representative of Trigonotarbida (Arachnida), both from the Westphalian A. In the study of Van der Heide (1951) all then known Carboniferous fossils of Euarthropoda (excluding scorpions and insects) of southern Limburg were listed, including many fragments of *Arthropleura* (†*Arthropleura* Meyer, 1854 – a presumed representative of Myriapoda Latreille, 1802) and an indefinite remain of another possible representative of Myriapoda.

We here report a new fossil millipede from the Carboniferous of the Netherlands. The specimen was collected in 2011 on the Westphalian A dumps of the former “Laura” coal mine in Eygelshoven, southern Limburg.

MATERIAL AND METHODS

MATERIAL

In the focus of the study is a single specimen, MAB13859A/B. It is preserved as part and counterpart. The specimen was found in 2011 by a private collector (Mr. Geert Willemsen, Middelbeers, the Netherlands) in the coal mine dump of mine “Laura” near Eygelshoven in Southern Limburg, the Netherlands. Accordingly, the fossil is of lowermost Westphalian age (about 313–304 million years before present), being part of the upper Carboniferous and Pennsylvanian. The specimen is deposited in the Oertijlmuseum, Boxtel, the Netherlands.

For comparison, a millipede from the Pleistocene was documented, ROMIP32380. The label of the specimen stated that it was determined as *Fontaria corrugata* (for a discussion on the validity of this name, see Hoffman 1957). It was found in Whitchurch Township, York County, Ontario. The specimen is deposited at the Royal Ontario Museum, Toronto, Canada.

DOCUMENTATION METHODS

Many Carboniferous fossils of the group Euarthropoda are challenging to be documented with minimum artefacts. A major challenge is the proper differentiation between the information provided by the relief and the colour (Haug *et al.* 2012a, 2015; Hörnig *et al.* 2018). The standard method of using oblique light is not well suited for such an approach. Even, shadow-free lighting provides a good access to the colour information, while stereo imaging provides an independent access to the relief. Stereo images can be produced directly from a specimen by recording two images from two slightly differing angles (Haug *et al.* 2012a, b, 2015; Hörnig *et al.* 2018). It is often necessary for stereo images to document the entire specimen with one image instead of applying composite imaging (Haug *et al.* 2011a), as stitching (merging) of stereo images to a panorama can lead to severe artefacts in depth perception.

The specimen at hand is very elongate, documenting it with a single image for a stereo image would have led to a rather low resolution. Yet, there are alternatives to physical stereo images. Instead, it is possible to extract a surface information of a specimen by depth from defocus (Haug *et al.* 2011b, 2012b, 2013a, b) and produce stereo images from the virtual surfaces resulting from these.

Part and counterpart of the specimen were documented as a composite image (Haug *et al.* 2011a) on a Keyence VHX-6000 digital microscope, under cross-polarised light and unpolarised ring light (Haug *et al.* 2019). Several exposure times were combined (HDR, cf. Haug *et al.* 2013c). Based on depth from defocus, a surface was calculated. Of this, stereo images were produced.

The specimen for comparison was documented as a physical stereo image. The camera used was a Canon EOS Rebel T3i with a MP-E 65 mm lens.

SYSTEMATICS

Myriapoda Latreille, 1802
Diplopoda de Blainville *in* Gervais, 1844
†Archipolypoda Scudder, 1882

Lauravolsella n. gen.

[urn:lsid:zoobank.org:act:BB3144B2-66D7-411E-BB72-1659B1D8ED2D](https://zoobank.org/act:BB3144B2-66D7-411E-BB72-1659B1D8ED2D)

DERIVATION OF NAME. — A combination of Laura, the name of the former colliery at Eygelshoven, and the Latin word for forceps alluding to the shape of the anal region.

TYPE SPECIES. — *Lauravolsella willemeni* n. gen., n. sp.

DIAGNOSIS. — As for the type species as it is monotypic.

Lauravolsella willemeni n. gen., n. sp. (Figs 1; 2; 4)

[urn:lsid:zoobank.org:act:6554B9A8-E52E-45FA-BD7F-F9F60F67DA68](https://zoobank.org/act:6554B9A8-E52E-45FA-BD7F-F9F60F67DA68)

DERIVATION OF NAME. — After the surname of the collector.

HOLOTYPE. — MAB13859A/B.

TYPE LOCALITY AND AGE. — Coal mine dump of mine “Laura” near Eygelshoven in Southern Limburg, the Netherlands, lowermost Westphalian (Westphalian A).

DIAGNOSIS. — Elongate millipede with 20 diplosegments. Each diplosegment with a posteriorly drawn out diplotergites, three times the length of the ventral region of each diplosegment.

DESCRIPTION

Elongate specimen, longer than wide, at least 4.5 times, total length *c.* 40 mm (Figs 1; 2). Mostly ventral details accessible. Subdivision into discrete units apparent: one anterior region, a series of 20 sub-similar units, and a terminal end. The segmented appearance of the body indicates that the specimen is a representative of Euarthropoda; representatives of this group have a body usually differentiated into an anterior head and a posterior trunk.

Anterior region not well preserved with many details, therefore remains partly unclear. It is wider than long. Possibly this



FIG. 1. — New millipede *Lauravolsella willemeni* n. gen., n. sp., MAB13859A/B: **A, B**, part B; **A**, cross-polarised light; **B**, reflected light under ring illumination; **C**, part A, reflected light under ring illumination. Scale bars: 3 mm.

region represents the original head region and some trunk segments. The anterior edge of this region is gently rounded, further posterior elevations and depressions may represent remains of original mouthparts. Alternatively, the elevated region (Fig. 2C marked green) may represent the not well preserved head, and the depression (Fig. 2C marked cyan) may represent a not well delineated trunk segment. At least one pair of insertion areas of presumably locomotory (ambulatory) appendages is apparent at the posterior part of the anterior region. These are most likely arising from a trunk segment, not well delineated from the rest of the anterior region.

All of the following 20 units are subsimilar in appearance (Figs 1; 2A-C). Each unit is wider than long, about four times. Medially a sternal sclerite is apparent, occupying about 20% of the area. The sternal sclerite is deeply subdivided into two equal parts, an anterior one and a posterior one, by a distinct groove (Fig. 2D). Along the groove, four evenly spaced pores apparent. Adjacent to the sternal sclerites insertion areas of appendages or proximal parts of appendages apparent, two pairs per unit. These insertions/proximal parts are of similar dimensions as an anterior/posterior part of a sternal sclerite. Further later-




FIG. 2. — New millipede *Lauravolsella willemeni* n. gen., n. sp., MAB13859A/B: **A**, stereo image of part A; **B**, stereo image of part B; **C**, colour-marked version of B; **D**, **E**, stereo images of details of part B; **D**, trunk region showing the leg insertions and sternal sclerites; **E**, terminal end. Abbreviations: **d1-d20**, diplosegments 1-20; **te**, trunk end. Scale bars: A-C, 3 mm; D, E, 2 mm.

ally smooth region. Far lateral parts of the dorsal region of the unit visible in some of the units. More elongated than ventral region, up to three times.

Further posterior units slightly more slender. Very posterior ones slightly backwards curved, partly surrounding the posterior end.

Terminal end drop-shaped with tip facing backwards, slightly longer than wide (Fig. 2E). Three distinct sclerites apparent; antero-median sclerite small circular; other sclerites elongate left and right (symmetrically similar), anteriorly surrounding

the small sclerite, posteriorly leaving an elongate triangular gap between them.

DISCUSSION

THE NEW FOSSIL: A REPRESENTATIVE OF THE LINEAGE OF DIPLOPODA

The fossil is unfortunately incompletely preserved in several aspects. Only the very proximal parts of the appendages are




FIG. 3. — Schematic representations of the proximal leg arrangement (upper rows, proximal leg element in **darker grey**, sternite in **lighter grey**) and anal region (lower rows): **A**, new fossil; **B**, Polydesmida Leach, 1815 (based on Gilgado *et al.* 2015: fig. 10D); **C**, Polyxenida Lucas, 1840 (based on Huynh & Veenstra 2018: fig. 6B); **D**, Spirostreptida Brandt, 1833 (based on Likhitrakarn *et al.* 2018: fig. 5O, U); **E**, Siphonophorida Newport, 1844 (based on Marek *et al.* 2012: fig. 23); **F**, Julida Leach, 1814 (Reboleira & Enghoff 2018: fig. 3C); **G**, Callipodida Pocock, 1894 (based on Liu & Tian 2015: fig. 5); **H**, Platydesmida Cook, 1895 (based on Shorter *et al.* 2018: fig. 5D); **I**, Stemmiulida Pocock, 1894 (based on Mauriès *et al.* 2010: fig. 7); **J**, Glomerida Leach, 1814 (based on Antić & Makarov 2021: fig. 1C).

accessible, and especially the anterior body region does not provide many details. Yet, the proximal leg region as well as the terminal end is already quite informative.

As described above, two pairs of appendages correspond to a single tergite in the here described fossil. This arrangement is a diplosegment in the wider sense, as for example known in Pauropoda. In Diplopoda, the diplosegments are further derived into a single structure (diplosegments *sensu stricto*). While diplosegments are known in different lineages of Euarthropoda (e.g. in Scutigermorpha or in certain fossil representatives, e.g. †*Sinoburius lunaris* Hou, Ramsköld & Bergström, 1991; Schmidt *et al.* 2021), diplosegments along (almost) the entire body occur only in Pauropoda and Diplopoda (e.g. Sierwald & Bond 2007; Janssen 2011; Koch 2015). Unfortunately, the anterior body region of the new fossil is not well preserved; in millipedes, we should expect three single segments instead of diplosegments after the head region. While these are not apparent in the fossil, the preservation well explains this lack.

A further structural similarity of the fossil to extant representatives of Diplopoda is the arrangement of the anal region. There are two prominent anal vales (“paraprocts”) and an anterior smaller sclerite (“ventral scale”, “hypoproct”) delimiting the anal opening. Such an arrangement is well apparent in the fossil. It seems that in other lineages of Myriapoda such an arrangement is not developed. Especially in Pauropoda,

no such prominent arrangement of sclerites around the anus is present (Tiegs 1947: 278, text fig. 24A; Scheller 1988: 12, fig. 6; Bu 2020: 56, fig. 1G).

All these observations support the interpretation of the new fossil as a representative of Diplopoda or at least as a derivative of the lineage towards the group (“stem-lineage”). As pointed out, the majority of Carboniferous millipede fossils have so far been interpreted as representatives of the group Archipolypoda. Archipolypoda is characterised by the presence of paired pores on the sternites, often with valve-like structures closing these. Also the new fossil has distinct openings in the sternites, making it likely that the fossil is also a representative of Archipolypoda. Still, it has some morphological details that appear quite unique among the Palaeozoic fossils, such as the very long posteriorly elongated tergites. In addition, it possesses features shared with different extant ingroups of Diplopoda. We briefly highlight some of these.

CHARACTER DISTRIBUTION WITHIN DIPLOPODA

Two aspects are very apparent in the new fossil (Fig. 3A):

- the sclerite anterior to the anus (“ventral scale”, “hypoproct”) is rather small, more or less circular;
- the proximal leg elements are in quite some distance to each other, between them is a distinct sternite.




FIG. 4. — Schematic restorations of new millipede *Lauravolsella willemeni* n. gen., n. sp. and comparison: **A**, restoration in lateral view; **B**, restoration in dorsal view; **C**, supposed enrolment posture; **D**, partly enrolled millipede from the Pleistocene for comparison of posture, red-cyan stereo image of ROMIP32380. Scale bar: 2 mm.

Among extant ingroups of Diplopoda, a small sclerite anterior to the anus occurs, for example in representatives of Callipodida Pocock, 1894 (Liu & Tian 2015: 126, fig. 5) (Fig. 3G) and Stemmiulida Pocock, 1894 (Mauriès *et al.* 2010: 75, fig. 7) (Fig. 3I). In other ingroups of Diplopoda, the sclerite anterior to the anus is either not apparent, as in representatives Polyxenida Lucas, 1840 (Huynh & Veenstra 2018: 143, fig. 6B) (Fig. 3C) or Platydesmida Cook, 1895 (Shorter *et al.* 2018: 30, fig. 5D) (Fig. 3H), or the sclerite is quite broad as in representatives of Polydesmida Leach, 1815 (Gilgado *et al.* 2015: 27, fig. 10D) (Fig. 3B), Spirostreptida Brandt, 1833 (Likhitrakarn *et al.* 2018: 65, fig. 5U) (Fig. 3D), Julida Leach, 1814 (Reboleira & Eng-hoff 2018: 97, fig. 3C) (Fig. 3F), or Chordeumatida Pocock, 1894 (Reboleira *et al.* 2018: 4, fig. 1D).

In most ingroups of Diplopoda, the insertions of the left and right legs are very close to each other, with almost no distance between them and often without a distinct sternite or a rather concealed one as in representatives of Callipodida (Liu & Tian 2015: 126, fig. 5) (Fig. 3G), Glomerida Leach, 1814 (Antić & Makarov 2021: 396, fig. 1C) (Fig. 3J), Spiro-streptida (Likhitrakarn *et al.* 2018: 65, fig. 5O) (Fig. 3D), or Spirobolida Bollman, 1893 (Wesener *et al.* 2009: 31, fig. 16A).

If a more prominent sternite is present in cases in which the legs are close to each other, it may be positioned anterior to the legs as in representatives of Polyxenida (Huynh & Veenstra 2018: fig. 6B) (Fig. 3C), Siphonophorida Newport, 1844 (Marek *et al.* 2012: 94, fig. 23) (Fig. 3E), or Chordeumatida (Reboleira *et al.* 2018: 4, fig. 1B). Alternatively, it may be positioned more posteriorly, “squeezed” in between from posterior as in representatives of Platydesmida (Shorter *et al.* 2018: 30, fig. 5D) (Fig. 3H). A broader sternite region between leg insertions, as in the fossil, is present in representatives of Polydesmida (Golovatch *et al.* 2013: 38, fig. 14G; Gilgado *et al.* 2015: 27, fig. 10D) (Fig. 3B).

In other Palaeozoic millipedes, prominent sternite regions seem to be present, yet the details of the leg insertions remain unclear for some specimens (e.g. Almond & Lawson 1985: 231, fig. 2). In some known representatives of Archipolypoda, the sclerite anterior to the anus is quite broad (Wilson 2005b: 1099, text fig. 3), unlike in the new fossil. Similar to the new fossil, in other representatives of Archipolypoda the region between the leg insertion seems very wide (Wilson 2005b: 1099, text fig. 2). Different from the new fossil, it appears separated into two halves by a distinct median groove.

Overall, the new fossil shows a mixture of features characteristic for Archipolypoda, of specialisations of its own, but also of some features seen in different extant millipedes. Apparently, convergent evolution seems to have been common within millipedes.

FUNCTIONAL ASPECTS OF THE NEW FOSSIL

The strongly posteriorly drawn out (diplo-)tergites of the diplosegments appear quite special on a first glance. Yet, it seems that there is indeed a functional explanation for these structures.

Some millipedes can perform a fully enclosed enrolment (e.g. Glomerida: Antić & Makarov 2021: 398, fig. 3; 405, fig. 9H; Sphaerotheriida: Wesener & Sierwald 2005: 597, fig. 63; Wesener & Koenig 2016: 19, fig. 11C, E-H; 20, fig. 12A, C-E, G; 21, fig. 13A; 22, fig. 14A, B, F-H; some forms of Polydesmida: Golovatch 2003: 48, fig. 24) likely for defensive purposes. Also many millipedes that cannot fully enroll perform a less tightly closed spiral type of enrolment (e.g. Sierwald & Bond 2007: 404, fig. 2C, I; Wesener *et al.* 2009: 32, fig. 17; 67, fig. 37B, C; Shorter *et al.* 2018: 25, fig. 1D).

Enrolling is not a simple task: the dorsal side of the body needs to be stretched, the ventral side to be compressed. For the latter there is often a distance between the sternites with softer membrane (e.g. Golovatch 2003: 56, fig. 46; Golovatch *et al.* 2013: 58, fig. 30H).

In the new fossil, there is apparently no distance between the sternal sclerites, no compression is possible here. Enrolling would be limited to the dorsal stretching. Strong stretching would expose soft membranous areas, if the diplo-tergites would be of a normal length. The strongly elongated diplo-tergites hence will most likely provide protection when enrolling.

For estimating how such a posture could look like we need to assume certain aspects of the body which are not directly accessible: the height of the body was assumed as being equal to the width of the appendage insertions (Fig. 4A), as many enrolling millipedes indeed have a body that is relatively high. The exact shape of the posterior edge of the diplo-tergite was assumed to be slightly concave (Fig. 4B). We assumed maximum tilting for adjacent segments until the anterior edge of the posterior segment was still covered by the drawn out diplo-tergite of the anterior segment. Based on these assumptions the animal could have enrolled quite effectively (Fig. 4C *vs.* 4D). We also assumed the presence of a collum which allows to fold the head further inwards. It is indeed possible that part of the anterior region of the fossil represents the collum.

Overall, it seems quite likely that the elongated diplo-tergites served for protection in conjunction with an enrolment behaviour. Yet, no real extant counterpart of such an arrangement is known.

Acknowledgements

We thank Greg Edgecombe, London, and Joseph T. Hannibal, Cleveland, for sharing their thoughts on the fossil, Hans Steur, Ellecom, the Netherlands, for items of literature, and

Geert Willems, Middelbeers, the Netherlands, for donating the studied specimen. We are grateful to Alessandro Minelli, Padova, and an anonymous reviewer for helpful comments. JTH is kindly supported by the Volkswagen Foundation in the course of a Lichtenberg professorship. J. Matthias Starck, Munich, is thanked for long-term support. We thank all people providing low-cost, open-access, and open-source software.

REFERENCES

- ALMOND J. E. & LAWSON J. D. 1985. — The Silurian-Devonian fossil record of the Myriapoda. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* 309 (1138): 227–237. <https://doi.org/10.1098/rstb.1985.0082>
- ANTIĆ D. Ž. & MAKAROV S. E. 2021. — *Macedomeris*, a new monotypic doderiinine genus (Diplopoda, Glomerida) from a cave in North Macedonia, Balkan Peninsula. *Zootaxa* 4908 (3): 393–408. <https://doi.org/10.11646/zootaxa.4908.3.4>
- BRAUCKMANN C. & GRÖNING E. 1996. — Ein neuer ober-karbonischer Insekten-Flügel aus Süd-Limburg (Palaeodictyoptera: Breyeriidae, Westfalium, Niederlande). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 2: 72–82. <https://doi.org/10.1127/njgpm/1996/1996/72>
- BU Y. 2020. — First record of the family Colinauropodidae (Myriapoda, Pauropoda) in China, with the description of three new species. *ZooKeys* 947: 53–70. <https://doi.org/10.3897/zookeys.947.53723>
- BUDD G. E., HÖGSTRÖM A. E. & GOGIN I. 2001. — A myriapod-like arthropod from the Upper Cambrian of East Siberia. *PalZ* 75: 37–41. <https://doi.org/10.1007/BF03022596>
- EDGECOMBE G. D. 2015a. — Diplopoda — fossils, in MINELLI A. (ed.), *Treatise on Zoology – Anatomy, Taxonomy, Biology. The Myriapoda, Vol. 2*. Brill, Leiden: 337–351. https://doi.org/10.1163/9789004188273_015
- EDGECOMBE G. D. 2015b. — Diplopoda — phylogenetic relationship, in MINELLI A. (ed.), *Treatise on Zoology – Anatomy, Taxonomy, Biology. The Myriapoda, Vol. 2*. Brill, Leiden: 353–362. https://doi.org/10.1163/9789004188273_016
- GILGADO J. D., ENGHOF H., TINAUT A. & ORTUNO V. M. 2015. — Hidden biodiversity in the Iberian Mesovoid Shallow Substratum (MSS): new and poorly known species of the millipede genus *Archipolydesmus* Attems, 1898 (Diplopoda, Polydesmidae). *Zoologischer Anzeiger – A Journal of Comparative Zoology* 258: 13–38. <https://doi.org/10.1016/j.jcz.2015.06.001>
- GOLOVATCH S. I. 2003. — A review of the volvatory Polydesmida, with special reference to the patterns of volvation (Diplopoda). *African Invertebrates* 44 (1): 39–60.
- GOLOVATCH S. I., GEOFFROY J. J., STOEY P. & VAN DEN SPIEGEL D. 2013. — Review of the millipede family Opisotretidae (Diplopoda, Polydesmida), with descriptions of new species. *ZooKeys* 302: 13–77. <https://doi.org/10.3897/zookeys.302.5357>
- HANNIBAL J. T. 1995. — Modified legs (clasping appendages?) of Carboniferous euphoberiid millipeds (Diplopoda: Euphoberiida). *Journal of Paleontology* 69 (5): 932–938. <https://doi.org/10.1017/S0022336000035587>
- HANNIBAL J. T. & FELDMANN R. M. 1981. — Systematics and functional morphology of oniscomorph millipedes (Arthropoda: Diplopoda) from the Carboniferous of North America. *Journal of Paleontology* 55: 730–746.
- HANNIBAL J. T. & KRZEMIŃSKI W. 2005. — A palaeosomatid millipede (Archipolypoda: Palaeosomatida) from the Carboniferous (Namurian A) of Silesia, Poland. *Polskie pismo entomologiczne* 74: 205–217.
- HANNIBAL J. T. & MAY W. J. 2021. — Permian millipedes from the Fort Sill fissures of southwestern Oklahoma, with comments

- on allied taxa and millipedes preserved in karstic environments. *Journal of Paleontology* 95 (3): 586-600. <https://doi.org/10.1017/jpa.2020.100>
- HAUG C. & HAUG J. T. 2014. — Defensive enrolment in mantis shrimp larvae (Malacostraca: Stomatopoda). *Contributions to Zoology* 83 (3): 185-194. <https://doi.org/10.1163/18759866-08303003>
- HAUG J. T., HAUG C., KUTSCHERA V., MAYER G., MAAS A., LIEBAU S., CASTELLANI C., WOLFRAM U., CLARKSON E. N. K. & WALOSZEK D. 2011a. — Autofluorescence imaging, an excellent tool for comparative morphology. *Journal of Microscopy* 244 (3): 259-272. <https://doi.org/10.1111/j.1365-2818.2011.03534.x>
- HAUG C., MAYER, G., KUTSCHERA V., WALOSZEK D., MAAS A. & HAUG J. T. 2011b. — Imaging and documenting gammarideans, in GERHARDT A., BLOOR M. & MILLS C. L. (eds), *Gammarus*: important taxon in freshwater and marine changing environments. *International Journal of Zoology* 2011: 380829. <https://doi.org/10.1155/2011/380829>
- HAUG C., VAN ROY P., LEIPNER A., FUNCH P., RUDKIN D. M., SCHÖLLMANN L. & HAUG J. T. 2012a. — A holomorph approach to xiphosuran evolution – a case study on the ontogeny of *Euproops*. *Development Genes & Evolution* 222: 253-268. <https://doi.org/10.1007/s00427-012-0407-7>
- HAUG J. T., MAYER G., HAUG C. & BRIGGS D. E. G. 2012b. — A Carboniferous non-onychophoran lobopodian reveals long-term survival of a Cambrian morphotype. *Current Biology* 22 (18): 1673-1675. <https://doi.org/10.1016/j.cub.2012.06.066>
- HAUG J. T., MÜLLER C. H. G. & SOMBKE A. 2013a. — A centipede nymph in Baltic amber and a new approach to document amber fossils. *Organisms Diversity & Evolution* 13: 425-432. <https://doi.org/10.1007/s13127-013-0129-3>
- HAUG C., NYBORG T. & VEGA F. J. 2013b. — An exceptionally preserved upogebiid (Decapoda: Reptantia) from the Eocene of California, in VEGA F. J. & GARASSINO A. (eds), *Mesozoic and Cenozoic Crustacea: updating their systematics*. *Boletín de la Sociedad Geológica Mexicana* 65 (2): 235-248. <https://doi.org/10.18268/BSGM2013v65n2a5>
- HAUG C., SHANNON K. R., NYBORG T. & VEGA F. J. 2013c. — Isolated mantis shrimp dactyli from the Pliocene of North Carolina and their bearing on the history of Stomatopoda, in VEGA F. J. & GARASSINO A. (eds), *Mesozoic and Cenozoic Crustacea: updating their systematics*. *Boletín de la Sociedad Geológica Mexicana* 65 (2): 273-284. <https://www.jstor.org/stable/24921224>
- HAUG J. T., LABANDEIRA C. C., SANTIAGO-BLAY J. A., HAUG C. & BROWN S. 2015. — Life habits, hox genes, and affinities of a 311 million-year-old holometabolous larva. *BMC Evolutionary Biology* 15 (208). <https://doi.org/10.1186/s12862-015-0428-8>
- HAUG J. T., MÜLLER P. & HAUG C. 2019. — A 100-million-year old predator: a fossil neuropteran larva with unusually elongated mouthparts. *Zoological Letters* 5 (29). <https://doi.org/10.1186/s40851-019-0144-0>
- HOFFMAN R. L. 1957. — The status of *Fontaria coriacea* Koch and of *Polydesmus corrugatus* Wood: A most regrettable tangle of names in the Diplopoda (Polydesmida: Xystodesmidae). *Proceedings of the Biological Society of Washington* 70: 183-188.
- HÖRNIG M. K., HAUG C., SCHNEIDER J. W. & HAUG J. T. 2018. — Evolution of reproductive strategies in dictyopteran insects – clues from ovipositor morphology of extinct roachoids. *Acta Palaeontologica Polonica* 63 (1): 1-24. <https://doi.org/10.4202/app.00324.2016>
- HUYNH C. & VEENSTRA A. A. 2018. — Two new *Lophoturus* species (Diplopoda, Polyxenida, Lophoproctidae) from Queensland, Australia. *ZooKeys* 741: 133-154. <https://doi.org/10.3897/zookeys.741.21814>
- JANSSEN R. 2011. — Diplosegmentation in the pill millipede *Glomeris marginata* is the result of dorsal fusion. *Evolution & Development* 13 (5): 477-487. <https://doi.org/10.1111/j.1525-142x.2011.00504.x>
- KOCH M. 2015. — Diplopoda – general morphology, in MINELLI A. (ed.), *Treatise on Zoology-Anatomy, Taxonomy, Biology. The Myriapoda*, Vol. 2. Brill, Leiden: 7-67. https://doi.org/10.1163/9789004188273_003
- LAURENTIAUX D. 1950. — Les insectes houillers du Limbourg hollandais. *Mededelingen van de Geologische Stichting Serie C4*: 13-22.
- LAURENTIAUX-VIEIRA F. & LAURENTIAUX D. 1961. — *Prothelyphonus neerlandicus* nov. sp., Uropyge du Westphalien du Limbourg Hollandais. *Mededelingen van de Geologische Stichting* 13: 29-34.
- LIKHITRAKARN N., GOLOVATCH S. I., SRISONCHAI R., BREHIER F., LIN A., SUTCHARIT C. & PANHA S. 2018. — Two new species of the millipede family Cambalopsidae from Myanmar (Diplopoda, Spirostreptida). *ZooKeys* 760: 55-71. <https://doi.org/10.3897/zookeys.760.24837>
- LIU W. & TIAN M. 2015. — Two new cave-dwelling species of the millipede genus *Paracortina* Wang & Zhang, 1993 from southern China (Diplopoda, Callipodida, Paracortinidae). *ZooKeys* 517: 123-140. <https://doi.org/10.3897/zookeys.517.9949>
- MAREK P. E., SHEAR W. A. & BOND J. E. 2012. — A redescription of the leggiest animal, the millipede *Illacme plenipes*, with notes on its natural history and biogeography (Diplopoda, Siphonophorida, Siphonorhinidae). *ZooKeys* 241: 77-112. <https://doi.org/10.3897/zookeys.241.3831>
- MAURIÈS J. P., GOLOVATCH S. I. & GEOFFROY J. J. 2010. — Un nouveau genre et une nouvelle espèce de l'ordre Stemmiulida du Viet-Nam (Diplopoda) [A new genus and species of the order Stemmiulida from Vietnam (Diplopoda)]. *Arthropoda Selecta* 19 (2): 73-80. <https://doi.org/10.15298/arthsel.19.2.03>
- PRUVOST P. 1927. — Sur une aile d'insecte fossile trouvée au sondage de Gulpen. *Jaarverslag Geologisch Bureau voor het nederlandse Mijngedied te Heerlen* 1926: 76-77.
- RACHEBOEUF P. R., HANNIBAL J. T. & VANNIER J. 2004. — A new species of the diplopod *Amyiniylspes* (Oniscomorpha) from the Stephanian Lagerstätte of Montceau-les-Mines, France. *Journal of Paleontology* 78 (1): 221-229. <https://tinyurl.com/3j2srbd7>
- REBOLEIRA A. S. P. & ENGHOFF H. 2018. — First continental troglobiont *Cylindroiulus* millipede (Diplopoda, Julida, Julidae). *ZooKeys* 795: 93-103. <https://tinyurl.com/43sdjhsx>
- REBOLEIRA A. S. P., ENGHOFF H. & SANTAMARIA S. 2018. — Novelty upon novelty visualized by rotational scanning electron micrographs (rSEM): Laboulbeniales on the millipede order Chordeumatida. *PLoS ONE* 13: e0206900. <https://doi.org/10.1371/journal.pone.0206900>
- RUDOLF N. R., HAUG C. & HAUG J. T. 2016. — Functional morphology of giant mole crab larvae: a possible case of defensive enrollment. *Zoological Letters* 2 (17). <https://doi.org/10.1186/s40851-016-0052-5>
- SCHELLER U. 1988. — *The Pauropoda (Myriapoda) of the Savannah River Plant, Aiken, South Carolina*. Savannah River Plant National Environmental Research Park Program (Report no. SRO-NERP-17), Athens GA: 1-99. <https://doi.org/10.2172/5034022>
- SCHMIDT M., LIU Y., HOU X., HAUG J. T., HAUG C., MAI H. & MELZER R. R. 2021. — Intraspecific variation in the Cambrian: new observations on the morphology of the Chengjiang euarthropod *Sinoburius lunaris*. *BMC Ecology and Evolution* 21 (127). <https://doi.org/10.1186/s12862-021-01854-1>
- SHEAR W. A. 1993. — Myriapodous arthropods from the Viséan of East Kirkton, West Lothian, Scotland. *Earth and Environmental Science Transactions of The Royal Society of Edinburgh* 84 (3-4): 309-316. <https://doi.org/10.1017/S026359330000612X>
- SHEAR W. A. 1998. — The fossil record and evolution of the Myriapoda, in FORTEY R. A. & THOMAS R. H. (eds.), *Arthropod Relationships*. Chapman & Hall (The Systematics Association Special Volume Series; 55), London: 211-219.
- SHEAR W. A. & EDGECOMBE G. D. 2010. — The geological record and phylogeny of the Myriapoda. *Arthropod Structure & Development* 39 (2-3): 174-190. <https://doi.org/10.1016/j.asd.2009.11.002>

- SHORTER P. L., HENNEN D. A. & MAREK P. E. 2018. — Cryptic diversity in *Andrognathus corticarius* Cope, 1869 and description of a new *Andrognathus* species from New Mexico (Diplopoda, Platydesmida, Andrognathidae). *ZooKeys* 786: 19–41. <https://doi.org/10.3897/zookeys.786.27631>
- SIERWALD P. & BOND J. E. 2007. — Current status of the myriapod class Diplopoda (millipedes): taxonomic diversity and phylogeny. *Annual Review of Entomology* 52: 401–420. <https://doi.org/10.1146/annurev.ento.52.111805.090210>
- STEUR H. & VAN HOUTEM T. 2012. — *Eophrynus*, een spinachtige uit het Carboon van Zuid-Limburg. *Grondboor and Hamer* 66 (6): 444–447.
- TESAKOV A. S. & ALEKSEEV A. S. 1992. — Myriapod-like arthropods from the Lower Devonian of central Kazakhstan. *Paleontological Journal* 26: 18–23.
- TIEGS O. W. 1947. — The development and affinities of the Pauropoda, based on a study of *Pauropus silvaticus*. *Journal of Cell Science* 88 (3): 275–336. <https://doi.org/10.1242/jcs.s3-88.3.275>
- VAN DER HEIDE S. 1951. — Les Arthropodes du terrain houiller du Limbourg méridional (excepté les Scorpions et les insectes). *Mededelingen van de Geologische Stichting*, series C IV: 3–5.
- WESENER T. & KOENIG Z. F. A. 2016. — The giant pill-millipedes, order Sphaerotheriida: an annotated species catalogue with morphological atlas and list of apomorphies (Arthropoda: Diplopoda). *Bonn Zoological Bulletin Suppl.* 63: 1–104.
- WESENER T. & SIERWALD P. 2005. — The giant pill-millipedes of Madagascar. *Proceedings of California Academy of Science* 56: 557–599.
- WESENER T., ENGHOFF H. & SIERWALD P. 2009. — Review of the Spirobolida on Madagascar, with descriptions of twelve new genera, including three ‘fire millipedes’ (Diplopoda). *ZooKeys* 19: 1–128. <https://doi.org/10.3897/zookeys.19.221>
- WILSON H. M. 2005a. — Zosterogrammida, a new order of millipedes from the Middle Silurian of Scotland and the Upper Carboniferous of Euramerica. *Palaeontology* 48 (5): 1101–1110. <https://doi.org/10.1111/j.1475-4983.2005.00498.x>
- WILSON H. M. 2005b. — A new genus of archipolypodan millipede from the Coseley Lagerstätte, Upper Carboniferous, UK. *Palaeontology* 48 (5): 1097–1100. <https://doi.org/10.1111/j.1475-4983.2005.00496.x>
- WILSON H. M. 2006a. — Aggregation behaviour in juvenile millipedes from the Upper Carboniferous of Mazon Creek, Illinois. *Palaeontology* 49 (4): 733–740. <https://doi.org/10.1111/j.1475-4983.2006.00567.x>
- WILSON H. M. 2006b. — Juliformian millipedes from the Lower Devonian of Euramerica: implications for the timing of millipede cladogenesis in the Paleozoic. *Journal of Paleontology* 80 (4): 638–649. [https://doi.org/10.1666/0022-3360\(2006\)80\[638:JMF TLD\]2.0.CO;2](https://doi.org/10.1666/0022-3360(2006)80[638:JMF TLD]2.0.CO;2)
- WILSON H. M. & ANDERSON L. I. 2004. — Morphology and taxonomy of Paleozoic millipedes (Diplopoda: Chilognatha: Archipolypoda) from Scotland. *Journal of Paleontology* 78 (1): 169–184. <https://tinyurl.com/3emds28r>
- WILSON H. M. & HANNIBAL J. T. 2005. — Taxonomy and trunk-ring architecture of pleurojulid millipedes (Diplopoda: Chilognatha: Pleurojulida) from the Pennsylvanian of Europe and North America. *Journal of Paleontology* 79 (6): 1105–1119. [https://doi.org/10.1666/0022-3360\(2005\)079\[1105:TATAOP\]2.0.CO;2](https://doi.org/10.1666/0022-3360(2005)079[1105:TATAOP]2.0.CO;2)
- WILSON H. M., DAESCHLER E. B. & DESBIENS S. 2005. — New flat-backed archipolypodan millipedes from the Upper Devonian of North America. *Journal of Paleontology* 79 (4): 738–744. [https://doi.org/10.1666/0022-3360\(2005\)079\[0738:NFAMFT\]2.0.CO;2](https://doi.org/10.1666/0022-3360(2005)079[0738:NFAMFT]2.0.CO;2)

*Submitted on 24 February 2022;
accepted on 13 March 2023;
published on 20 October 2023.*