

Stratigraphic correlations between the continental and marine Tethyan and Peri-Tethyan basins during the Late Carboniferous and the Early Permian

Alain IZART ⁽¹⁾, Denis VASLET ⁽²⁾, Céline BRIAND ⁽¹⁾, Jean BROUTIN ⁽³⁾,
Robert COQUEL ⁽⁴⁾, Vladimir DAVYDOV ⁽⁵⁾, Martin DONSIMONI ⁽²⁾,
Mohammed EL WARTITI ⁽⁶⁾, Talgat ENSEPBAEV ⁽⁷⁾, Mark GELUK ⁽⁸⁾,
Nathalya GOREVA ⁽⁹⁾, Naci GÖRÜR ⁽¹⁰⁾, Nayyer IQBAL ⁽¹¹⁾, Geroi JOLTAEV ⁽⁷⁾,
Olga KOSSOVAYA ⁽¹²⁾, Karl KRAINER ⁽¹³⁾, Jean-Pierre LAVEINE ⁽⁴⁾,
Maria MAKHLINA ⁽¹⁴⁾, Alexander MASLO ⁽¹⁵⁾, Tamara NEMIROVSKAYA ⁽¹⁵⁾,
Mahmoud KORA ⁽¹⁶⁾, Raissa KOZITSKAYA ⁽¹⁵⁾, Dominique MASSA ⁽¹⁷⁾,
Daniel MERCIER ⁽¹⁸⁾, Olivier MONOD ⁽¹⁹⁾, Stanislav OPLUSTIL ⁽²⁰⁾,
Jörg SCHNEIDER ⁽²¹⁾, Hans SCHÖNLAUB ⁽²²⁾, Alexander STSCHEGOLEV ⁽¹⁵⁾,
Peter SÜSS ⁽²³⁾, Daniel VACHARD ⁽⁴⁾, Gian Battista VAI ⁽²⁴⁾,
Anna VOZAROVA ⁽²⁵⁾, Tuvia WEISSBROD ⁽²⁶⁾, Albin ZDANOWSKI ⁽²⁷⁾
(see appendix 2 for addresses)



343

Izart A. *et al.* 1998. — Stratigraphic correlations between the continental and marine Tethyan and Peri-Tethyan basins during the Late Carboniferous and the Early Permian, in Crasquin-Soleau S., Izart A., Vaslet D. & De Wever P. (eds), *Peri-Tethys: stratigraphic correlations 2*, *Geodiversitas* 20 (4) : 521-595.

ABSTRACT

The compilation of detailed stratigraphic, sedimentologic and paleontologic data resulted in stratigraphic correlations of marine and continental areas outcropping today in the Tethyan and Peri-Tethyan domains: (1) the base of the Moscovian would correspond to the base of the Westphalian C in the Peri-Tethyan domain and to the base of the Westphalian B in the Tethyan domain; (2) the Kasimovian, the Gzhelian and the Orenburgian would correspond in the northern Peri-Tethyan domain and Tethyan domain (Carnic Alps) respectively to the early Stephanian, the late Stephanian and the Autunian *p.p.*, in the southern Peri-Tethyan domain to an undifferentiated time interval. The boundary between the Stephanian and the Autunian was recognized in the Donets Basin with some doubts; (3) the Asselian, Sakmarian, Artinskian and Kungurian would correspond in all the domains to the Autunian *p.p.* and the Saxonian that remain difficult to separate.

KEY WORDS

Peri-Tethys,
biostratigraphy,
Carboniferous,
Permian,
Tethys,
stratigraphic correlations.

RÉSUMÉ

Corrélations stratigraphiques dans les bassins téthysiens et péri-téthysiens au Carbonifère supérieur et au Permien inférieur. Des corrélations sont proposées entre les domaines marins et continentaux du Paléozoïque supérieur affleurant aujourd'hui dans le domaine téthysien et sur les plate-formes qui le bordent au nord et au sud: (1) la base du Moscovien correspondrait dans le domaine péri-téthysien à la base du Westphalien C et dans le domaine téthysien à la base du Westphalien B ; (2) le Kasimovien, le Gzhélien et l'Orenburgien correspondraient dans le domaine nord péri-téthysien et dans le domaine téthysien (Alpes carniques) respectivement au Stéphanien inférieur, supérieur et Autunien *p.p.*, dans le domaine sud péri-téthysien à un intervalle de temps indifférencié, la correspondance n'étant pas établie précisément. La limite entre le Stéphanien et l'Autunien a été reconnue dans le bassin du Donetz avec incertitudes ; (3) l'Assélien, le Sakmarien, l'Artinskien, le Kungurien correspondraient dans tous les domaines à l'Autunien *p.p.* et au Saxonien, qui restent difficile à différencier.

MOTS CLÉS

Péri-Téthys,
biostratigraphie,
Carbonifère,
Permien,
Téthys,
corrélations stratigraphiques.

INTRODUCTION

In the frame of the IGCP 343, a synthesis of the published and original data from eastern and western Europe, northern Africa and Arabia is presented. Three litho- and chronostratigraphic cross-sections (location Fig. 1) show the results in three domains outcropping nowadays in the northern Peri-Tethyan domain (Fig. 2, Inset Fig. 1), in the Tethyan domain (Fig. 3, Inset Fig. 2) and in the southern Peri-Tethyan domain (Fig. 4, Inset Fig. 3). Biostratigraphic correlations are proposed in the northern Peri-Tethyan domain (Fig. 5) between the marine eastern basins (Russia, Ukraine) and the continental western basins (France, Germany), and are extended to the other domains (Figs 6, 7). Moreover, the latitudinal correlations between the continental basins and the correlations based on the sequence stratigraphy allow to test the climatic, eustatic and tectonic factors. The radiochronological data chart of Hess & Lippolt (1986) and of Menning (1995) are chosen for the Late Carboniferous and for the Permian respectively.

BIOSTRATIGRAPHIC CORRELATIONS BETWEEN THE MARINE AND CONTINENTAL DOMAINS

Correlations are proposed for the Bashkirian-Moscovian and Namurian-Westphalian time intervals in the northern Peri-Tethyan domain (Fig. 8). Sinitsyn *et al.* (1978), Yablokov *et al.* (1978), Aisenverg *et al.* (1978), Makhlina *et al.* (1979), Solovieva (1985), Solovieva *et al.* (1985a, b), Kagarmanov & Donakova (1990), Izart *et al.* (1996), Izart *et al.* (1998), Makhlina *et al.* (1997), Vachard & Maslo (1996), Einoir *et al.* (1996), Briand *et al.* (1998), Zhamoida (in press), Ensebaev *et al.* (this volume) described the Carboniferous of the Ural, Moscow, Donets and Precaspian (Kazakhstan) Basins. The Bashkirian and Moscovian show thin and medium bedded limestones in the Moscow and Ural Basins. During this time, the Donets was a paralic basin with thick alternations of fluvial sandstone, paleosol, coal, marine limestone and claystone, deltaic siltstone and sandstone. In eastern Europe, fusulinid, conodont and coral




FIG. 1. — Location of cross-sections on the paleogeographic map of the North Hemisphere during the Lower Permian modified after Scotese & Langford (1995). Land in white, shallow sea in stipple, ocean in waves, active plate boundaries in bold lines. **AB**, cross-section of the northern Peri-Tethyan Basins; **CD**, cross-section of the Tethyan Basins; **EF**, cross-section of the southern Peri-Tethyan Basins.

biozones were defined for the Bashkirian, Moscovian, Kasimovian, Gzhelian and marine Early Permian (Fig. 5). See also for the fusulinids, Davydov (1996) and the table in Krainer & Davydov (this volume) and for the corals Kossovaya (1996) and Kossovaya (this volume). In the paralic basins of western Europe, the Namurian and the Westphalian present thick alternations of fluvial sandstone, paleosol, coal, marine, lagoonal and lacustrine claystone, deltaic siltstone and sandstone. The flora allows to defi-

ne a biostratigraphy and a chronostratigraphy framework (Fig. 5). The distribution of flora for the North France coal basin (Laveine 1987) and microflora for western Europe (Clayton *et al.* 1977) are presented for the Namurian and Westphalian. The basins of Lorraine (Donsimoni 1981), the Netherlands (Geluk), the Ruhr (Fiebig 1969; Josten 1991), Poland (Zdanowski & Zakowa 1995), the Czech (Oplustil & Pesek this volume) and Slovak (Vozarova this volume) Republics, the Caucasus (Chernyavsky *et al.*

AGE (Ma)	Continental STAGES	FRANCE	GERMANY	DONETS	MOSCOW	URAL	Marine STAGES	
274								
277	Saxonian		SAALE			Irensk Philippovsk	Kungurian	LOWER PERMIAN
283			Hornburg			Saraninsk Sarginsk Irginsk Burtsevsk	Artinskian	
290	Autunian	AUTUN Millery Surmoulin		Kramatorsk		Sterlitamak Tastubsk	Sakmarian	
296		Muse Igornay Epinac	Sennewitz Halle	Slavjansk Nikitovsk Kartamysh	Sokolvesarsk	Shikansk Kholodnolozk	Asselian	
300		SAINT- ETIENNE Couronnement Janon	Mansfeld	Mironovsk	Melekhovo Noginsk	Zianchurinsk	Orenburgian	
302				Kalinovsk	Pavlov-Posad Amerevo Rechitsy		Gzhelian	
305		Stephanian	Gier	Grillenber	Torezk	Dorogomilovo Khamovnich Kreyyakino	Abzanovsk	
308	Westphalian D	N FRANCE Assise de Bruay	RUHR Ibbenburen	Sanjarovsk Sabovsk	Myachkovo Podolsk	Bolshekynsk	Moscovian	
311	Westphalian C		Lembeck Dorsten	Marievsk Kamensk	Kashira Vereia	Kirovsk		
313	Westphalian B	A. d'Anzin	Horst Essen	Krasnodonsk		Asatausk	Bashkirian	
315	Westphalian A	A.de Vicoigne	Bochum Witten	Makeevsk Zuevsk	Upper Aza	Tashatinsk		
317	Namurian C	A. de Flines	Sprockhövel	Blagodatnensk Manuilovsk	Lower Aza	Askynbashsk		
319	Namurian B		Vorhalle Hagen	Feninsk		Akavask Siuransk		

FIG. 2. — The formations of the northern Peri-Tethyan Basins.

1978) and North Turkey (Zonguldak; Kerey *et al.* 1985; Görür *et al.* 1997) are also considered in this study. The Donets Basin is the only

northern Peri-Tethyan Basin, where the correlation between the marine and continental domains has been attempted. In this basin, the

AGE (Ma)	Continental STAGES	SPAIN		ITALY	CARNIC ALPS	IRAN	Marine STAGES	
274								
277	Saxonian	PALENCIA	PICOS DE EUROPA				Kungurian	LOWER PERMIAN
					Goggau Tressdorf		Artinskian	
283		unnamed			Trogkofel Upper Pseudo -Schwagerina	Dorud	Sakmarian	
290	Autunian				Grenzland		Asselian	
296			Rio Marina		Lower Pseudo -Schwagerina Carnizza Auernig Corona Pizzul		Orenburgian	UPPER CARBONIFEROUS
300	Stephanian	Villablino Sabero						
302			Barruelo	Puentelles	Spirifer		Kasimovian	
305	Cantabrian	Branosera	Gamonedo		Meledis Bombaso	Gheselghaleh		
308	Westphalian D	Central ASTURIAS Sama	Picos de Europa	San Antonio			Moscovian	
311	Westphalian C	Lena						
313	Westphalian B			Carpineta				
315	Westphalian A							
317	Namurian C	Valdeteja	Valdeteja					
319	Namurian B						Bashkirian	

FIG. 3. — The formations of the Tethyan Basins.

AGE (Ma)	Continental STAGES	MOROCCO	ALGERIA	TUNISIA	LIBYA Cyrenaica	OMAN	Marine STAGES	
274								
277	Saxonian	unnamed				Lower Gharif	Kungurian	LOWER PERMIAN
283				unnamed limestones	unnamed limestones		Artinskian	
290				unnamed limestones KRP2			Sakmarian	
296	Autunian		unnamed limestones KRP1	unnamed limestones	Al Khlata	Asselian		
300	Stephanian	Senhadja					Orenburgian	
302			unnamed limestones KRC3				Gzhelian	
305			MEZARIF		Kasimovian			
308	Westphalian D	JERADA	Nekheila	unnamed limestones KRC2			Moscovian	UPPER CARBONIFEROUS
311	Westphalian C	Assise de Jerada	Carbonates de base					
313	Westphalian B	F. inférieure de Jerada		unnamed limestones	unnamed limestones		Bashkirian	
315	Westphalian A	Schiste supérieur						
317	Namurian C	Schiste inférieur						
319	Namurian B							

FIG. 4. — The formations of the southern Peri-Tethyan Basins.

flora are closed to western Europe ones (Fissunencko & Laveine 1984), but some differences could be sometimes noted. The microflora presents a local biozonation (Coquel *et al.* 1984). At the boundary between the Westphalian B and C, a disharmony is observed in the appearance of species more precocious or later in the Donets Basin than in western Europe. The base of the Moscovian would be inside the Westphalian B or at the base of the Westphalian C (Fig. 8).

Correlations are proposed for the Kasimovian-Gzhelian-Orenburgian and the Stephanian-Autunian in the northern Peri-Tethyan domain. The Kasimovian, the Gzhelian and the Orenburgian show thin limestones in the Moscow Basin and thick paralic facies in the Donets Basin. The Kasimovian, the Gzhelian and the Orenburgian (Fig. 5) present fusulinid, conodont and coral biozones in the Moscow, Ural and Donets Basins. Davydov (1990, 1992) moved the *Ultradaixina bosbytauensis* biozone from the Asselian to the late Gzhelian. The boundaries of this biozone present an uncertainty, because Davydov defined this biozone in the carbonated marine facies of the pre-Donets Basin in Russia, whereas the Donets Basin in Ukraine shows lagoonal facies without fusulinids. Then, Davydov (1996) put the *Daixina sokensis* and *Ultradaixina bosbytauensis* biozones in the Orenburgian, between the Gzhelian and the Asselian. The boundary between the Carboniferous and the Permian is located at the top of the Orenburgian. The Stephanian, defined in the limnic basin of Saint-Étienne, was subdivided by Doubringer *et al.* (1995) into a early Stephanian (Barruelian) corresponding to the Stephanian A and an late Stephanian (Forezian) corresponding to the Stephanian B and C with a precise biozonation of flora (Fig. 5). The Stephanian presents breccia along the sides of the basin and lacustrine claystone and coal in the center (Becq-Giraudon *et al.* 1995, Mercier this paper). The early Stephanian exhibits hygrophytic Lycophytes and Filicophytes (*Pecopteris* sp., *Astherotheca lamuriana*, Fig. 5) and presents Westphalian affinities. The late Stephanian shows hygrophytic Pteridosperma-phytes and Lycophytes (*Odontopteris* sp., *Pecopteris* sp., *Sphenopteris* sp., *Alethopteris zeilleri*, *Sphenophyllum angustifolium*) and

some occurrences of xerophytic plants (*Autunia conferta*, *Lodevia nicklesii*). The biozonation of Clayton *et al.* (1977) applies in Saint-Étienne with some occurrences of xerophytic spores with *Potonieisporites novicus* and *Vittatina* sp. Alternation of hygrophytic and xerophytic floras and microfloras are observed through the late Stephanian in numerous basins (Broutin *et al.* 1986, 1990). The basins of Lorraine (Donsimoni 1981), the Saale (Schneider & Gebhardt 1993), Poland (Zdanowski, this paper), the Czech (Oplustil this paper) and Slovak (Vozarova this paper) Republics, the Caucasus (Chernyavsky *et al.* 1978) and North Turkey (Kerey *et al.* 1985) are also considered. In the Donets Basin (Fig. 8), the biozonation of flora is local (Boyarina, in press; Izart *et al.* 1998) with some common plants as *Astherotheca lamuriana* in the late Stephanian. There were alternations of hygrophytic and xerophytic flora and microflora (*Lodevia nicklesii*, *Potonieisporites novicus*) in the Orenburgian (Stschegolev in Aisenverg *et al.* 1978; Stschegolev & Kozitskaya 1984; Izart *et al.* 1998). The Kasimovian (Fig. 8) would be equivalent to the early Stephanian. The Gzhelian (Fig. 8) would be the equivalent of the late Stephanian, the Orenburgian of the early Autunian *p.p.* if we refer to the appearance of the xerophytic plants, or of the late Stephanian if we refer to the acme of the xerophytic plants. The beginning of the sedimentation in the Stephanian and Autunian Basins that depends on their tectonic opening will be heterochronous in western Europe.

Correlations are proposed between the marine and continental Early Permian in the northern Peri-Tethyan domain. The marine Early Permian was defined in the Ural Basin (Chuvashov 1993). The Asselian, the Sakmarian, the Artinskian Stages have a carbonate facies whereas the Kungurian is evaporitic. Biozones of fusulinids, conodonts and corals were defined (Fig. 8). The new Permian chronostratigraphic subdivisions (Yugan 1996; Yugan *et al.* 1997) were utilized. In the Moscow Basin, the Asselian is carbonated and in the Donets Basin, the Asselian and the Sakmarian exhibit red claystone and evaporites. The Autunian, defined in the limnic basin of Autun (Fig. 5), shows fluvial sandstone and

WESTERN EUROPE (FRANCE - GERMANY)						EASTERN EUROPE (DONETS - MOSCOW - PERM)										
Age (Ma)	Sequences (1)	Goniatites (2)	Flora (3)	Microflora (4)	CONTINENTAL STAGES	Sequences (1)	Flora (5)	Microflora (6)	Foraminifera (7)	Conodonts (8)	Corals (9)	MARINE STAGES				
274	SA				SAXONIAN	KU			<i>Nodosaria sexangulata</i> <i>Parafusulina solidissima</i>	<i>Neostreptognathodus pequoppensis</i>		KUNGURIAN	LOWER PERMIAN			
277						AR			<i>Nodosaria pugioidea</i>					<i>Lophophyllidium sp.</i>		
283									<i>Hemigordius saranaensis</i>							
	AU							<i>Parafusulina solidissima</i>	<i>Sweetognathodus whitei</i>	<i>Protolonsdaleiastraea juresanensis</i>	ARTINSKIAN					
								<i>Pseudofusulina juresanensis</i>		<i>Protolonsdaleiastraea juresanensis</i>						
<i>Pseudofusulina pedissequa</i>																
290						S3			<i>Pseudofusulina urdalensis</i>	<i>Sweetognathodus primus</i>	<i>Protolonsdaleiastraea longiseptata</i>	SAKMARIAN				
						S2			<i>Pseudofusulina verneuilli</i> <i>Pseudofusulina uralica</i>	<i>Neogondolella lata</i>	<i>P. biseptata</i> <i>Timania schmidtii</i>					
									S1	<i>Pseudofusulina moelleri</i>	<i>Neogondolella uralensis</i>			<i>Kleopatrina magnifica</i>		
296					AUTUNIAN	A3			<i>Sphaeroschwagerina sphaerica</i>	<i>Streptognathodus postfusus</i>	<i>Kleopatrina pseudoelegans</i>	ASSELIAN				
						A2			<i>Sphaeroschwagerina moelleri</i>	<i>Streptognathodus fusus</i> <i>Streptognathodus constrictus</i>	<i>Tschussosvenskia captiosa</i>					
						A1			<i>Sphaeroschwagerina fusiformis</i>	<i>Streptognathodus cristellaris</i>	<i>Ferganophyllum uralicum</i>					

Autunia conferta
Rhacophyllum schenkii

Disaccites striatiti
Potonielsponites spp.
Vitalina spp.

Autunia conferta

Potonielsponites "horvici-bhardwaji"
Vitalina spp.




Fig. 5. — Biostratigraphy and chronostratigraphy of the northern Peri-Tethyan Basins. 1, Izart (this paper); 2, Bouroz *et al.* (1978); 3, Laveine (1987), Broutin (this paper); 4, Clayton *et al.* (1977), Coquel & Broutin (this paper); 5, Fissunenkeno & Laveine (1984), Stschegolev & Broutin (this paper); 6, Coquel *et al.* (1984), Inosova *et al.* (1976), Broutin (this paper); 7, Solovieva *et al.* (1985a, b), Davydov (1990, 1992), Chuvashov (1993), Vachard & Maslo (this paper); 8, Alekseev, Goreva, Kozitskaya & Nemirovskaya (this paper); 9, Kossovaya (this paper).

SPAIN (PALENCIA-CENTRAL ASTURIAS)							SPAIN (PICOS DE EUROPA)			CARNIC ALPS					
Age (Ma)	Sequences (1)	Flora (2)	Microflora (3)	CONTINENTAL STAGES	Foraminifera (4)	MARINE STAGES	Sequences (1)	Foraminifera (5)	MARINE STAGES	Sequences (1)	Foraminifera (6)	Flora (7)	MARINE STAGES		
274	SA	Autunia conferta		SAXONIAN						S			KUNGURIAN	Autunia conferta	LOWER PERMIAN
277											Schwagerina vulgaris Praeparafusulina lutugini		ARTINSKIAN		
283	Paraschwagerina inflata Schwagerina moelleri			SAKMARIAN											
290											ASSELIAN				
296	AU			AUTUNIAN						A	Sphaeroschwagerina glomerosa				




Fig. 6. — Biostratigraphy and chronostratigraphy of the Tethyan Basins. 1, Izart (this paper); 2, Wagner & Winkler-Prins (1985), Martinez Garcia (1991); 3, Leyva *et al.* (1985), Coquel & Rodriguez (1995); 4, Lys (1988), Granados *et al.* (1985); 5, Villa (this paper); 6, Krainer (1992, 1993), Vai & Venturini (1997); 7, Krainer (this paper).

MOROCCO - ALGERIA - TUNISIA - LIBYA - EGYPT									SAUDI ARABIA		OMAN		
Age (Ma)	Sequences (1)	Goniatites (2)	Flora (3)	Microflora (4)	CONTINENTAL STAGES	Sequences (1)	Foraminifera (5)	MARINE STAGES	Sequences (1)	Microflora of Arabia (6)	Microflora of Oman (7)	STAGES	
274	SA				SAXONIAN			KUNGURIAN				KUNGURIAN	
277								ARTINSKIAN				ARTINSKIAN	
283													
290	AU		Lodevia nicklesi Rhaetophyllum schenkii		Potoniesporites "novius bhardwajii" Vittatina spp.	S	Schwagerina sp. Staffellidae	SAKMARIAN	S			SAKMARIAN	
296								ASSELIAN				A	
											Cycadopites cymbatus Protohaploxyrinus spp. Vittatina spp. Kingslacopites subcircularis	ASSELIAN	LOW PERMIAN

Lodovia nicklesi
Rhachiphyllum schenkii

Potoniesporites "novius bhardwaji"
Vittatina spp.

Cycadopiles cymbatus
Prothaploxypinus spp.
Vittatina spp.
Kingiapolpites subcircularis

LOWER PERMIAN





Fig. 7. — Biostratigraphy and chronostratigraphy of the southern Peri-Tethyan Basins. 1, Izart (this paper); 2, Deleau (1951), Owodenko (1976); 3, Deleau (1951), Nedjari (1982), Desteuq *et al.* (1988), Broutin & El Wartiti (this paper); Aassoumi (1994); 4, Doubinger & Fabre (1983), Aassoumi (1994); 5, Lys (1988), Nedjari (1982), Massa & Vachard (1979), Kora (this paper); 6, Owens & Turner (1995); 7, Love (1994).



lacustrine claystone with xerophytic flora and microflora (*Autunia conferta*, *Rhachiphyllum schenkii*, *Potonieisporites* sp., *Vittatina* sp.). The Saar-Nahe Basin presents red fluvial sandstone and lacustrine black shale (Stapf 1997). The Saale Basin in Germany (Schneider & Gebhardt 1993; Schneider *et al.* 1994, 1995; Schneider 1996; Schneider & Rössler in press) exhibits two informal lithological units named early and late "Rotliegend". The early "Rotliegend" includes grey conglomerates and sandstones that would be dated to the Autunian by plants (*Autunia conferta*) and the late "Rotliegend" red conglomerates and sandstones that would be dated to the Saxonian by tracks of reptile. The boundary of these units presents an incertitude between the basins of North Germany and the Saale (Menning 1995). The basins of Poland (Zdanowski this paper), the Czech (Oplustil this paper) and Slovak (Vozarova this paper) Republics are also considered. In the Donets Basin (Fig. 5), the flora and microflora of the Asselian and Sakmarian Stages are xerophytic (Inosova *et al.* 1976) with *Autunia conferta*, *Potonieisporites* sp. and *Vittatina* sp. The climate changed from a humid tropical to a dry tropical climate (Parrish 1995). According to Schneider *et al.* (1994, 1995), the Kartamysh, Nikitovsk and Slavjansk formations would be equivalent to the early "Rotliegend" and the Kramatorsk formation to the late "Rotliegend" with some incertitudes. The Asselian would correspond to the Autunian *p.p.* and the Sakmarian, Artinskian and Kungurian to the Saxonian (Fig. 8).

Correlations are proposed in the Tethyan (Fig. 6) and southern Peri-Tethyan domains (Fig. 7). The Bashkirian and Moscovian are dated by fusulinids in the paralic basin of the Cantabrian zone in Palencia (Lys 1988a) and in the central Asturias (Granados *et al.* 1985), the limestones of the Picos de Europa (Villa 1985; Villa *et al.* 1993), Tuscany (Vai 1991, 1994; Pasini & Vai 1997), eastern Elbourz (Jenny *et al.* 1978; Jenny-Deshusses 1983), Mezarif, Algeria (Nedjari

FIG. 8. — Correlations of the continental and marine Late Carboniferous and Early Permian of the Tethyan and Peri-Tethyan Basins. Oblique lines: uncertainty.

1982; Lys 1988a), Tunisia (Lys 1988b), Turkey (Monod 1977; Lys 1988a), Libya (Massa & Vachard 1979; Vachard *et al.* 1993), Egypt (Kora 1995), Syria (Al Youssef & Ayed 1992). The Carnic Alps (Schönlaub 1992; Krainer 1993; Vai 1991; Vai & Venturini 1997; Krainer & Davydov, this volume) present only the late Moscovian. Some marine bands dated to the Moscovian are known in Morocco, Jerada (Owodenko 1976) and in Algeria, Bechar (Deleau 1951). The Kasimovian is found everywhere except in Algeria (Bechar, Mezarif). The Gzhelian and the Orenburgian are found everywhere except in Spain, Algeria (Bechar, Mezarif) and southern Tunisia (Lys 1988b). The marine Early Permian is found everywhere except in Spain, Algeria (Bechar, Mezarif) and Libya (Ghadames). The marine facies of the Late Carboniferous and Early Permian are unknown in Saudi Arabia. In Oman, the Early Permian presents marine and continental levels (Broutin *et al.* 1995; Angiolini *et al.* 1997).

The Westphalian is paralic in central Asturias, in Jerada (Morocco) and in Bechar (Algeria). The hygrophytic flora and microflora are dated the Westphalian A to D in central Asturias (Leyva *et al.* 1985a, b, c; Granados *et al.* 1985; Saenz de Santa Maria *et al.* 1985; Wagner & Alvarez 1991) and in Jerada (Owodenko 1976; Desteucq *et al.* 1988; Izart 1991), the Westphalian C and D in Bechar (Deleau 1951; Coquel *et al.* 1988). Elsewhere the base of the Moscovian could be located in Spain inside the Westphalian A or B (Wagner & Bowman 1983; Granados *et al.* 1985; Leyva *et al.* 1985a, b, c; Martinez Dias *et al.* 1985; Ginkel & Villa 1996), and in Algeria at the base of the Westphalian C (Lys 1988a) as in eastern Europe, but there is an objection: an eustatic event is synchronous everywhere in the world, or the appearance of plants in the Westphalian of Spain presents a disharmony. In Libya (Massa *et al.* 1980; Coquel *et al.* 1988), in Saudi Arabia (Owens & Turner 1995), in Oman (Love 1994) as in Gondwanaland, a xerophytic microflora with *Potonieisporites* sp. is known in the Westphalian D.

In the Cantabrian zone in Palencia, Bouroz *et al.* (1972), Wagner & Winkler-Prins (1985, 1991,

1993) defined the Cantabrian and the Stephanian in the interval attributed by Lys (1988a) to the late Moscovian (Myachkovian) and the Kasimovian. The comparison between the flora and microflora (Coquel & Rodriguez 1995) does not allow to differentiate the Cantabrian and the Stephanian. The precocious tectonic opening of the basin of the Cantabrian zone in Palencia would explain the presence of the Cantabrian in this basin and its absence in the Saint-Étienne Basin. The lower part of the Cantabrian could be connected with the Westphalian D and the upper part with the early Stephanian. In the Carnic Alps, Krainer (1993) reported in the Kasimovian and the Gzhelian hygrophytic flora dated to the Stephanian (*Pecopteris* sp., *Sphenophyllum* sp.). In Morocco, hygrophytic plants dated to the late Stephanian exist in the High Atlas (Beauchamp *et al.* 1986; Doubinger & Roy-Dias 1985, 1986; Aassoumi 1994). In Oman, Love (1994) described microflora *Microbaculispora* known in the Gondwanaland. The equivalence between the Kasimovian-Gzhelian and the Stephanian is established in the Tethyan domain, but not in the southern Peri-Tethyan domain.

The Autunian presents xerophytic plants (*Autunia conferta*) in the Carnic Alps (Krainer 1993) and in Spain where in addition the Saxonian exists (Martinez-Garcia 1991). The Autunian and the Saxonian exhibit xerophytic flora and microflora with *Lodevia nicklesii*, *Rhachiphyllum schenkii*, *Potonieisporites* sp. and *Vittatina* sp. in central Morocco (Broutin *et al.* 1987; El Wartiti *et al.* 1986, 1990; Aassoumi 1994). Doubinger & Fabre (1983) attributed to the Autunian a microflora with *Torispora* sp., *Potonieisporites* sp. at the top of the Bechar Series. In Oman, Love (1994) distinguished the biozone *Cycadopites* dated to the Asselian and Sakmarian and the biozone *Kingiacolpites* dated to the late Sakmarian to Kungurian, known in Gondwanaland. In the Salt Range (Pakistan), the biozone *Cycadopites* (Iqbal 1993 and this volume) was found at the base of the Warcha Formation, corresponding to the Sakmarian-Artinskian interval. Everywhere, the correlations are imprecise between the marine and continental Permian.

LATITUDINAL CORRELATIONS

The stratigraphic correlations provided by the plants of the continental domain will depend on the latitudinal location of each domain. Following the position of the continents according to Scotese & McKerrow (1990), this hypothesis can be tested on the basis of the relations between the hygrophytic plants and the equatorial-humid tropical climate and between the xerophytic plants and the dry tropical-desert climate. The geographic zonation of climate has been described in term of floral biomes for the Early Permian (Sakmarian) and Late Permian (Kazanian) by Ziegler (1990) and Ziegler *et al.* (1997). The biomes are: (1) equatorial and tropical everwet, (2) tropical and subtropical summerwet, (3) coastal and inland tropical desert, (4) winterwet, (5) western and eastern warm temperate, (6) western and eastern cool temperate, (7) midlatitude desert, (8) cold temperate, (9) arctic and (10) glacial. This approach has been adapted from a climate study of the Recent. The sediments gave also informations about the climates: coal in various climates (cold to tropical), reef in tropical climate, evaporite in dry tropical or desert climate, tillites in cold climate. Numerous paleoclimate studies exist for the Permian: Parrish (1995), Kutzbach & Gallimore (1989), Kutzbach & Ziegler (1993), Barron & Fawcett (1995); Crowell 1995. All these studies exhibit from Late Carboniferous to Trias an increase of the aridity in the humid tropical zone, that was explained by the perfect symmetry of the continental surfaces at the equator, that implied a warm and dry climate with monsoons. In the northern Peri-Tethyan domain, the location of western and eastern Europe was between the equator and 5°N during the Westphalian, between 5°N and 10°N during the Early Permian (Fig. 1). All the basins were located inside the same latitudinal zone and the latitudinal correlations are excellent. The Westphalian presents hygrophytic plants and coal in the everwet biome, the Stephanian alternation of hygrophytic and xerophytic plants (Saint-Étienne, Saale, Donets) and alternation of red beds and grey beds with coal (Lorraine, Saale, Czech Republic, Slovakia, Donets) explained by period of mon-

soon in the summerwet biome. The Autunian exhibits xerophytic plants growing on the slopes of basins and alternation of red beds and lacustrine black shale (Autun, Lorraine, Saar-Nahe, Saale, Czech Republic and Slovakia) or coal (Donets) in the summerwet biome. The Saxonian presents xerophytic plants, red beds, calcretes and evaporites (Saale, Czech Republic, Slovakia, Donets, Russian platform) in the subtropical desert biome.

In the Tethyan domain (Fig. 1), Spain, Italy and the Carnic Alps were located at 5°S during the Westphalian and Stephanian and at the equator during the Early Permian. The Westphalian and Stephanian present hygrophytic plants and coal in the everwet biome, the Autunian xerophytic plants growing on the slopes of basins and red beds in the summerwet biome, the Saxonian xerophytic plants and red beds in the subtropical desert biome in spite of its location at the equator. The presence of this subtropical biome at the equator can be explained by the perfect symmetry of the continental surfaces at the equator (Parrish 1995). The correlations are excellent between the northern Peri-Tethyan and Tethyan domains. The presence of two opposite climates at the equator during the Westphalian and the Early Permian makes the correlations easier.

In the southern Peri-Tethyan domain (Fig. 1), North Africa was located at 10°S during the Westphalian and at the equator during the Early Permian. Libya, Egypt and Saudi Arabia were located at the 25°S during the Westphalian and 20°S during the Early Permian; Oman and the Salt Range were located at 50°S during the Westphalian and at 40°S during the Early Permian. In North Africa, the Westphalian and the Stephanian present hygrophytic plants and coal in the everwet biome, the Autunian xerophytic plants growing on the slopes of basins and red beds in the summerwet biome, the Saxonian xerophytic plants and red beds in the subtropical desert biome in spite of its location at the equator and for the same reason as the Tethyan domain. In Libya, Egypt and Saudi Arabia, the plants were xerophytic during the Westphalian and the Early Permian. In Oman and the Salt Range as in all Gondwanaland, the plants lived in a cool to temperate climate. These different

latitudinal locations make the correlations difficult between Europe and Gondwanaland.

During the Late Carboniferous and Early Permian, three realms are defined (Ziegler 1990): the tropical Cathaysian (biome 1) in China and Euramerican-Atlantic in Europe and North America (biome 1 during Carboniferous and biomes 2 and 3 during Permian) realm, the North temperate Angaran realm (biomes 4 to 8) in Siberia and the southern temperate Gondwanan realm (biomes 5 to 10) in Gondwana. Biome 1, represented by tropical rainforests populated by arborescent hygrophytic flora, was known in China during the Carboniferous and Permian and in Europe and North America during the Late Carboniferous. Biome 2 was represented by *Callipteris* and primitive conifers (*Walchia*) during the Early Permian in Europe, North Africa and North America. Biome 6 was represented by *Glossopteris* and Biome 8 by *Gangamopteris* in Gondwana.

CORRELATIONS FOUND ON THE SEQUENCE STRATIGRAPHY

A sequence stratigraphy of the Carboniferous and Permian was built by Ross & Ross (1987, 1994, 1995). Second order, third order (Figs 5-7), fourth order and high frequency sequences were defined in eastern Europe in Moscow Basin (Briand *et al.* 1998), the Donets Basin (Izart *et al.* 1996 and 1998) and the Central Ural Basin (Izart *et al.* 1998), and in the western Europe in paralic and limnic basins (Izart & Vachard 1994), in the foreland basin of Asturias and in the rift basin of the Carnic Alps (Samankassou 1997; Vai & Venturini 1997; Krainer & Davydov, this volume), in the intracontinental basins or the marine platform of North Africa (Izart 1991).

For the Late Carboniferous, the same number of third order and second order sequences was found in the Moscow and Donets Basins and a different number in western Europe. In the Moscow and Donets Basins, the Moscovian, Kasimovian and Gzhelian-Orenburgian deposits each form a second order sequence, the Moscovian is subdivided into four third order

sequences, the Kasimovian into three or four sequences and the Gzhelian-Orenburgian into four or five sequences. In western Europe, the Westphalian presents four third order sequences, the Stephanian two, the Autunian one, the Saxonian one. The number of sequences is different from Ross & Ross (1987), who used fourth order sequences. In eastern Europe, they resulted from the eustasy that produced synchronous sequences and tectonics. In western Europe, the sequences are tectonically controlled and heterochronous with eastern Europe.

For the Late Carboniferous, the fourth order sequences (FOS, 400 000 y.) and high frequency sequences (HFS, 40 000 y. to 250 000 y.) are variable everywhere, as shown by Izart & Vachard (1994) and these results: for Moscovian, eleven FOS and forty HFS in Moscow and eighteen FOS and one hundred HFS in Donets; for Kasimovian, eight FOS and twenty HFS in Moscow, seven FOS and twenty-six HFS in Donets; for Gzhelian-Orenburgian, nine FOS and twenty-four HFS in Moscow, sixteen FOS and fifty-two HFS in Donets; for Westphalian, ten FOS in western Europe and for Westphalian C, four FOS and sixteen HFS in England; four FOS and forty HFS in North France and Germany. The number of FOS is different from Ross & Ross (1987) and the number is variable everywhere. In the Moscow Basin, eustasy prevails certainly over tectonics and in the Donets and western Europe Basins, tectonics prevail over eustasy. The average duration of HFS in the Moscow and western Europe Basins is near the periodicity of eccentricity (100 000 y.) and in the Donets Basin near the periodicity of obliquity (40 000 y.). However, a better accuracy of time by radiochronology is needed to calibrate these sequences.

The Early Permian deposits in Central Ural form two second order sequences, one during Asselian and one during the Sakmarian and Artinskian, five third order sequences for the Asselian, three for the Sakmarian, two composite for Artinskian and one composite for the Kungurian. The Early Permian deposits in the Donets Basin form one second order sequence during the Asselian and the Sakmarian, three third order sequences during the Asselian and one during the

Sakmarian. The number of sequences is weakly different from Ross & Ross (1987, 1994, 1995): five for Asselian, four or five for the Sakmarian, three or four for the Artinskian and one composite or two for the Kungurian. These sequences are controlled by eustasy and tectonics in the Ural foreland Basin and the Donets Basin.

For the Early Permian, the fourth order sequences (FOS) and high frequency sequences (HFS) are variable everywhere: for the Asselian, eleven FOS and eighteen HFS in Donets and a number superior to twenty-four HFS in Central Urals; for Sakmarian, ninety-two HFS in Central Urals; for the early Artinskian (Bursevkian-Irginian), seventeen HFS in Central Urals. These sequences are controlled by eustasy and tectonics. The average duration of HFS is near the periodicity of eccentricity (100 000 y.).

The tectonic control was important in the Asturias and the eustatic control rules over in the Carnic Alps. The sequences are heterochronous in the Asturias and synchronous with eastern Europe in the Carnic Alps. The tectonics prevails in the continental basins of Morocco, the eustasy in the marine basins in Morocco, Algeria, Tunisia, Libya and Egypt. These sequences of the southern Peri-Tethyan domain are heterochronous in the intracontinental basins and synchronous with eastern Europe in the marine basins. The beginning of the sedimentation in the continental stages of the Late Carboniferous and Early Permian are heterochronous according to the tectonic phases affecting each basin, even though it is synchronous in the marine stages that are under the control of the eustasy.

CONCLUSION

Stratigraphic correlations (Fig. 8) are proposed between the marine and continental facies in the northern Peri-Tethyan, Tethyan and southern Peri-Tethyan domains. The base of the Moscovian would be located between the late part of the Westphalian B and the base of the Westphalian C in the northern Peri-Tethyan domain, between the bases of the Westphalian A and C in the Tethyan domain, at the base of the Westphalian C in the southern Peri-Tethyan

domain. The Moscovian would be equivalent to the Westphalian C and D in the northern and southern Peri-Tethyan domains and to an imprecise time interval (Westphalian A to Cantabrian *p.p.*) in the Tethyan domain. The Kasimovian would correspond to the early Stephanian in the northern Peri-Tethyan domain, to the Cantabrian *p.p.* and the early Stephanian in the Tethyan domain. The equivalence between the Kasimovian and the early Stephanian is not established in the southern Peri-Tethyan domain. The Gzhelian would be equivalent to the late Stephanian and the Orenburgian to the lower part of the Autunian in the northern Peri-Tethyan domain with an uncertainty, because we can choose either the appearance of xerophytic plants, or the acme for the definition of the base of the Autunian. In the Carnic Alps, the Gzhelian would be equivalent to the late Stephanian. The equivalence between the Gzhelian and the late Stephanian is not established in the southern Peri-Tethyan domain. The marine Early Permian would be equivalent to the continental Early Permian in all the domains, the Autunian and the Saxonian remain difficult to separate. A latitudinal good correlation is observed on the continents except near Gondwanaland. A good correlation linked with the eustasy is observed in the marine domain and a heterochrony in the beginning of the sedimentation linked with the tectonics is reported in the continental domain.

Acknowledgements

This paper, financed by the Peri-Tethys Program and BRGM, is the final report of the theme "Stratigraphic correlations of the Late Carboniferous and Early Permian" of the IGCP 343 "Stratigraphic correlations of Peri-Tethyan epicratonic Basins". We thank the reviewers of this paper, peculiarly Professor J.-B. Vai and Professor J. Dercourt for their advices.

REFERENCES

- Aassoumi H. 1994. — *Les paléoflores du Permien du Maroc Central. Implications biostratigraphiques, paléocéologiques et paléogéographiques pour le domaine*

- péri-téthysien occidental*. Thèse, Mémoires des Sciences de la Terre de l'université Pierre-et-Marie-Curie, Paris, France, 94.16, 226 p.
- Aisenverg D. E., Lagutina V. V., Levenstein M. L. & Popov V. S. 1978. — Field excursion guidebook for the Donets Basin, in *Compte Rendu du 8^e Congrès international sur la stratigraphie du Carbonifère, Moscow, 1975*. Nauka, Moscow, 360 p.
- Al Laboun A. A. 1993. — *Lexicon of the Paleozoic and lower Mesozoic of Saudi Arabia*. Ph.D. Thesis, Jeddah, Arabia, 102 p.
- Al Youssef W. & Ayed H. 1992. — Evolution of upper paleozoic sequences and the application of stratigraphy as a tool for hydrocarbon exploration in Syria. *Proceedings of the eleventh petroleum exploration and production conference, Cairo 1992* 2: 636-657.
- Angiolini L., Bucher H., Pillevuit A., Platel J.-P., Roger J., Broutin J., Baud A., Marcoux J. & Al Hashmi H. 1997. — Early Permian (Sakmarian) brachiopods from southeastern Oman. *Geobios*, Lyon 30 (3) : 379-406.
- Barron E. J. & Fawcett P. J., 1995. — The climate of Pangea: a review of climate model simulations of the Permian: 37-52, in Scholle P. A., Peryt T. M. & Ulmer-Scholle D. S. (eds), *The Permian of Northern Pangea*, Volume 1: *Paleogeography, Paleoclimates, Stratigraphy*. Springer, Berlin.
- Beauchamp J., Doubinger J., Roy-Dias C. & Sagon J.-P. 1986. — L'Autuno-Stéphanien de la région de Marrakech (Maroc). Implication paléobotanique et paléoclimatique. Comparaison avec les régions limitrophes. *Annales de la Société géologique du Nord* 106 : 145-153.
- Becq-Giraudon J.-F., Mercier D. & Jacquemin H. 1995. — Postface, in Doubinger J., Vetter P., Langiaux J., Galtier J. & Broutin J., La flore fossile du bassin houiller de Saint-Étienne. *Mémoires du Muséum national d'histoire naturelle*, Paris 164, 356 p.
- Bouroz A., Buisine M., Chalard J., Dalinval A. & Dolle P. 1964. — Bassin houiller du Nord et du Pas de Calais, in *5^e Congrès international de stratigraphie et de géologie du Carbonifère, Paris 1964*, Voyage d'étude n°1, 30 p.
- Bouroz A., Knight J. A., Wagner R. H. & Winkler-Prins C. F. 1972. — Sur la limite Westphalien-Stéphanien et sur les subdivisions du Stéphanien inférieur *sensu lato* : 241-261, in *Compte Rendu du 7^e Congrès international de stratigraphie et de géologie du Carbonifère, Krefeld 1971*, Volume 1.
- Bouroz A., Einor O. L., Gordon M., Meyen S. V. & Wagner R. H. 1978. — Propositions pour une classification chronostratigraphique internationale du Carbonifère. *Revue de l'industrie minière* 60 (10) : 469-483.
- Boyarina N. 1997. — Phytocoenosis characteristics of Late Carboniferous-Early Permian for the Donets Basin of Ukrain: 213-218, in *Compte Rendu du 13^e Congrès international de stratigraphie et de géologie du Carbonifère et du Permien, Krakow 1995*.
- Briand C., Izart A., Vaslet D., Vachard D., Makhlina M., Goreva N., Kossovaya O. & Jaroshenko A. 1998. — Sequence stratigraphy of the Moscovian, Kasimovian and Gzhelian in the Moscow basin. *Bulletin de la Société géologique de France* 169 (1) : 35-52.
- Broutin J., Doubinger J., Langiaux J. & Primey D. 1986. — Conséquences de la coexistence de flores à caractères stéphanien et autunien dans les bassins limniques d'Europe occidentale. *Mémoires de la Société géologique de France Nouvelle Série* 149 : 15-25.
- Broutin J., El Wartiti M., Freytet P., Heyler D., Larhib M. & Morel J.-L. 1987. — Nouvelles découvertes paléontologiques dans le bassin détritico-carbonaté permien de Tiddas (Maroc central). *Comptes Rendus de l'Académie des Sciences*, Paris 305 (IIa) : 143-148.
- Broutin J., Doubinger J., Farjanel G., Freytet P., Kerp H., Langiaux J., Lebreton M.-L., Sebban S. & Satta S. 1990. — Le renouvellement des flores au passage Carbonifère Permien : approches stratigraphique, biologique, sédimentologique. *Comptes Rendus de l'Académie des Sciences*, Paris 311 (IIa) : 1563-1569.
- Broutin J., Roger J., Platel J.-P., Angiolini L., Baud A., Bucher H., Marcoux J. & Al Hashmi H. 1995. — The Permian Pangea. Phytogeographic implications of new palentological discoveries in Oman (Arabian Peninsula). *Comptes Rendus de l'Académie des Sciences*, Paris 321 (IIa) : 1069-1086.
- Chernyavsky G. V., Egorov A. I., Kharlov A. L., Redichkin N. A., Rosentuler A. M., Sharudo J. I. & Sokolov V. A. 1978. — Field excursion guidebook for the Donets basin and North Caucasus, in *Compte Rendu du 8^e Congrès international sur la stratigraphie du Carbonifère, Moscow, 1975*. Nauka, Moscow, 104 p.
- Chuvashov B. I. 1993. — General characteristics of the Permian deposits of the Urals, in Chuvashov B. I. & Nairn A. E. M. (eds), Permian system: guide to geological excursion in the uralian type localities, *Occasional Publications ESRI*, New Series 10: 3-23.
- Clayton G., Coquel R., Doubinger J., Gueinn K. J., Loboziak S., Owens B. & Streel M. 1977. — Carboniferous miospores of western Europe: illustration and zonation. *Mededelingen Rijks Geologische Dienst* 29: 1-71.
- Coquel R., Loboziak S., Owens B. & Teteriuk V. K. 1984. — Comparaison entre la distribution des principales microspores-guides du Namurien et du Westphalien en Europe occidentale et dans le bassin du Donetz (URSS) : 443-446, in *Compte Rendu du 9^e Congrès international de stratigraphie et de géologie du Carbonifère, Urbana 1979*, Volume 2.

- Coquel R., Doubinger J. & Massa D. 1988. — Nouvelles données palynologiques sur l'intervalle carbonifère Viséen/Moscovien bassin de Rhadamès (Libye). *Revue de l'Institut français du pétrole* 43 (1) : 3-16.
- Coquel R. & Rodriguez R. M. 1995. — Étude palynologique du Cantabrien et du Barruélien dans les régions de Barruelo et de Tejerina (Nord-ouest Espagne). *Sciences géologiques Bulletin*, Strasbourg 48 (1-3) : 49-61.
- Crowell J. C. 1995. — The ending of the late Paleozoic ice age during the Permian period: 62-74, in Scholle P. A., Peryt T. M. & Ulmer-Scholle D. S. (eds), *The Permian of Northern Pangea*, Volume 1: *Paleogeography, Paleoclimates, Stratigraphy*. Springer, Berlin.
- Davydov V. I. 1990. — Biostratigraphic zones of the Gzhelian in the Donbas and Pre-Donets basins according to the Fusulinids: 52-69, in Bogdanova J. N. & Bugrova E. M., *Problems of Modern Micropaleontology, Proceedings of the 34th Session of the All Union Paleontological Society, Baku, 1988* [in Russian].
- 1992. — Subdivision and correlation of Upper Carboniferous and Lower Permian deposits of the Donets Basin and Pre-Donets trough based on fusulinids. *Soviet Geology* 5: 53-61 [in Russian].
- 1996. — Fusulinid biostratigraphy and the correlation of Moscovian-Guadalupian North American, Tethyan and Boreal (Russian Platform/Uralian) standards. *Permian Newsletter of SCPS* 29: 47-51.
- Deleau P. 1951. — Les bassins houillers du Sud Oranais dans la région de Colomb-Béchar, Abadla. *Bulletin du Service de la Carte géologique d'Algérie* 2^e série, 20, 277 p.
- Desteuq C., Izart A. & Potherat P. 1988. — Étude sédimentologique du Carbonifère du bassin de Jerada (Maroc oriental). *Annales de la Société géologique du Nord* 107 : 203-210.
- Donsimoni M. 1981. — Synthèse géologique du bassin houiller lorrain. *Mémoire BRGM* 117, 100 p.
- Doubinger J. & Fabre J. 1983. — Mise en évidence d'un Autunien à flore mixte dans le bassin de Béchar-Abadla (Sahara occidental algérien). *Pollen et Spores* 25 (1) : 91-116.
- Doubinger J. & Roy-Dias C. 1985. — La paléoflore du Stéphanien de l'oued Zat (Haut Atlas de Marrakech, Maroc). *Geobios*, Lyon 18 : 573-586.
- 1986. — La paléoflore autuno-stéphanienne de la région de Demnate (Est de Marrakech, Maroc). *Geobios*, Lyon 19 (6) : 549-571.
- Doubinger J., Vetter P., Langiaux J., Galtier J. & Broutin J. 1995. — La flore fossile du bassin houiller de Saint-Étienne. *Mémoires du Muséum national d'Histoire naturelle*, Paris 164, 356 p.
- Eino O. L., Nemirovskaya T. I. & Solovieva M. N. (eds) 1996. — European part of the USSR: 23-180, in Wagner R. H., Winkler-Prins C. F. & Granados L. F. (eds), *The Carboniferous of the World*, Volume 3. Spanish geological Survey, Madrid.
- El Wartiti M., Broutin J. & Freytet P. 1986. — Premières découvertes paléontologiques dans les séries rouges carbonatées permienne du bassin de Tiddas (Maroc central). *Comptes Rendus de l'Académie des Sciences*, Paris 303 (IIa) : 263-268.
- El Wartiti M., Broutin J., Freytet P., Lahrib M. & Toutin-Morin N. 1990. — Continental deposits in Permian basins of the Mesetian Morocco. *Journal of African Earth-Sciences* 10 (1-2): 361-368.
- Fiebig H. E. R. 1969. — Das Namur C und Westphal im nieder-heinisch-westfälischen Steinkohlengbiet: 79-89, in *Compte Rendu du 6^e Congrès international de stratigraphie et de géologie du Carbonifère*, Sheffield 1967.
- Fissunen O. P. & Laveine J.-P. 1984. — Comparaison entre la distribution des principales espèces-guides végétales du Carbonifère moyen dans le bassin du Donetz (URSS) et les bassins du Nord-Pas de Calais et de Lorraine (France) : 95-100, in *Comptes Rendus du 9^e Congrès international de stratigraphie et de géologie du Carbonifère*, Urbana 1979, Volume 1.
- Geluk M. 1997. — Palaeogeographic maps of Moscovian and Arktinskian; contributions from the Netherlands, in Crasquin-Soleau S. & De Wever P. (eds), *Peri-Tethys: stratigraphic correlations*, *Geodiversitas* 19 (2) : 229-234.
- Ginkel A. C. van 1965. — Spanish Carboniferous fusulinids and their significance for correlations purposes. *Leide Geologische Mededelingen* 34: 137-225.
- Ginkel A. C. van & Villa E. 1996. — Palaeontological data of the San Emiliano formation (Cantabrian mountains, Spain) and their significance in the Carboniferous chronostratigraphy. *Geobios*, Lyon 29 (2) : 149-170.
- Görür N., Monod O., Okay A. I., Sengör C. A. M., Tüysüz O., Yigitbas E., Sakinc M. & Akkoc R. 1997. — Palaeogeographic and tectonic position of the Carboniferous rocks of the western Pontides (Turkey) in the frame of the Variscan belt. *Bulletin de la Société géologique de France* 168 (2) : 197-205.
- Granados L. F., Solovieva M. N., Reitlinger E. A. & Martinez-Diaz C. 1985. — The Bashkirian-Moscovian boundary problem in the Asturias (Northwest Spain): 27-34, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère*, Madrid, 1983, Volume 1. Instituto geológico y minero de España, Madrid.
- Hess J. C. & Lippolt H. J. 1986. — ⁴⁰Ar / ³⁹Ar ages of tonstein and tuff sandines: new calibration points for the improvement of the Upper Carboniferous time scale. *Isotope Geoscience* 59: 143-154.
- Inosova K. I., Kruzina A. H. & Schvatsman E. G. 1976. — *Atlas of Microspores and Pollens from the Upper Carboniferous and Lower Permian of the*

- Donets Basin*. Nedra, Moscow, 176 p. [in Russian].
- Iqbal N. 1993. — *Le Permien inférieur du Salt Range (Pakistan)*. Thèse, université de Nancy. I, Nancy, France, 227 p.
- Izart A. 1991. — Les bassins carbonifères de la Meseta marocaine, étude sédimentologique et approche du contexte structural. Part de la tectonique et de l'eustatisme. *Géologie méditerranéenne* 18 (1-2) : 61-72.
- Izart A., Briand C., Vaslet D., Vachard D., Coquel R. & Maslo A. 1996. — Stratigraphy and sequence stratigraphy of the Moscovian in the Donets Basin. *Tectonophysics* 268: 189-209.
- Izart A., Briand C., Vaslet D., Vachard D., Broutin J., Coquel R., Maslo A., Maslo N. & Kozitskaya R. 1998. — Stratigraphy and sequence stratigraphy of the Upper Carboniferous and Lower Permian in the Donets Basin, in Crasquin-Soleau S. & Barrier E. (eds), *Peri-Tethys Memoir 3: stratigraphy and evolution of Peri-Tethyan platforms*, *Mémoires du Muséum national d'Histoire naturelle*, Paris 177 : 9-33.
- Izart A., Kossovaya O., Vachard D. & Vaslet D. 1998. — Sequence stratigraphy of the Lower Permian along the Kosva river (Gubakha area, Central Urals, Russia). *Permian Newsletter of SCPS* 31: 17-20.
- Izart A. & Vachard D. 1994. — Subsidence tectonique, eustatisme et contrôle des séquences dans les bassins namuriens et westphaliens de l'Europe de l'ouest, de la CEI et des USA. *Bulletin de la Société géologique de France* 165 (5) : 499-514.
- Jenny J., Jenny-Deshusses C., Stampfli G. & Lys M. 1978. — La formation de Gheselghaleh, nouvelle unité lithologique du Carbonifère de l'Elbourz oriental (Iran). *Eclogae Geologiae Helveticae* 71/2 : 297-312.
- Jenny-Deshusses C. 1983. — *Le Permien de l'Elbourz central et oriental (Iran) : stratigraphie et micropaléontologie (Foraminifères et Algues)*. Thèse, université de Genève, Genève, Suisse.
- Josten K. H. 1991. — Die Steinkohlenfloren Nordwestdeutschlands. *Fortschrift Geologie Rhein und Westfalen* 32: 434 p.
- Kagarmanov A. K. & Donakova L. M. 1990. — *Decisions of the Regional Stratigraphic Conference on the Middle and Upper Paleozoic of the Russian Platform, Leningrad 1988, Carboniferous System*. VSEGEI, Leningrad, 92 figs [in Russian].
- Kerey I. E., Kelling G. & Wagner R. H. 1985. — An outline stratigraphy and palaeobotanical records from the middle Carboniferous rocks of northwestern Turkey. *Annales de la Société géologique du Nord* 105 : 203-216.
- Kora M. 1995. — Carboniferous macrofauna from Sinai, Egypt: biostratigraphy and palaeogeography. *Journal of African Earth-Sciences* 20: 37-51.
- Kossovaya O. 1996. — Correlations of Uppermost Carboniferous and Lower Permian Rugose Coral zones from the Urals to Western North America. *Palaios* 11: 71-82.
- Krainer K. 1992. — Facies, sedimentationprozesse und paläogeographie im Karbon der Ost- und Süd Alpen. *Jahrbuch Geologie B.A.* 135 (1): 99-193.
- Krainer K. 1993. — Late- and post-Variscan sediments of the eastern and southern Alps: 537-564, in Rainer J. F. von & Neubauer F. (eds), *Pre-Mesozoic Geology on the Alps*. Springer, Berlin.
- Kutzbach J. E. & Gallimore R. G. 1989. — Pangean climates: megamonsoons of the megacontinent. *Journal of Geophysical Research* 94: 3341-3357.
- Kutzbach J. E. & Ziegler A. M. 1993. — Simulation of the Late Permian climate and biomes with an asthenosphere-ocean model: comparisons with observations. *Philosophical Transactions of the Royal Society of London* 341 (B): 327-340.
- Laveine J.-P. 1987. — La flore du bassin houiller du Nord de la France, biostratigraphie et méthodologie. *Annales de la Société géologique du Nord* 106 : 87-94.
- Leyva F., Granados L. F., Solovieva M. N., Reitlinger M. N., Martinez-Diaz C., Laveine J.-P., Loboziak S., Brousmiche C., Candilier A. M. & Horvath V. 1985a. — La estratigrafia del Carbonifero medio en el sector central de la unidad estructural de la Sobia-Bodon (Cuencas de Quiros-Teverga): 213-230, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère, Madrid, 1983*, Volume 1. Instituto geológico y minero de España, Madrid.
- Leyva F., Granados L. F., Solovieva M. N., Laveine J.-P., Lys M., Loboziak S., Martinez-Diaz C., Brousmiche C., Candilier A. M. & Garcia Cortes A. 1985b. — La estratigrafia del Carbonifero medio en el area Los Tornos-Villoria-Colladona (sector oriental de Cuenca central): 231-248, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère, Madrid, 1983*, Volume 1. Instituto geológico y minero de España, Madrid.
- Leyva F., Granados L. F., Solovieva M. N., Laveine J.-P., Lys M., Loboziak S., Martinez-Diaz C., Brousmiche C., Candilier A. M., Garcia Cortes A. & Esnaola J. M. 1985c. — La estratigrafia del Carbonifero medio en la zona de Campo de Caso-Tanes (sector central de la region de Mantos): 249-268, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère, Madrid, 1983*, Volume 1. Instituto geológico y minero de España, Madrid.
- Love C. F. 1994. — The palynostratigraphy of the Hauski group (Westphalian-Artinskian) in Oman: 23-39, in Simmons M. D. (ed.), *Micropaleontology and Hydrocarbon Exploration in the Middle East. Arabia*. Chapman and Hall, London.
- Lützner H. (ed.) 1987. — *Sedimentary and volcanic Rotliegendes of the Saale-Depression. Symposium on Rotliegendes on Central Europe, Erfurt. Excursion*

- Guidebook*. Academy of Sciences of the GDR, Central Institute Physics Earth, 197 p.
- Lys M. 1988a. — Biostratigraphie du Carbonifère et du Permien en Mésogée. *Document BRGM* 147, 315 p.
- Lys M. 1988b. — Biostratigraphie des dépôts marins du Carbonifère et du Permien du Sud Tunisien ; micropaléontologie (foraminifères) et paléobiogéographie. *Bulletin du Centre de Recherche, Exploration et Production Elf-Aquitaine*, Pau 12 (2) : 601-659.
- Makhlina M. X., Koulikova A. M. & Nikitina N. A. 1979. — Structure, Biostratigraphy and Paleogeography of the upper Carboniferous in the Moscow syncline, in Makhlina M. X. & Chik E. M. (eds), *Stratigraphy, Paleontology and Paleogeography of the Carboniferous of the Moscow syncline, Trudy Mineralogy and Geology of Russia*: 25-69 [in Russian].
- Makhlina M. X., Solovieva M. N. & Goreva N. V. 1997. — On the biostratigraphy and cyclostratigraphy of the Moscovian stage in the type area, in Crasquin-Soleau S. & De Wever P. (eds), *Peri-Tethys: stratigraphic correlations, Geodiversitas* 19 (2) : 221-228.
- Martinez Diaz C., Granados L. F., Leyva F., Laveine J.-P., Solovieva M. N., Reitlinger E. A., Gervilla M., Loboziak S., Brousmiche C., Candilier A. M., Pendas F. & Horvath V. 1985. — Aportaciones a la cronoestratigrafía del Carbonífero medio de Asturias y nuevos datos para un intento de correlación de las escalas marinas y continentales: 269-280, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère, Madrid, 1983*, Volume 1. Instituto geológico y minero de España, Madrid.
- Martinez-Garcia E. 1991. — Hercynian syn-orogenic and post-orogenic successions in the Cantabrian and Palentian zones (North West Spain): comparison with the western European occurrences. *Giornale di Geologia* 3 (53-1): 209-228.
- Massa D. & Vachard D. 1979. — Le Carbonifère de Libye occidentale. Biostratigraphie et Micropaléontologie. Position dans le domaine téthysien d'Afrique du Nord. *Revue de l'Institut français du Pétrole* 31 (1) : 3-65.
- Massa D., Coquel R., Loboziak S. & Taugourdeau-Lantz J. 1980. — Essai de synthèse stratigraphique et palynologique du Carbonifère en Libye occidentale. *Annales de la Société géologique du Nord* 99 : 429-442.
- Menning M. 1995. — A numerical time scale for the Permian and Triassic periods: an integrated time analysis: 77-97, in Scholle P. A., Peryt T. M. & Ulmer-Scholle D. S. (eds), *The Permian of Northern Pangea*, Volume 1: *Paleogeography, Paleoclimates, Stratigraphy*. Springer, Berlin.
- Monod O. 1977. — *Recherches géologiques dans le Taurus occidental au Sud de Beyşehir (Turquie)*. Thèse, université Paris-Sud, Orsay, France, 442 p.
- Nedjari A. 1982. — Les formations Permo-Carbonifères du bassin du Mezarif-Nekheila (Sud-Ouest Oranais, Algérie), études stratigraphique et sédimentologique, approche géochimique. *Bulletin de la Sous-Direction de la Géologie d'Algérie* 49 : 132 p.
- Owens B. & Turner N. 1995. — Late Westphalian palynomorphs from northern Saudi Arabia. *Review of Paleobotany and Palynology* 89: 125-137.
- Owodenko B. 1976. — Le bassin houiller de Jerada (Maroc oriental). Essai de synthèse et de paléogéographie. *Notes et Mémoires du Service géologique du Maroc* 207bis, 148 p.
- Parrish J. T. 1995. — Geologic evidence of permian climate: 54-61, in Scholle P. A., Peryt T. M. & Ulmer-Scholle D. S. (eds), *The Permian of Northern Pangea*, Volume 1: *Paleogeography, Paleoclimates, Stratigraphy*. Springer, Berlin.
- Pasini M. & Vai G. B. 1997. — Review and updating of the Moscovian to Artinskian marine rocks in peninsular Italy, in Crasquin-Soleau S. & De Wever P. (eds), *Peri-Tethys: stratigraphic correlations, Geodiversitas* 19 (2) : 187-191.
- Ramsbottom W. H., Calver M. A., Eagar R. M. C., Hodson F., Holliday D. W., Stubblefield C. J. & Wilson R. B., 1978. — A correlation of Silesian Rocks in the British Isles. *Special Report – Geological Society of London* 10, 81 p.
- Ross C. A. & Ross J. R. P. 1987. — Late paleozoic sea levels and depositional sequences. *Cushman Foundation for Foraminiferal Research Special Publication* 24: 137-149.
- 1994. — Permian Sequence Stratigraphy and fossil Zonation, in Embry A. F., Beauchamp B. & Glass D. (eds), *Pangea: Global Environments and Resources, Memoir – Canadian Society of Petroleum Geologists* 17: 219-231.
- 1995. — Permian Sequence Stratigraphy: 98-123, in Scholle P. A., Peryt T. M. & Ulmer-Scholle D. S. (eds), *The Permian of Northern Pangea*, Volume 1: *Paleogeography, Paleoclimates, Stratigraphy*. Springer, Berlin.
- Saenz De Santa Maria J. A., Luque C., Gervilla M., Laveine J.-P., Loboziak S., Brousmiche C., Coquel R. & Martinez-Diaz C. 1985. — Aportacion al conocimiento estratigrafico y sedimentologico del Carbonifero productivo de la cuenca central Asturiana: 303-326, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère, Madrid, 1983*, Volume 1. Instituto geológico y minero de España, Madrid.
- Samankassou E. 1997. — Cyclic sedimentation patterns of the lower Pseudoschwagerina limestone (latest Carboniferous, Carnic Alps). Abstracts: 127, in *Compte Rendu du 13^e Congrès international de stratigraphie et de géologie du Carbonifère et du Permien, Krakow 1995*.
- Schönlaub H. P. 1992. — Stratigraphy, Biogeography, and Paleoclimatology of the Alpine Paleozoic and its implications for plate movements. *Jahrbuch Geologie B. A.* 135 (1): 381-418.


- Schneider J. W. & Gebhard U. 1993. — Litho- und Biofaziesmuster in intra- und extramontanen Senken des Rotliegend (Perm, Nord- und Ostdeutschland). *Geologisches Jahrbuch A* 127: 57-98.
- Schneider J. W., Rössler R. R. & Gaitsch B. G. 1994. — Time lines of the Late Variscan volcanism – a holostatigraphic synthesis. *Zeitblatt für Geologie und Paläontologie* 1: 477-490.
- Schneider J. W., Rössler R. R. & Gaitsch B. G. 1995. — Proposal for a combined reference section of the central european continental Carboniferous and Permian for correlations with marine standard sections. *Permian Newsletter of SCPS* 26: 26-32.
- Schneider J. W. 1996. — Biostratigraphie des kontinentalen Oberkarbon und Perm im Thüringer Wald, SW-Saale-Senke-Stand und Probleme. *Beiträge zur Geologie von Thüringen Neue Folge* 3: 121-151.
- Schneider J. W. & Rössler R. R. in press. — Stratigraphie des Oberkarbons: Biostratigraphie der Rotfolgen. *DGMK-Forschungsbericht*, Hamburg 459 (3/3): 66 p.
- Scotese C. R. & Langford R. P. 1995. — Pangea and the Paleogeography of the Permian: 3-19, in Scholle P. A., Peryt T. M. & Ulmer-Scholle D. S. (eds), *the Permian of Northern Pangea*, Volume 1: *Paleogeography, Paleoclimates, Stratigraphy*. Springer, Berlin.
- Scotese C. R. & McKerrow W. S. 1990. — Revised world maps and introduction, in McKerrow W. S. & Scotese C. R. (eds), *Paleozoic paleogeography and biogeography*, *Memoir – Geological Society of London* 12: 1-21.
- Sinitsyn I. I., Sinitsyna Z. A., Aleksandrov V. A., Furdij R. S., Shamov D. F. & Vissarionova A. Ya. 1978. — Field excursion guidebook for the Carboniferous sections of South Urals (Bashkiria), in *Compte Rendu du 8^e Congrès international sur la stratigraphie du Carbonifère*, Moscow, 1975. Nauka, Moscow, 183 p.
- Solovieva M. N. 1985. — Correction of the USSR general Carboniferous scale in connection with restudying the Moskovian stage stratotype and a new model for the correlation of the lower Moskovian substage: 21-26, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère*, Madrid, 1983, Volume 1. Instituto geológico y minero de España, Madrid.
- Solovieva M. N., Chizhova V. A., Einor O. L., Grigorieva A. D., Reitlinger E. A. & Vdovenko M. V. 1985a. — Review of recent data on the carboniferous stratigraphy of the USSR: 3-10, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère*, Madrid, 1983, Volume 1. Instituto geológico y minero de España, Madrid.
- Solovieva M. N., Fisunen O. P., Goreva N. V., Barskov I. S., Gubareva I. S., Dzhenchuraeva A. V., Dalmatskaia I. I., Ivanova E. A., Poletaev V. I., Popov A. V., Rummyantseva Z. S., Teteruk V. K. & Shik E. M. 1985b. — New data on stratigraphy of the Moskovian stage: 11-20, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère*, Madrid, 1983, Volume 1. Instituto geológico y minero de España, Madrid.
- Stapf K. G. 1997. — Rotliegend lacustrine sediments of the Saar-Nahe Basin. Excursion A2, 18th IAS meeting. *Gaia heidelbergensis* 4: 15-28.
- Stschegolev A. K. & Kozitskaya R. I. 1984. — The paleontological basis of the project of the Upper Carboniferous standard scale in Europe and central Asia, in Menner V. V. & Grigorieva A. D. (eds), *Upper Carboniferous of the USSR, Transactions of Stratigraphical Committee of USSR*, Moscow 13: 107-113 [in Russian].
- Vachard D., Massa D. & Strank A. 1993. — Le Carbonifère du sondage A1-37 (Cyrénaïque, Libye), analyse biostratigraphique, conséquences paléogéographiques. *Revue de Micropaléontologie* 36 (2): 165-186.
- Vachard D. & Maslo A. 1996. — Précisions biostratigraphiques et micropaléontologiques sur le Bashkirien d'Ukraine (Carbonifère moyen). *Revue de Paléobiologie* 15: 357-383.
- Vai G. B. 1991. — Palaeozoic strike-slip rift pulses and palaeogeography in the circum-mediterranean tethan realm. *Palaeogeography, Palaeoclimatology, Palaeoecology* 87: 223-252.
- 1994. — Crustal evolution and basement elements in the Italian area: palaeogeography and characterization. *Bollettino di Geofisica Teorica ed Applicata* 36 (141-144): 411-434.
- Vai G. B. & Venturini C. 1997. — Moscovian and Artinskian rocks in the frame of the cyclic Permo-Carboniferous deposits of the Carnic Alps and related areas, in S. Crasquin-Soleau S. & De Wever P., (eds), *Peri-Tethys: stratigraphic correlations*, *Geodiversitas* 19 (2): 221-228.
- Villa E. 1985. — Foraminiferos de la region oriental de Asturias (Cordillera Cantabrica, N de España): 333-344, in *Compte Rendu du 10^e Congrès international de stratigraphie et de géologie du Carbonifère*, Madrid, 1983, Volume 1. Instituto geológico y minero de España, Madrid.
- Villa E., Davydov V. I., Konovalova M. V. & Remizova S. 1993. — Studies on the base of the *Protriticites* zone. A report on SCCS project group 5. *Annales de la Société géologique de Belgique* 116: 323-331.
- Wagner R. H. & Alvarez-Vasquez C. 1991. — Floral characterization and biozones of the Westphalian D Stage in NW Spain. *Neues Jahrbuch Geologie und Paläontologie* 183: 171-202.
- Wagner R. H. & Bowman M. B. J. 1983. — The position of the Bashkirian/Moskovian boundary in West European chronostratigraphy. *Newsletter Stratigraphy*, Berlin 12: 132-161.
- Wagner R. H. & Winkler-Prins C. F. 1985. — The

- Cantabrian and Barruelian stratotypes: a summary of basin development and biostratigraphic information, in Lemos De Sousa M. J. & Wagner R. H. (eds), Papers on the Carboniferous of the Iberian Peninsula, *Annale Facultad Ciencias*, Porto suppl. 64: 359-410.
- 1991. — Major subdivisions of the Carboniferous system: 213-245, in *Compte Rendu du 11^e Congrès international de stratigraphie et de géologie du Carbonifère, Beijing 1987*, Volume 1.
- 1993. — General overview of the Carboniferous stratigraphy. *Annales de la Société géologique de Belgique* 116 (1) : 163-174.
- Yablokov V. S., Leonenko V. S., Mikhailova E. V., Belskaya T. N. & Reitlinger E. A. 1978. — Field excursion guidebook for the Carboniferous sequences of the Moscow basin, in *Compte Rendu du 8^e Congrès international sur la stratigraphie du Carbonifère, Moscow, 1975*. Nauka, Moscow, 176 p.
- Yugan J. 1996. — A global chronostratigraphic scheme for the Permian system, two decades of the Permian subcommission. *Permophiles Newsletter of SCPS* 28: 4-9.
- Yugan J., Wardlaw B. R., Glenister B. F. & Kotlyar G. V. 1997. — Permian chronostratigraphic subdivisions. *Episodes* 20 (1): 10-15.
- Zdanowski A. & Zakowa H. (eds) 1995. — The Carboniferous system in Poland. *Prace Państwowego Instytutu Geologicznego*, Warszawa 148, 215 p.
- Zhamoyda A. I. (ed.) in press. — Information on the decisions of the commission on the Carboniferous Stratigraphy. *Russian interdepartemental Stratigraphic Committee*, Saint-Petersburg issue 29 [in Russian].
- Ziegler A. M. 1990. — Phytogeographic patterns and continental configurations during the Permian period, in McKerrow W. S. & Scotese C. R. (eds), Paleozoic paleogeography and biogeography, *Memoir – Geological Society of London* 12: 363-379.
- Ziegler A. M., Hulver M. L. & Rowley D. B. 1997. — Permian world topography and climate: 111-146, in Martini I. P. (ed.), *Late Glacial and Postglacial Environmental Changes, Quaternary, Carboniferous-Permian and Proterozoic*. Oxford University Press, Oxford.

*Submitted for publication on 30 April 1997;
accepted on 15 December 1997.*

APPENDIX 1

Late carboniferous and Early Permian logs and tables.
Northern Peri-Tethyan Basins (logs 1-36); Tethyan Basins (logs 37-49);
Southern Peri-Tethyan Basins (logs 50-62).




MAP. — Location of the logs or sections presented in the appendix 1. Map realized by the Peri-Tethys cartographic team. Pakistan table (62) is not located on the map. 1, log or section; 2, synthetic log or section.

LEGEND

	Clayey Limestone		Gypsum
	Bioclastic Limestone		Potash
	Limestone		Halite
	Limestone		Coal with paleosol
	Reef Limestone		Coal
	Outer Platform Limestone		Marine Claystone
	Limestone with chert		Marine Claystone
	Limestone with chert		Fluvial or deltaic Claystone
	Nodular Limestone		Claystone
	Pebbly Limestone		Silty Claystone
	Sandy Limestone		Sandy Claystone
	Primary Dolomite		Siltstone
	Secondary Dolomite		Turbidites
	Dolomite		Fluvial Sandstone
	Clayey Dolomite		Conglomerate
	Accurate location of a fossil in the log		Conglomerate
	Red colour of the sediments known with certainty located at the left side of the column		Breccia
			Volcanic Rocks

LATE CARBONIFEROUS OF THE SAINT-ÉTIENNE BASIN




LOG. 1. — Late Carboniferous of the Saint-Étienne Basin, coordinated by Mercier and Broutin. 1, Mercier (this paper); 2, Broutin (this paper). Saint-Étienne, France, 4°24'E - 45°27'N; Firminy, ex-SW, 4°15'E - 45°24'N; La Fouillouse, ex-NW, 4°19'E - 45°30'N; W Saint-Chamond, 4°30'E - 45°29'N; Rive-de-Gier, ex-NE, 4°37'E - 45°31'N.

AUTUNIAN OF THE AUTUN BASIN


Autun	Formations	Beds	Macroflora	Palynomorphs	Western European Stages			
 125 m 0	Millery	Les Télots bituminous beds (10 beds)	<i>Rhachiphyllum schenkii</i> <i>Rhachiphyllum pellatii</i> <i>Lodevia bibractensis</i>		UPPER AUTUNIAN	A U T U N I A N		
	Surmoulin	Surmoulin bituminous bed	<i>Gracilopteris raymondii</i> <i>Baiera raymondii</i>					
	Muse	Muse bituminous bed Lally bituminous bed	<i>Autunia conferta</i> <i>Autunia naumanii</i> <i>Walchia piniformis</i> <i>W. goeppertiana</i>	<i>Disaccites striatiti</i> <i>Vittatina</i> spp. <i>Disaccites non-striati</i>	LOWER AUTUNIAN			
			<i>Culmitschia frondosa</i>	<i>Potonieisporites novicus-bhardwaji</i> complex <i>Vesicaspora</i> spp.				
			<i>Sphenophyllum angustifolium</i> <i>Alethopteris zeilleri</i>	<i>Vittatina costabilis</i>				
	Igornay	Igornay bituminous bed Moloy bituminous bed	<i>Pecopteris plumosa</i> <i>Odontopteris brardii</i>	<i>Florinites</i> spp. <i>Monoletes</i> spores <i>Trilete</i> spores	UPPER STEPHANIAN			
	Epinac							
	BASEMENT							

Log. 2. — Autunian of the Autun Basin, coordinated by Broutin, after Broutin *et al.* (1996). Autun, France, 4°30'E - 47°N.

LATE CARBONIFEROUS OF THE LORRAINE BASIN




LOG. 3. — Late Carboniferous of the Lorraine Basin, coordinated by Donsimoni, Laveine and Coquel, Sainte-Fontaine, 6°48'E - 49°30'N. 1, Donsimoni (this paper); 2, Laveine (this paper); 3, Coquel (this paper).




LOG. 4. — Late Carboniferous of the Lorraine Basin, coordinated by Donsimoni, Laveine and Coquel, Faulquemont, 6°38'E - 49°02'E. 1, Donsimoni (this paper); 2, Laveine (this paper); 3, Coquel (this paper).

LATE CARBONIFEROUS OF THE LORRAINE BASIN




LOG. 5. — Late Carboniferous of the Lorraine Basin, coordinated by Donsimoni, Laveine and Coquel, La Houve, 6°38'E - 48°45'N. 1, Donsimoni (this paper); 2, Laveine (this paper); 3, Coquel (this paper).



LOG. 6. — Westphalian C and D of the North France Basin, coordinated by Izart, Laveine and Coquel. 1, Izart after Bourroz *et al.* (1964); 2, Laveine (this paper); 3, Coquel (this paper). Hénin, North France, 3°E - 50°30'N.

THE NETHERLANDS COAL BASIN



LOG. 7. — The Netherlands coal Basin, coordinated by Gelük, after Gelük (1997). Tubbergen-8, 6°53'E - 52°26'N.

LATE CARBONIFEROUS OF THE NORTHERN ENGLAND BASIN

Staffordshire	Formations	Marine bands	Coal seams	Goniatites	Pelecypods Esteriids	Macroflora	Palynomorphs	Western European Stages
	Clent							AUTUNIAN
	Enville							STEPHANIAN
	Keele							
	Newcastle							
	Etruria						<i>Thymospora obscura</i>	WESTPHALIAN D
		Cambriense		<i>Anthracoceas cambriense</i>	<i>A. phillipsii</i>		<i>T. securis</i>	
		Shotton						
		Edmondia	Highman		<i>Upper similis- pulchra</i>		<i>Vestispora magna</i>	WESTPHALIAN C
80 m 0	Coal Measures	Aegir	Wales	<i>Anthracoceas aegiranum</i>				

WESTPHALIAN B

Log. 8 — Late Carboniferous of the Northern England Basin, coordinated by Izart, Izart (this paper) after Ramsbottom *et al.* (1978). Stafford, North England, 2°W - 52°50'N.


WESTPHALIAN OF THE RUHR COAL BASIN

Ruhr	Marine bands	Coal seams	Goniatites	Pelecypods Esteriids	Macroflora	Palynomorphs	Western European Stages
		Xanten Walkür Volker Undine		<i>A. tenuis</i>	<i>Alethopteris michaui</i>		
		Tristan Siegfried Rubezahl					
		Parsifal Odin			<i>Alethopteris grandini</i>		
		Niebelung			<i>Mariopteris robusta</i>		
		Midgard Loki Kobold Iduna Hagen Gudrun			<i>Neuropteris chalardi</i> <i>Linopteris regniezi</i>		WESTPHALIAN C
		Freya Erda Chriemhilt Baldur	<i>Anthracoceas aegiranum</i>	<i>A. phillipsii</i>	<i>Reticulopteris münsteri</i>	<i>Torispora securis</i>	
80 m 0	Aegir						

WESTPHALIAN B


Log. 9. — Germany Basins. Westphalian of the Ruhr coal Basin, coordinated by Süß after Fiebig (1969) and Josten (1991). Lippermulde 1, Ruhr, Germany, 6°52'E - 51°37'N.

WESTPHALIAN D OF IBSENBUHREN/OSNABRÜCK AREA (GERMANY)




LOG. 10. — Germany Basins. Westphalian of the Ibbenbüren/Osnabrück area coordinated by Süss, after Josten (1991). Ibbenbüren, Osnabrück, 8°E - 52°20'N.

STEPHANIAN AND AUTUNIAN OF THE SAALE (EASTERN GERMANY) BASIN




Log. 11. — Germany Basins. Stephanian and Autunian of the Saale (eastern Germany) Basin, coordinated by Izart, Izart (this paper) after Schneider & Rossler (in press); 1, Broutin (this paper); 2, Lützner (1987). Halle, NE Saale, 12°E - 51°30'N; Tambach, SW Saale, 10°36'E - 50°48'N.

LATE CARBONIFEROUS OF THE UPPER SILESIA BASIN



LOG. 12. — Poland Basins, coordinated by Izart. Late Carboniferous of the upper Silesian Basin, Izart (this paper) after Zdanowski & Zakowa (1995). Katowice, Poland, 19°E - 50°15'N.

LATE CARBONIFEROUS OF THE LUBLIN BASIN





LOG. 13. — Poland Basins, coordinated by Izart. Late Carboniferous of the Lublin Basin, Izart (this paper) after Zdanowski & Zakowa (1995). Lublin, Poland, 22°45'E - 51°15'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE LOWER SILESIAN BASIN CZECH REPUBLIC

Brou-1	Formations (1)	Macroflora (1)	Palynomorphs (1)	Western European Stages (1) (2)		
	Broumov	<i>Autunia conferta</i> <i>Autunia naumanii</i> <i>Walchia piniformis</i>	<i>Potonieisporites novicus</i> <i>Florinites</i>	AUTUNIAN	AUTUNIAN	
	Chvalec	<i>Autunia naumanii</i> <i>Alethopteris zeilleri</i> <i>Odontopteris brardii</i>	<i>Cadiospora magna</i> <i>Angulisporites splendidus</i>	C	UPPER STEPHANIAN	
		Odolov	<i>Sphenophyllum oblongifolium</i> <i>Sphenophyllum thoni</i> v. minor	<i>Potonieisporites novicus</i> <i>Verrucosisporites grandiverrucosus</i>		STEPHANIAN B
			<i>Sphenophyllum emarginatum</i> <i>Pseudomariopteris ribeyroni</i>	<i>Vestispora fenestrata</i> <i>Vestispora quaesita</i>	A	LOWER STEPHANIAN
	Zacler	<i>Sphenophyllum myriophyllum</i> <i>Sphenophyllum cuneifolium</i>	<i>Torispora securis</i> <i>Westphalensisporites striatus</i>	CANTABRIAN	WESTPHALIAN D	
		<i>Laveineopteris tenuifolia</i> <i>Paripteris linguaeifolia</i>	<i>Dictyotrites bireticulatus</i> <i>Savistrisporites cantotus</i>			
	100m 0				WESTPHALIAN B	


LOG. 14. — Czech Republic Basins, coordinated by Oplustil, Pesek, Martinek, Simunek & Drabkova. Late Carboniferous and Early Permian of the lower Silesian Basin. 1, Oplustil *et al.* (this volume); 2, this paper. Broumov, Czech Republic, 16°21'E - 50°35'N.

LATE CARBONIFEROUS IN THE CENTRAL AND WESTERN BOHEMIA

OB-1	Formations		Macroflora	Palynomorphs	Western European Stages	
					(1)	(2)
	Line		<i>Sphenophyllum angustifolium</i> <i>Sphenophyllum thoni</i> v. <i>minor</i> <i>Pecopteris densifolia</i> <i>Ernestiodendron filiciforme</i> <i>Callipteridium pteridium</i> <i>Odontopteris brardii</i> <i>Odontopteris osmundaeformis</i>		C	UPPER
	Slany	Otruby	<i>Sphenophyllum oblongifolium</i> <i>Sphenophyllum longifolium</i> <i>Sphenophyllum thoni</i> v. <i>minor</i> <i>Pseudomariopteris ribeyroni</i> <i>Odontopteris intermedia</i>	<i>Granulatisporites granifer</i>	STEPHANIAN B	STEPHANIAN
		Malesice		<i>Verrucosisporites grandiverrucosus</i>		
		Jelenice		<i>Kosankeisporites elegans</i>		
	Tynec		<i>Sigillaria brardii</i> <i>Sphenophyllum thoni</i> v. <i>minor</i> <i>Nemejcopteris feminaeformis</i>		A	LOWER STEPHANIAN
	Nyrani		<i>Sphenophyllum emarginatum</i> <i>Dicksonites pluckenetii</i> <i>Ptychocarpus unitus</i>	<i>Vestispora fenestrata</i> <i>Vestispora quaesita</i>	CANTABRIAN	WESTPHALIAN D
			<i>Praecallipteridium rubescens</i> <i>Sphenophyllum emarginatum</i>	<i>Torispora securis</i> <i>Punctatosporites minutus</i> <i>Microreticulosporites nobilis</i> <i>Laevigatosporites minimus</i>		
	Kladno					
	Radnice		<i>Laveineopteris tenuifolia</i> <i>Sphenophyllum cuneifolium</i>	<i>Knoxisporites pustulatus</i> <i>Dictyotritetes</i>	WESTPHALIAN C	BASEMENT

LOG. 15. — Czech Republic Basins, coordinated by Oplustil, Pesek, Martinek, Simunek & Drabkova. Late Carboniferous in the central and western Bohemian. 1, Oplustil *et al.* (this volume); 2, this paper. Otruby, Czech Republic, 14°06'E - 50°15'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE BOSKOVICE FURROW




Log. 16. — Czech Republic Basins, coordinated by Oplustil, Pesek, Martinek, Simunek & Drabkova. Late Carboniferous and Early Permian of the Boskovice Furrow, after Oplustil *et al.* (this volume). Boskowitz, Czech Republic, 16°25'E - 49°10'N.


CARBONIFEROUS AND EARLY PERMIAN OF THE KRKONOSE BASIN

Pé-1	Formations	Macroflora	Palynomorphs	Western European Stages	
				(1)	(2)
	Proseč	<i>Autunia conferta</i>	<i>Vesicaspora</i> spp. <i>Vittatina costabilis</i> <i>Gardenaisporites glomus</i> <i>Knoxisporites glomus</i> <i>Endosporites formosus</i>	AUTUNIAN	AUTUNIAN
	Vrchlabí	<i>Autunia conferta</i> <i>Ernestiodendron filiciforme</i>	<i>Potonieisporites novicus</i> <i>Vittatina costabilis</i> <i>Protohaploxipinus</i> spp. <i>Hamia pollinites</i>		
	Semily	<i>Ernestiodendron filiciforme</i> <i>Callipteris pteridium</i> <i>Alethopteris zeilleri</i> <i>Odontopteris osmudaeformis</i>	<i>Cadiopora magna</i> <i>Spinisporites spinosus</i> <i>Potonieisporites novicus</i>	C	UPPER STEPHANIAN
	Syřenov	<i>Pecopteris unita</i> <i>Annularia stellata</i> <i>Sphenophyllum emarginatum</i> <i>Sphenophyllum oblongifolium</i>	<i>Cadiospora magna</i> <i>Endosporites formosus</i> <i>Verrucosisporites sinensis</i> <i>Gillespieisporites</i> spp.	B	
	Kumburk	<i>Alethopteris</i> sp. <i>Pecopteridium costei</i> <i>Nemejcopteris feminaeformis</i>	No microflora	A CANTABRIAN	LOWER STEPHANIAN
				WESTPHALIAN D	

Log. 17. — Czech Republic Basins, coordinated by Oplustil, Pesek, Martinek, Simunek & Drabkova. Carboniferous and Early Permian of the Krkonose Basin. 1, Martinek *et al.* this volume; 2, this paper. Proseč, Czech Republic, 15°41'E - 50°34'N.




LOG. 18. — Czech Republic Basins, coordinated by Oplustil, Pesek, Martinek, Simunek & Drabkova. Late Carboniferous and Early Permian of the Krusne hory Mountains, after Oplustil *et al.* (this volume). Krusne Hory, Czech Republic, 13°24'E - 50°37'N.




LOG. 19. — Slovak Republic Basins, coordinated by Vozarova. Moscovian of the northern Gemic unit (Slovak Republic), after Vozarova (this paper). Northern Gemic, Slovak Republic, 20°30' E - 48°55' N.

AUTUNIAN AND SAXONIAN OF THE NORTHERN GEMERIC UNIT



LOG. 20. — Slovak Republic Basins, coordinated by Vozarova. Autunian and Saxonian of the northern Gemic unit (Slovak Republic), after Vozarova (this paper). North Gemic, Slovak Republic, 20°30'E - 48°55'N.

LATE CARBONIFEROUS OF THE ZEMPLINIC UNIT


Zemplinic	Formations	Macroflora	Palynomorphs	Western European Stages
	Kasov	<i>Dadoxylon</i> sp.	<i>Columinisporites ovalis</i> <i>Thymospora perrucosa</i> <i>Vestispora fenestrata</i> <i>Cyclogranisporites pergranulus</i>	C
	Trna	<i>Stigmaria ficoides</i> <i>Pecopteris cyathea</i> <i>Cordaites borassifolius</i> <i>Asterotheca arborescens</i> <i>Alethopteris bohémica</i> <i>Sphenophyllum oblongifolium</i> <i>Annularia pseudostellata</i>	<i>Columinisporites</i> <i>Lycospora granulata</i> <i>Torispora laevigata</i> <i>Dendosporites plicatus</i> <i>Endosporites</i> div. sp. <i>Torispora securis</i> <i>Thymospora thiesseii</i> <i>Thymospora perrucosa</i>	STEPHANIAN B
	Luhyna	<i>Calamites cistii</i> <i>Pecopteris cf. miltonii</i> <i>Asterophyllites trichomatosus</i>	<i>Lycospora pusilla</i> <i>Laevigatosporites</i> sp. <i>Dendosporites</i> div. sp.	A
100m 0	Cerhov		<i>Triquitres tricuspis</i> <i>Microreticulatisporites sulcatus</i> <i>Cyrratiradites trizonarius</i>	WESTPHALIAN C-D

LOG. 21. — Slovak Republic Basins, coordinated by Vozarova. Late Carboniferous of the Zemplinic unit (Slovak Republic), after Vozarova (this paper). Zemplinic, Slovak Republic, 21°44'E - 48°30'N.

AUTUNIAN AND SAXONIAN OF THE ZEMPLINIC UNIT


Zemplinic	Formations	Macroflora	Palynomorphs	Western European Stages
	Cernochov		<i>Lueckisporites virkkiae</i> <i>Striatites richteri</i>	
			<i>Lueckisporites virkkiae</i> <i>Jugasporites delasauei</i> <i>Limitisporites rectus</i> <i>Lunatisporites div. sp.</i> <i>Klausipollenites schaubegeri</i> <i>Falcisporites zapfei</i>	THURINGIAN ?
	Cejkov		<i>Vittatina div. sp.</i> <i>Florinites div. sp.</i>	SAXONIAN
			<i>Potonieisporites radius</i> <i>Potonieisporites novicus</i> <i>Nuskoisporites dulhuntyi</i>	AUTUNIAN
25 m 0				

Log. 22. — Slovak Republic Basins, coordinated by Vozarova. Autunian and Saxonian of the Zemplinic unit (Slovak Republic), after Vozarova (this paper). Zemplinic, Slovak Republic, 21°44'E - 48°30'N.




LOG. 23. — Slovak Republic Basins, coordinated by Vozarova. Stephanian of the Hronic unit (Slovak Republic), after Vozarova (this paper). Hronic, Slovak Republic, 19°58'E - 49°03'N.

GEODIVERSITAS • 1998 • 20 (4)




AUTUNIAN AND SAXONIAN OF THE SOUTHERN GEMERIC UNIT

S Gemic	Formations	Macroflora	Palynomorphs	Western European Stages
	Roznava			
			<i>Potonieisporites novicus</i>	
			<i>Striatodisaccites</i>	
			<i>Gnetaceapollenites</i>	
			<i>Vittatina sp.</i>	SAXONIAN
			<i>Vittatina costabilis</i>	-
			<i>Leiotrilites infragranulatus</i>	AUTUNIAN
			<i>Triquitrites additus</i>	
			<i>Reticulatisporites reticulocingulum</i>	


LOG. 25. — Slovak Republic Basins, coordinated by Vozarova. Autunian and Saxonian of the southern Gemic unit, after Vozarova (this paper). Southern Gemic, Slovak Republic, 20°18'E - 48°40'N.

LATE CARBONIFEROUS OF THE NORTHERN TURKEY BASIN




LOG. 26. — Late Carboniferous of the Northern Turkey Basin, coordinated by Izart, Izart (this paper) after Görür *et al.* (1997) and Kerey *et al.* (1985). Zonguldak, North Turkey, 31°30'E - 41°15'N.

LATE CARBONIFEROUS OF THE CAUCASUS BASIN




LOG. 27. — Late Carboniferous of the Caucasus Basin, coordinated by Izart, Izart (this paper) after Chernyavsky *et al.* (1978). Labinskoe, Russia, 42°E - 44°N.

VEREIAN AND KASHIRIAN OF THE DONETS BASIN




LOG. 28. — Donets Basin (Ukraine), coordinated by Izart et al. (1996, 1998). Vereian and Kashirian of the Donets Basin. 1, Izart et al. (1996); 2, Vachard & Maslo (this paper); 3, Kozitkaya (this paper); 4, Laveine (this paper); 5, Coquel (this paper). Boreholes 771/849, 38°30'E - 48°12'N.


PODOLSKIAN AND MYACHKOVIAN OF THE DONETS BASIN




LOG. 29. — Donets Basin (Ukraine), coordinated by Izart et al. (1996, 1998). Podolskian and Myachkovian of the Donets Basin. 1, Izart et al. (1996); 2, Vachard & Maslo (this paper); 3, Kozitkaya (this paper); 4, Laveine (this paper); 5, Coquel (this paper). Boreholes 771/849, 38°30'E - 48°12'N.




LOG. 30. — Donets Basin (Ukraine), coordinated by Izart *et al.* (1996, 1998). Kasimovian of the Donets Basin. 1, Izart *et al.* (1996); 2, Vachard & Maslo (this paper); 3, Kozitkaya (this paper); 4, Stchegolev (this paper); 5, Broutin (this paper). Kalinovo cross-section, 38°48'E - 48°36'N.




LOG. 31. — Donets Basin (Ukraine), coordinated by Izart *et al.* (1996, 1998). Gzhelian and Orenburgian of the Donets Basin. 1, Izart *et al.* (1996); 2, Vachard & Maslo (this paper); 3, Kozitkaya (this paper); 4, Stchegolev (this paper); 5, Broutin (this paper). Kalinovo cross-section, 38°48'E - 48°36'N.



LOG. 32. — Donets Basin (Ukraine), coordinated by Izart *et al.* (1996, 1998). Early Permian of the Donets Basin. 1, Izart *et al.* (1996); 2, Vachard & Maslo (this paper); 3, Stchegolev (this paper); 4, Broutin (this paper). Ivanovo-Darievka cross-section, 37°30'E - 48°30'N.



LOG. 33. — Moscow Basin, coordinated by Briand *et al.* Moscowian of the Moscow Basin. 1, Briand *et al.* (1998); 2, Makhliina *et al.* (1997); 3, Goreva (this paper); 4, Kossovaya (this paper). Borehole 4K, 37°30'E - 55°N; Borehole 5K, 38°E - 55°45'N.



LOG. 34. — Moscow Basin, coordinated by Briand *et al.* Kasimovian and Gzhelian of the Moscow Basin. 1, Briand *et al.* (1998); 2, Makhlina *et al.* (1997); 3, Goreva (this paper); 4, Kossovaya (this paper). Boreholes 1/42, 35°50'E - 56°42'N; Borehole 7K, 38°30'E - 56°N.


SEQUENCES		URAL (PERM) BASIN			
Gubakha	2nd order	Formations	Fusulinids	Eastern European Stages	
		Urminsk			ARTINSKIAN
			<i>Eoparafusulina lutugini</i>	Irginian	
			<i>Schwagerina concavatus</i>	Burtsevkian	
		Limestone Formation	<i>Schwagerina uralensis</i>	Sterlitamakian	SAKMARIAN
			<i>Schwagerina verneuilli</i> <i>Schwagerina uralica</i>	Tastubskian	
			<i>Schwagerina moelleri</i>		
			<i>Sphaeroschwagerina sphaerica</i>	Shikhanian	
			<i>Sphaeroschwagerina moelleri</i>	Kholodnologian	
	<i>Sphaeroschwagerina fusiformis</i>				
		Limestone Formation	<i>Ultradaxina bosbytauensis</i>		ORENBURGIAN
			<i>Daixina sokensis</i>		GZHELIAN
			<i>Jigulites jigulensis</i>		
			<i>Triticites rossicus</i>		KASIMOVIAN
			<i>Triticites acutus</i>		
		Limestone Formation	<i>Montiparus montiparus</i>		MOSCOVIAN
			<i>Protriticites pseudomontiparus</i>		
			<i>Fusulina cylindrica</i>	Myachkovian	
			<i>Fusulinella colaniae</i>	Podolskian	
			<i>Fusulinella subpulchra</i> <i>Aljutovella priscoidea</i>	Kashirian	
			<i>Aljutovella aljutovica</i>	Vereian	

LOG. 35. — Moscovian to Artinskian in Central Ural Basin (Perm), coordinated by Izart *et al.*, after Chuvachov (1993). Gubakha, Russia, 57°30'E - 58°50'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE EASTERN PRECASPIAN BASIN

Janajol	Formations	Fusulinids	Conodonts	Eastern European Stages			
				KUNGURIAN			
				ARTINSKIAN			
				<i>Schwagerina moelleri</i>	<i>Gondolella biselli</i>	SAKMARIAN	
				Janatan	<i>Sphaeroschwagerina moelleri, fusiformis</i>	<i>Streptognatodus wabaunsensis</i>	ASSELIAN
	Janaturmyce	<i>Daixina sokensis</i>	<i>Streptognatodus elongatus</i> <i>Idiognathodus sp.</i>	ORENBURGIAN GZHELIAN KASIMOVIAN			
		<i>Montiparus montiparus</i>					
	Kulakchy			Myachkovian	M O S C O V I A N		
	Jaiyndy			Podolskian			
	Alibek						
				Sinelnikov		Kashirian	
	Vereian					BASHKIRIAN	
200m							
0							

LOG. 36. — Late Carboniferous and Early Permian of the Eastern Precaspian Basin (Kazakhstan), coordinated by Ensepbaev, after Ensepbaev (this paper). Janajol, Kazakhstan, 57°E - 50°N.



Log. 37. — Late Carboniferous of northwestern Spain Basins, coordinated by Izart. 1, Bouroz *et al.* (1972), Wagner & Winkler-Prins (1985); 2, Lys (1988a); 3, Coquel (this paper); 4, this volume. La Pernia, Spain, 4°30'W - 43°N; Guardo, Spain, 4°50'W - 42°53'N; Tejerina, Spain, 5°01'W - 42°55'N; Castilleria, Spain, 4°28'W - 42°56'N; Barruelo, Spain, 4°16'W - 42°53'N; Sabero, Spain, 5°10'W - 42°50'N; Villablino, Spain, 6°10'W - 42°58'N.

MOSCOWIAN STRATIGRAPHIC UNITS IN FOUR STRUCTURAL UNITS OF THE CANTABRIAN ZONE (SPAIN)



La Sobia Unit	Central Asturian Coalfield	Ponga Unit	Picos de Europa Unit	BIOSTRATIGRAPHY		CHRONOSTRATIGRAPHY	
LITHOSTRATIGRAPHY	LITHOSTRATIGRAPHY	LITHOSTRATIGRAPHY	LITHOSTRATIGRAPHY	Fusulinid Zones (1)	Characteristic assemblages (2)	Russian horizons (3)	Stages

LOG. 38. — Cantabrian Basin (Spain), coordinated by Villa. Moscovian stratigraphic subdivisions in four structural units of the Cantabrian zone (Spain). 1, after van Ginkel (1965); 2, 3, Villa (this paper). Mieres, Spain, 5°47'W - 43°16'N.

KASIMOVIAN OF THE PICOS DE EUROPA (CANTABRIAN ZONE, SPAIN)


Las Llacierías Section		Gamonedo -Cabrales Area		BIOSTRATIGRAPHY	CHRONOSTRATIGRAPHY	
LITHOSTRATIGRAPHY		LITHOSTRATIGRAPHY		Fusulinid Zones (1)	Russian horizons (2)	Stages
		Cavandi Fm.			YAUZSKY	K
		Puentellés Fm.		<i>Rauserites</i>	DOROGOMILOVSKY	A
	?					S
						I
				<i>Montiparus</i>	KHAMOVNICHESKY	M
	Covadonga Beds (ex "Puentellés Fm.")	Dobros Lmst.				O
		Gamonedo Beds				V
						I
				<i>Protriticites</i>	KREVIKINSKY	A
	Picos de Europa Fm.					N
					MYACHKOVSKY (uppermost part)	MOSCOVIAN
		Gamonedo Lmst.				

LOG. 39. — Cantabrian Basin (Spain), coordinated by Villa. Kasimovian of the Picos de Europa (Cantabrian zone, Spain). 1, according to Villa (1985, modified); 2, Villa (this paper). Picos de Europa, Spain, 4°50'W - 43°15'N.




LOG. 41. — Italy Basins, coordinated by Pasini & Vai. Early Permian of the southern Tuscan shelf, Pasini & Vai (1997).

LOG. 40. — Italy Basins, coordinated by Pasini & Vai. Moscovian of the southern Tuscan Shelf, Pasini & Vai (1997).




LOG. 42. — Moscovian of the southern Tuscan Basin, Pasini & Vai (1997), Borehole PC33b (Tuscany, Italy).

MOSCOVIAN OF THE CARNIC ALPS (Pramollo Basin)




Log. 43. — Austria boundary, coordinated by Vai, Pasini & Venturi. Moscovian of the Carnic Alps (Pramollo Basin), Pasini (this paper).


KASIMOVIAN AND EARLY GZHELIAN OF THE CARNIC ALPS (Pramollo Basin)



LOG. 44. — Austria boundary, coordinated by Vai, Pasini & Venturi. Kasimovian and early Gzhelian of the Carnic Alps (Pramollo Basin). Vai & Venturi (1997).



LOG. 45. — Austria boundary, coordinated by Vai, Pasini & Venturi. Late Gzhelian of the Carnic Alps (Pramollo Basin), Vai & Venturi (1997).




LOG. 46. — Austria boundary, coordinated by Vai, Pasini & Venturi. Early Permian of the Carnic Alps (Pramollo Basin), Vai & Venturi (1997).

LATE CARBONIFEROUS OF THE CARNIC ALPS


SEQUENCES						
Carnic Alps	4th order	3rd order	Formations	Fusulinids	Macroflora	West & East European Stages
			Lower Pseudo-schwagerina Limestone (Schulterkofel)	<i>Ul. dahtidzhumica</i> <i>Ul. bosbytauensis</i> <i>Occidentoschwagerina alpina</i> <i>Sch. ulukensis</i> <i>Ruzhencevites ferganensis</i>		ORENBURGIAN
			Carnizza	<i>Pseudofusulinoides regularis</i> <i>Triticites immutabilis</i> <i>T. paraduplex</i> <i>T. perlongus</i> <i>T. turkestanensis</i>	<i>Sphenophyllum oblongifolium</i> <i>Aphlebia elongata</i> <i>Odont. alpina</i> <i>Odont. brardii</i> <i>Pec. feminaeformis</i>	
			Auernig	<i>Schagonella gigantea</i> <i>Pseudofusulina devexa acallosa</i> <i>P. multiseptata</i> <i>P. paraconcinna</i> <i>Dutkevitchia dastarensis</i> <i>D. kargalensis</i> <i>D. ruzhencevi</i>	<i>Aphlebia elongata</i> <i>Alethopt. bohémica</i> <i>Odontopteris brardii</i> <i>Pec. feminaeformis</i> <i>Callipt. pteridium</i> <i>Pec. ex gr. arborescens-schlotheimii</i>	
			Corona	<i>Daixina alpina</i> <i>D. ex gr. admirabilis</i> <i>Schagonella spp.</i> <i>Pseudofusulina multiseptata</i>	<i>Alethopteris bohémica</i> <i>Odontopteris alpina</i> <i>Odontopteris brardii</i> <i>Pec. feminaeformis</i> <i>Pseudomariopteris busquetii</i> <i>Sigillaria brardii</i> <i>Sphenophyllum oblongifolium</i>	STEPHANIAN C
			Pizzul	<i>Daixina alpina</i> <i>D. sokensis</i> <i>D. naviculaeformis</i> <i>D. sakmarensis</i> <i>Triticites oryziformis</i> <i>Jigulites jigulensis</i> <i>Schagonella spp.</i> <i>Daixina sp.</i>	<i>Pseudomariopteris busquetii</i> <i>Sphenophyllum oblongifolium</i>	
			Meledis	<i>Rauserites sp.</i> <i>Ferganites aff. ferganensis</i> <i>Montiparus montiparus</i> <i>Protriticites pseudomontiparus</i>	<i>Sphenophyllum angustifolium</i> <i>Sphenophyllum oblongifolium</i>	KASIMOVIAN
			Bombaso	<i>Protriticites ovatus</i> <i>Quasifusulinoides quasifusulinoides</i>	<i>Linopteris neuropteroides</i> <i>Neuropteris ovata</i> <i>N. scheuchzeri</i> <i>Sphenophyllum oblongifolium</i>	CANTABRIAN
	T	R T R				MOSCOVIAN

LOG. 47. — Austria boundary, coordinated by Krainer & Davydov. Late Carboniferous of the Carnic Alps, Krainer (this paper).




LOG. 48. — Austria boundary, coordinated by Vai, Pasini, Venturi, Krainer & Davydov. Early Permian of the Carnic Alps, Krainer (this paper). Auernig, Carnic Alps, 13°17'E - 46°33'N; Trogkofel, Carnic Alps, 13°13'E - 46°35'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF IRAN BASIN




LOG. 49. — Late Carboniferous and Early Permian of eastern Elbourz Basin (Iran), coordinated by Jenny & Jenny-Deshusses, after Jenny & Jenny-Deshusses (this paper). Gheselghaleh, Iran, 53°E - 34°N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE CENTRAL MOROCCO BASIN




LOG. 50. — Morocco Basins: Late Carboniferous and Early Permian of the Central Morocco Basin, coordinated by El Wartiti and Broutin, after El Wartiti & Broutin (this paper). Ait Ziffa, 7°30'W, 31°27'N; Bou Achouch, 5°44'W, 33°42'N; Khenifra, 5°40'W, 33°N; Tiddas, 6°16'W, 33°34'N; Haouz, 7°10'W, 32°N.

WESTPHALIAN C OF THE MOROCCAN BASIN




Log. 51. — Morocco Basins: Westphalian of the eastern Morocco Basin, coordinated by Izart, after Izart (this paper). Jerada, 2°W - 34°30'N.




Log. 52. — Westphalian of the southern Algeria Basins, coordinated by Izart, after Nedjari (1982). Kenadza, southern Algeria, 2°W - 31°45'N; Mezarif, Southern Algeria, 1°45'W - 31°45'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE SOUTH TUNISIA BASIN



Log. 53. — Late Carboniferous and Early Permian of the southern Tunisia Basin, coordinated by Massa and Lys. 1, Massa (this paper); 2, Lys (this paper). Kirchaou, 10°40'E - 33°N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE GHADAMES BASIN



Log. 54. — Libya Basins, coordinated by Massa, Vachard and Coquel. Late Carboniferous and Early Permian of the Ghadames Basin (Libya). 1, Massa (this paper); 2, Vachard (this paper); 3, Coquel (this paper). Ghadames, Libya, 10°30'E - 29°N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF THE CYRENAIC BASIN

	Formations (1)	Fusulinids (2)	Palynomorphs	Eastern European Stages
A1-19				
		not described		EARLY PERMIAN
A1NC92		A1-37		
		<i>Nodosaria cf. shikhanica</i>	<i>Strotersporites indicus</i> <i>Protohaploxypinus goraiensis</i>	GZHELIAN
A1-37	125 m 0	<i>Fusulinella ex gr. colaniae</i>	<i>Plicatipollenites malabarensis</i>	late MOSCOVIAN
		<i>Syzrania bella</i> <i>Schubertella obscura (mosquensis)</i> <i>Eostaffella acuta</i>	<i>Cannanoropollis janakii</i>	early MOSCOVIAN
				BASHKIRIAN

Log. 55. — Libya Basins, coordinated by Massa, Vachard and Coquel. Late Carboniferous and Early Permian of the Cyrenaic Basin (Libya). 1, Massa (this paper); 2, Vachard *et al.* (1993). A1-37 borehole, 22°53'E - 30°44'N; A1NC92 borehole, 22°08'E - 31°N; A1-19 borehole, 23°40'E - 31°N.


LATE CARBONIFEROUS AND EARLY PERMIAN OF THE NORTH EAST EGYPT BASIN

SEQUENCES

NE Egypt	Formations	Foraminifera	Conodonts	Corals	Brachiopods	Macroflora	Eastern European Stages
							KUNGURIAN
							ARTINSKIAN
	Aheimer				<i>Rhipidomella cordialis</i>	<i>Cordaites sp.</i>	SAKMARIAN
		<i>Hemigordius haritoni</i>	<i>Streptognathodus spp.</i>	<i>Bothrophyllum pseudoconicum</i>	<i>Anthracospirifer sp.</i>	<i>Callipteris conferta</i>	ASSELIAN
							GZHELIAN
	Abu Durba		<i>Idiognathodus Neognathodus</i>		<i>Neospirifer cameratus</i>		KASIMOVIAN
		<i>Glomospira Nodosinella</i>				<i>Calamites sp.</i>	MOSCOVIAN
40 m 0							
T	R						


Log. 56. — Late Carboniferous and Early Permian of the northeastern Egypt Basin, coordinated by Kora, after Kora (1995). Abu Durba, Egypt, 33°18'E - 28°35'N; Wadi Aheimer, Egypt, 32°23'E - 29°30'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF SOUTHERN TURKEY

Bademli	Formations (1)	Foraminifera (2)	Eastern European Stages
	Cevizli Limestone	<i>Sphaeroschwagerina sphaerica</i> <i>Quasifusulina cayeuxi</i> <i>Tubiphytes obscurus</i>	ASSELIAN
		<i>Daixina sokensis</i> <i>Triticites sp.</i>	ORENBURGIAN GZHELIAN
20 m 0	Quartzites	<i>Fusulina paradistenta</i>	KASIMOVIAN
		<i>Fusulina cf. cylindrica</i>	MOSCOVIAN
		<i>Aljutovella sp.</i> <i>Staffellaeformes staffellaeformis</i>	

LOG. 57. — Late Carboniferous and Early Permian of the southern Turkey, coordinated by Monod. 1, Monod (1977); 2, Lys (1988). Bademli, southern Turkey, 32°E - 37°30'N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF SYRIA

Syria	Formations	Fauna - Flora	Eastern European Stages
	Heil		EARLY PERMIAN
	Najeeb	<i>Potonieisporites sp.</i> <i>S. triangulus</i>	GZHELIAN KASIMOVIAN
125 m 0	Sawanet		MOSCOVIAN
			BASHKIRIAN

LOG. 58. — Late Carboniferous and Early Permian of Syria, coordinated by Izart, after Al Youssef & Ayed (1992). Sawanet 2, Syria, 38°E - 34°10'N.

PERMIAN OF ISRAEL BASIN

Bakhtesh Qatan 2	Group	Formation	Palynomorphs	Western European Stage
	Negev	Sa'ad	<i>Potoniesporites novicus</i> <i>Distrialites insolitus</i> <i>Vittatina ovalis</i> <i>Plicatipollenites indicus</i> <i>Punctatoporites minutus</i> <i>Lophotriletes novicus</i> <i>Vittatina costabilis</i> <i>Protohaploxypinus microcorpus</i> <i>Laevigatosporites callosus</i>	AUTUNIAN

Log. 59. — Permian of Israel Basin, coordinated by Weissbrod, after Weissbrod (this paper). Well Bakhtesh Qatan 2, 35°E - 31°N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF SAUDI ARABIA

Well St 8	Formations	Palynomorphs	West & East European Stages	
	Unayzah		KUNGURIAN	E A R L Y P E R M I A N
			ARTINSKIAN	
			SAKMARIAN	
50 m 0	Berwath	<i>Thymospora obscura</i> <i>Torispota granulata</i> <i>Punctatosporites</i> spp. <i>Potoniesporites</i> spp.	WESTPHALIAN D	



Log. 60. — Late Carboniferous and Early Permian of Saudi Arabia, coordinated by Izart and Vaslet, after Owens & Turner (1995) and Al Laboun (1993). Well 8, Saudi Arabia, 42°E - 30°N.

LATE CARBONIFEROUS AND EARLY PERMIAN OF OMAN

Suwaihat 3	Formations	Palynomorphs	Brachiopods	West & East European Stages
	Saiwan = "early Gharif" (1)	<i>Kingiacolpites subcircularis</i>	<i>Cyrtella nagmargensis</i>	ARTINSKIAN LATE SAKMARIAN Sterlitamakskian
	Al Khlata	<i>Cycadopites cymbatus</i> <i>Protohaploxypinus</i> spp. "Vittatina"		SAKMARIAN ASSELIAN
		<i>Microbaculispora</i> spp. <i>Cristatisporites</i> spp. <i>Parasaccites</i> spp.		STEPHANIAN
		<i>Potonieisporites</i> spp. <i>Punctatisporites</i> spp.		EARLY STEPHANIAN LATE WESTPHALIAN

Log. 61. — Late Carboniferous and Early Permian of Oman, coordinated by Izart and Vaslet, after Broutin *et al.* (1995) and Angiolini *et al.* (1997); 1, Saiwan = ex "early Gharif" *sensu* Love (1994). Well Suwaihat 3, 55°E - 20°N.

EARLY PERMIAN OF SALT RANGE (PAKISTAN)

Salt Range	Formations	Fusulinids	Pelecypods	Palynomorphs	Eastern european Stages
	Amb	<i>Cancellina</i>		<i>Verrucosisporites cf planiverrucatus</i>	KUBERGANDIAN
		<i>Misellina</i>			BOLORIAN
	Sardhai				ARTINSKIAN ?
	Warcha				SAKMARIAN ?
	Dandot		<i>Eurydesma</i>	<i>Cycadopites cymbatus</i>	SAKMARIAN Tastubskian
	Tobra			<i>Protohaploxypinus limpidus</i>	ASSELIAN

CAMBRIAN

Log. 62. — Early Permian of Salt Range Basin (Pakistan), coordinated by Iqbal *et al.*, after Iqbal (this paper). Salt Range, 72°E - 32°N.

APPENDIX 2

- (1) Laboratoire de Géologie des Ensembles Sédimentaires, Université de Nancy I, B.P. 239, F-54506 Vandœuvre-les-Nancy (France)
- (2) BRGM-SGN, B.P. 6009, F-45060 Orléans cedex 2 (France)
- (3) Laboratoire de Paléobotanique, Université Pierre-et-Marie-Curie, 12 rue Cuvier, F-75252 Paris cedex 05 (France)
- (4) Laboratoire de Paléobotanique, Université des Sciences et Technologies Lille I, F-59655 Villeneuve d'Ascq (France)
- (5) Department of Geosciences, Boise State University, 1910 University Drive, Boise, Idaho 83725 (USA)
- (6) Faculté des Sciences, Département des Sciences de la Terre, Université Mohammed V, Avenue Ibn Batouta, B.P. 1014, Rabat (Morocco)
- (7) Faculté du Pétrole, département de Géologie du Pétrole et du Gaz, Institut polytechnique Kazakh, 22 rue Satpaev, 480013 Almaty (Kazakhstan)
- (8) Geological Survey, PO Box 157, 2000 AD Haarlem (The Netherlands)
- (9) Geological Institute of the Academy of Sciences of Russia, 7 per. Pyzhevsky, Moscow 109117 (Russia)
- (10) Madem Fakultesi, Jeoloji Bölümü, Tubitak-Gloek, ITU, Ayazaga, 80626, Istambul (Turkey)
- (11) Pakistan Museum of Natural History, 7/2 Islamabad (Pakistan)
- (12) All-russian Research Geological Institute (VSEGEI), Sredny pr. 74, Saint Petersburg (Russia)
- (13) Institut für Geologie und Paläontologie der Universität Innsbruck, Innrain 52, A-6020 Innsbruck (Austria)
- (14) Industrial Geological Amalgamation, Tsenter Geology, 39a, Warshawshoye Shosse, Moscow (Russia)
- (15) Institute of Geological Sciences, Ukraineian Academy of Sciences, 55b Chkalov Street, Kiev, 252601 (Ukraine)
- (16) Faculty of Sciences, Department of Geology, University of Mansoura, Mansoura 35516 (Egypt)
- (17) 6 rue J.-J.-Rousseau, F-92150 Suresnes (France)
- (18) École des mines de Paris, CGES-sédimentologie, 35 rue Saint-Honoré, F-77305 Fontainebleau cedex (France)
- (19) Département des Sciences de la Terre, Université d'Orléans, B.P. 6759, F-45067 Orléans 2 (France)
- (20) Department of Geology, Charles University, Albertov 6, 128 43 Praha 2 (Czech Republic)
- (21) Institute of Geology, Department of Paleontology, Freiberg University of Mining and Technology, Bernhard von Cotta Str. 2, D-09596 Freiberg (Germany)
- (22) Geologische Bundesanstalt, postfach 154, A-1031 Wien (Austria)
- (23) Geologisches Institut, Nussalle 8, D-53757 Bonn (Germany)
- (24) Dipartimento di Scienze geologiche, Università degli studi di Bologna, 67 via Zamboni, I-40127 Bologna (Italy)
- (25) Geological Survey, Mlynska dolma 1, 81704 Bratislava (Slovak Republic)
- (26) Geological Survey, 30 Malkhe Israel Street, Jerusalem 95501 (Israel)
- (27) Polish Geological Institute, Late Silesian branch, Sosnowiec (Poland).