

A new Pennsylvanian *Oriocrassatellinae* from Brazil and the distribution of the genus *Oriocrassatella* in space and time

Luiz E. ANELLI

Instituto de Geociências, Universidade de São Paulo,
Rua do Lago 562, CEP 05508-080 São Paulo, SP (Brazil)
anelli@usp.br

Marcello G. SIMÕES

Instituto de Biociências, Universidade Estadual Paulista,
CP 510, CEP 18618-000 Botucatu, SP (Brazil)
profmsgsimoes@gmail.com

Carlos Roberto GONZÁLEZ

CONICET, Fundación Miguel Lillo,
Miguel Lillo 251 (4000), San Miguel de Tucumán (Argentina)
crgonzalez@csnat.unt.edu.ar

Paulo A. SOUZA

Instituto de Geociências, Universidade Federal do Rio Grande do Sul,
Av. Bento Gonçalves, 9500, CEP 91540-000, Porto Alegre, RS (Brazil)
paulo.alves.souza@ufrgs.br

Anelli L. E., Simões M. G., González C. R. & Souza P. A. 2012. — A new Pennsylvanian *Oriocrassatellinae* from Brazil and the distribution of the genus *Oriocrassatella* in space and time. *Geodiversitas* 34 (3): 489-504. <http://dx.doi.org/10.5252/g2012n3a2>

ABSTRACT

Oriocrassatella Etheridge Jr., 1907 is a long range crassatellid bivalve genus well recognized in shallow waters of epeiric seas throughout the upper part of Paleozoic. The first occurrences of this genus are recorded in the sedimentary successions of the Gondwana, both in Australia and South America. However, the geographic and age distribution of *Oriocrassatella* in Late Mississippian deposits of Australia and Argentina may indicate an earliest Visean or even a pre-Visean origin for the genus. Following its origin in Early Carboniferous a complex paleobiogeographic history from Southern to Northern Hemisphere took place in the Permian. During its initial dispersal phase from Late Carboniferous to the Early Permian the genus thrived in cold water environments associated to the Late Paleozoic Gondwana glaciation. Shallow-water bottoms of the warm

KEY WORDS

Oriocrassatella,
bivalve,
Gondwana,
Upper Paleozoic,
glaciation,
paleobiogeography,
new species.

waters of the central Gondwana fringe and Laurussia were colonized by *Oriocrassatella* only during Early Permian times when the genus became cosmopolitan. A new species of this genus is described herein, *Oriocrassatella piauiensis* n. sp., recorded from the Piauí Formation, Pennsylvanian of the Parnaíba Basin. This new species may represent an early adaptation to warm waters. However, based on available data, species of this genus seem to have adapted definitely to warm water environments probably related the Late Pennsylvanian interglacial phases. In these phases, climatic barrier were interrupted allowing the faunal interchange and larval dispersion following a South to North migration route through the eastern margins of Gondwana and the eastern Paleotethys.

RÉSUMÉ

Nouvel Oriocrassatellinae du Pennsylvanien du Brésil et distribution spatio-temporelle d'Oriocrassatella.

Oriocrassatella Etheridge Jr., 1907 est un crassatellidé à large distribution stratigraphique, bien connu dans les mers épicontinentales en eaux peu profondes, caractéristique du Paléozoïque supérieur. Ce genre apparaît pour la première fois dans des dépôts sédimentaires du Gondwana, en Australie et en Amérique du Sud. D'après sa distribution spatio-temporelle dans des dépôts du Mississippien supérieur, *Oriocrassatella* serait originaire du début du Viséen, ou même pré-viséenne. Par la suite, depuis le Carbonifère inférieur jusqu'au Permien, ce genre a eu une histoire paléobiogéographique complexe allant de l'hémisphère Sud vers l'hémisphère Nord. Pendant la phase initiale de dispersion à partir du Carbonifère supérieur jusqu'au Permien inférieur, ce genre a prospéré dans un milieu dominé par des eaux froides du Gondwana, suite à la glaciation du Paléozoïque supérieur. Les eaux littorales chaudes du Gondwana central et de Laurussia ont été colonisées par *Oriocrassatella* uniquement durant le Permien inférieur, parce que ce genre est devenu cosmopolite. Une espèce nouvelle, *O. piauiensis* n. sp., est décrite de la Formation Piauí du Pennsylvanien du bassin du Parnaíba; elle pourrait représenter une première adaptation aux eaux chaudes. En outre, et en s'appuyant sur les données disponibles, d'autres espèces d'*Oriocrassatella* se seraient aussi définitivement adaptées à des milieux en eaux chaudes au cours des périodes interglaciaires du Pennsylvanien tardif, car des barrières climatiques ont disparu, favorisant des échanges de faune avec dispersion des larves, et migration du sud vers le nord le long des bordures est du Gondwana et de la Paléotéthys.

MOTS CLÉS

Oriocrassatella,
bivalve,
Gondwana,
Paléozoïque supérieur,
glaciation,
paléobiogéographie,
espèce nouvelle.

INTRODUCTION

Due to its paleoecological significance and paleogeographical distribution, the importance of *Oriocrassatella* Etheridge Jr., 1907 was highlighted by various authors (Boyd & Newell 1968; Dickins 1996; Archbold *et al.* 1996; Larghi 2005). Bivalve molluscs belonging to the cosmopolitan, long range

genus *Oriocrassatella* were first described by Etheridge Jr. (1907) from the Lower Permian of Western Australia. Later, Yakovlev (1928) erected the genus *Procrassatella*, a very similar crassatellacean from Russia, subsequently regarded by Newell (1958) as a junior subjective synonym of *Oriocrassatella*. At the present time, *Oriocrassatella* is a crassatellacean well-characterized by a hinge area with strong septate

nymphs, a large, well-defined triangular resilium and two cardinal teeth in each valve.

Until 1968, 19 occurrences of the genus were known, with only two species recognized in western Pangea (Boyd & Newell 1968: 29, table 1). In that contribution, the subfamily *Oriocrassatellinae* was erected to accommodate members of the family *Crassatellidae* Férussac, 1822, characterized, among other features, by the overlap of the left valve on right valve in front of beaks. In addition, these authors briefly discussed a possible evolutionary pathway for the genus referring to the presence of members of the family *Crassatellidae* in pre-Carboniferous times. However, the origin of the *Oriocrassatella* lineage is still uncertain due to its questionable phylogenetic affinity with the Devonian genera *Cypriocardella* Hall, 1858 and/or *Crassatellopsis* Beushausen, 1895.

Boyd & Newell (1968) were also the first authors to analyze the distribution of the genus in space and time. They suggested a possible Gondwanan origin in the Lower Carboniferous, and the dispersion to northern portions of Pangea during Permian times. Based on an extensive stratigraphical revision including new chronostratigraphic data, as well as about 18 newly recognized species, Archbold *et al.* (1996) reassessed the distribution of this genus asserting its Gondwanan origin in Lower Carboniferous and the posterior migration throughout Pangea. However, afterwards three new carboniferous species of *Oriocrassatella* were identified in South America (González 1997, 2002; Sterren 2005; see Figs 1, 2). A new species of *Oriocrassatella* is described herein from the Pennsylvanian of the Parnaíba Basin, Brazil, the tenth occurrence of the genus described in South America. A thorough revision of the records of this genus is presented based on their updated ages, including Argentinean and Australian data, offering new interpretations on the distribution of this *Crassatellidae* taxon in space and time, with significant paleobiogeographic implications.

ORIOCRASSATELLA IN SPACE AND TIME

According to the available data, around 42 occurrences of *Oriocrassatella* are known at the present




FIG. 1. — Paleogeographic distribution of *Oriocrassatella* Etheridge Jr., 1907 in the: **A**, Carboniferous; **B**, Lower Permian; **C**, Upper/Middle Permian. Paleogeography according to Scotese's Climate Story at www.scotese.com. Numbers according to references in Figure 2.

time. The genus was recorded in nearly 24 localities throughout the Pangea in Carboniferous and Permian strata (see Figs 1; 2).

Figure 2 displays the distribution of *Oriocrassatella* in space and time together with the Permo-Carboniferous glacial events. A simplified correlation of the Late Palaeozoic brachiopod zones from Argentina and eastern Australia is also shown. In this context we discuss the issues related to time and region of origin of the genus, and possible pathways that lead *Oriocrassatella* to reach and widespread in warm water environments throughout the Late Paleozoic.

THE CARBONIFEROUS FOSSIL RECORD OF *ORIOCRASSATELLA* IN THE GONDWANA

Nine occurrences are restricted to the Carboniferous, eight from Gondwanaland and one from Early Pennsylvanian Carboniferous in northern Laurussia (Muromtzeva 1974 in Biakov 2005). This record is puzzling, since all other Carboniferous occurrences are invariably recorded in Gondwana terrains (Figs 1; 2).


Australia

Two occurrences were described from Carboniferous strata in Australia. Records of *Oriocrassatella compressa* Maxwell, 1964, from Rands (Carboniferous) and Burnet formations (Lower Permian) were long considered to be the oldest for the genus in Gondwana (see Boyd & Newell 1968; Archbold et al. 1996). The Burnett Formation is probably Permian – it has *Eurydesma burnettensis* Waterhouse,

1986 that looks moderately close to *E. cordatum* Morris, 1845, and both are considered to be Early Asselian in age (Waterhouse pers. comm.). Overlying beds with *Bandoproductus macrospina* Waterhouse, 1986 are either latest Asselian (as generally favored for South Asian occurrences) or Early Tassubian (Sakmarian) (Waterhouse pers. comm.). In the Rands Formation *O. compressa* occurs below *Auriculispina levis*, and above *Levipustula levis* zones. *Auriculispina levis* appears to be the youngest known Carboniferous occurrence in eastern Australia, but the precise age of this record is open to conjecture. It could be Gzhelian or probably post-Moscovian, and certainly not as old as Namurian (Waterhouse pers. comm.). Hence, these data confine *O. compressa* to Upper Early to Middle Pennsylvanian beds. However, if *O. compressa* occurs within the post-Moscovian Rands Formation as well as in the upper part of the *Marginirugus barringtonensis* zone in the Faulkland Formation (Namurian after Peou & Engel 1979), the range of the species may be limited from Bashkirian to Moscovian, respectively Early to Middle Pennsylvanian.

Based on data above, the small *Oriocrassatella* sp. of Peou & Engel (1979) probably is the oldest known occurrence of this genus. This is because *Oriocrassatella* sp. occurs within the upper part of the *Marginirugus barringtonensis* zone of Faulkland Formation in the Rawdon Valley region (Australian Gondwana; Peou & Engel 1979; see Figs 1A, 2), being considered by Roberts et al. (1976) Upper Viséan to Lower Early Serpukhovian in age.

Fig. 2. — Distribution of *Oriocrassatella* Etheridge Jr., 1907 in space and time: 1–37, occurrences and ages according to: 1, Peou & Engel (1979); 2, González (2002); 3, *Oriocrassatella piauiensis* n. sp. described here, Campanha & Rocha-Campos (1979); 4, Sterren (2005), Azcuy et al. (2007); 5, González (1982); 6, González (1976), Sabattini et al. (1990), Azcuy et al. (2007); 7, Manceñido et al. (1976), Azcuy et al. (2007); 8, González (1997); 9, Maxwell (1964) (Bruce Waterhouse pers. comm. 2009); 10, Muromtzeva (1974); Biakov (2005, and written communication 2009); 11, Termier et al. (1974), Archbold et al. (1996); 12, Dickins (1963), Vanderhor et al. (2006); 13a–c, Reed (1932), Veevers (1995), Honegger et al. (1982); 14a, Dickins & Shah (1979), Stephenson et al. (2007); 14b, Dickins & Shah (1979), Angiolini et al. (2006); 14c, Larghi (2005); 15, Rocha-Campos (1970), Holz et al. (2008, 2010); 16, Dickins (1961, 1996); 17, Dickins (1963), Archbold et al. (1996); 18, Dickins & Skwarko (1981), Archbold et al. (1996), Charlton (2001); 19, Biakov (2005); 20a, b, Liu & Cui (1983), Archbold et al. (1996), Jin (2002); 21, Muromtseva & Guskov (1984), Archbold et al. (1996); 22, Kanev (1994), Shi et al. (2002); 23, Yakovlev (1902), Archbold et al. (1996); 24, Newell (1958); 25, Lobanova (1961), Archbold et al. (1996); 26, Boyd & Newell (1977), Archbold et al. (1996); 27, Newell (1955), Archbold et al. (1996); 28, Boyd & Newell (1968), Archbold et al. (1996); 29, Ciriacks (1963), Boyd & Newell (1968); 30, Suveizdas et al. (1975), Archbold et al. (1996); 31, Clarke (1987); 32a, b, Sanchez (1984 and pers. comm. 2009); 33, Newell (1958); 34, Dickins et al. (1989), Archbold et al. (1996); 35, Archbold et al. (1996); 36, Waner (1940), Archbold et al. (1996). Ages for glacial periods: in Argentina, according to González (1981, 1990, 2002) and González & Diaz-Saravia (2007); in Australia, to Roberts et al. (1995), Dickins (1996) and Fielding et al. (2008a, b); in Brazil, to Holz et al. (2008); in India, to Veevers & Tewari (1995) and Maejima et al. (2001). Ages for Biozones in Argentina and Australia from Kelly et al. (2001); for *Marginirugus barringtonensis*, Roberts et al. (1976).



South America

Coeval marine deposits in South America are found in western Argentina referred to the *Rugosochonetes-Bulahdelia* zone (Taboada 1997; see also Fig. 2), but *Oriocrassatella* was not found in these rocks, perhaps due to incomplete sampling or poor preservation. In western Argentina (San Eduardo Formation) the *Rugosochonetes-Bulahdelia* fauna occurs in beds immediately above the glacial phase El Paso II, and underlay the *Levipustula levis* zone (González 1990). In fact, Sterren (2005) identified as *Oriocrassatella*? sp. a specimen from the upper portion of the *Rugosochonetes-Bulahdelia* fauna (see Figs 1, 2), but it shows an incomplete hinge and an unusual profile for the genus.

The earliest known record of the genus in South America could be *Oriocrassatella andina* González, 2002, described from Serpukhovian-Bashkirian deposits of Western Argentina (see Figs 1A, 2; González 2002). *Oriocrassatella andina* is the only Oriocrassatellinae species known from the *Levipustula levis* zone (Serpukhovian-Bashkirian).

Oriocrassatella sp. (González 1997) from the Río del Peñón Formation belongs to the *Kochiproductus-Heteralosia* assemblage, the youngest faunal assemblage of the informal Aguanegran faunal stage. The same is also true for *Oriocrassatella* sp. (Manceñido *et al.* 1976). Both species are restricted to the Gzhelian (see Fig. 2[7; 8]).

Oriocrassatella sanjuanina? (González 1982) from the Agua del Jaguel Formation (southern Uspallata-Iglesia Basin) is probably a new species. It is much younger than *O. sanjuanina* (González 1976) from the Cerro Agua Negra Formation and belongs to the *Costatumulus amosi* fauna, restricted to the Upper Asselian-Lower Sakmarian, subsequent to the Early Permian glacial sequence, that should be restricted to early Asselian in western Argentina.

In Brazil, *Oriocrassatella piauiensis* n. sp., which is described below, was found in the Piauí Formation (Upper Carboniferous), Parnaíba Basin, Northeastern Brazil. The Piauí Formation crops out in a large area at the northeastern part of the Parnaíba Basin (Fig. 3A), Piauí State, Brazil. Several fossiliferous dolostones beds occur near the José de Freitas village, and are associated with sandstones, siltstones, and mudstones deposited in a shallow-

marine carbonatic shelf (Lima Filho 1991, 1999). One of the dolostones, informally named Mucambo (Fig. 3A-C), together with other three beds located in the region (see Anelli *et al.* 2006), contains the most diverse bivalve fossil record known from the Upper Paleozoic rocks of South America (Anelli 1999; Anelli *et al.* 2002, 2006, 2009). The Mucambo dolostone bed is Lower to Middle Pennsylvanian in age (Bashkirian to Lower Moscovian; Campanha & Rocha-Campos 1979). The fauna also includes commonly many gastropods, cephalopods, brachiopods and bryozoans. Trilobites are less common (Anelli 1999). The associated fauna shows clear affinities to other Upper Pennsylvanian South and North American warm water faunas. A Bashkirian to Lower Moscovian age attributed to the Piauí Formation is also supported by palynomorphs, from subsurface material (Dino & Playford 2002; Souza *et al.* 2010). An age younger than Middle Moscovian to this unit is discarded, taking into account the glacial event in Paraná Basin (Itararé Subgroup), that occurred after these times. Then, a significant gap separates the Piauí Formation and the overlying Pedra de Fogo Formation, comprising the Late Moscovian to Carboniferous/Permian boundary (Melo J.H.G. pers. comm.).

THE PERMIAN FOSSIL RECORD OF *ORIOCRASSATELLA* IN GONDWANA

The early portion of Early Permian was marked by an intense period of glaciation both in Argentina (Asselian to Early Sakmarian, González 2002) and Brazil (Asselian-Sakmarian; Holz *et al.* 2008; see Fig. 2) that may have restricted the occurrences of *Oriocrassatella* in southern South America. During the Early Permian (Cisuralian) there are only two scattered occurrences of the genus, being one in the post-glacial interval in western Argentina (Upper Asselian-Lower Sakmarian) and a second one in Brazil in post-glaciation rocks (Upper Sakmarian-Lower Artinskian, see both occurrences in Figures 1B and 2[5, 15]).

During the Early Permian, *Oriocrassatella* is well recorded in deposits of central Gondwana fringe (Oman, Afghanistan and India; Figs 1B, 2[11, 13a-c, 14a-c]) with seven species assigned. Species described in Oman and Afghanistan occur out of

glacial areas of perigondwanan regions during the Asselian and Sakmarian (Angiolini *et al.* 2007, 2009). Larghi (2005) described a dubious *Oriocrassatella* (see Figs 1B, 2[14c]) in the *Dickinsartella* fauna of the Saiwan Formation, Oman. According to this author, this fauna is indicative of climatic amelioration in North-Eastern Gondwanan fringe after the end of the Early Permian glacial events. Angiolini *et al.* (2006) stated that the progressive increase of carbonate deposits (Haushi Formation) where *Oriocrassatella* was recorded by Dickins & Shah (1979; see Figs 1B, 2[14b]) is correlative to this time of climatic warming during the Upper Sakmarian. Indeed, based on oxygen isotope composition of brachiopod shells Angiolini *et al.* (2009) suggested that low-latitude Early Permian ocean waters did not undergo significant cooling during the final episode of Gondwanan glaciations (Glacial III).

The Carboniferous and Permian glacial interval in India is just roughly understood from the contributions of Sen (1991), Sen & Banerji (1991) and Maejima *et al.* (2001). It seems that all occurrences known from India are present in rocks deposited after the glacial interval (Figs 1B, 2[13a-c, 14a]).

Three occurrences of *Oriocrassatella* are scattered throughout Permian in Eastern Australia in both glacial and interglacial intervals. However, this should be viewed with caution since the dating of some occurrences is not precise, lacking resolution to attribute them to worldwide accepted Permian stages (Figs 1B, C, 2[12, 16, 33]). In Western Australia and peripheral regions the genus is known only from post-glaciated deposits (Figs 1B, C, 2[17, 31, 34-36]), except by the Early Permian occurrence from Timor (Figs 1, 2[18]) an area apparently free from glaciers during this time (Angiolini *et al.* 2009).

THE PERMIAN FOSSIL RECORD OF *ORIOCRASSATELLA* IN LAURUSSIA

The oldest known record of the genus in the Laurussia refers to an isolated Middle Carboniferous occurrence in Kazakhstan (Muromtzeva 1974; Figs 1A, 2[10]), followed by an extensive hiatus throughout the entire Pennsylvanian. During the Early Permian (Asselian to Sakmarian) *Oriocrassatella* is present in only two localities in Laurussia (see

Figs 1B, 2[19, 20a, b]). After that, another gap in the stratigraphical distribution of the genus occurs during the entire Artinskian to Early Kungurian.

During the Upper Kungurian and Roadian the genus reached its maximum paleobiogeographic distribution in Laurussia (Figs 1B, 2[21-29]). Only one species was recorded in deposits of Early Lopingian age (Figs 1C, 2[30]) in the end of the Middle Permian and beginning of the Upper Permian. Indeed, the genus is declined significantly in the Upper Permian rocks of Laurussia.

SUMMARY AND FINAL REMARKS

The geographic and age distribution of the occurrences in Late Mississippian deposits of Australia and Argentina may indicate an earliest Visean or even a pre-Visean origin in Gondwana for the genus *Oriocrassatella* (see Fig. 2). Hence, the current knowledge of its distribution in space and time strongly supports Archbold *et al.* (1996) view, indicating a complex history of migration through time, from the Southern Hemisphere to Northern Hemisphere during the Permian.

Although the earliest known stratigraphic record of the genus appears in eastern Australia, a greater number of species is known from Lower and Upper portions of Carboniferous in South America indicating a putative South American origin for the genus during the earliest (Tournasian-Visean) Carboniferous times (see Fig. 2).

The earliest but isolated occurrence of the genus in eastern Gondwana (eastern Australia, see Figs 1A, 2[1]) during Early Carboniferous (Upper Visean-Lower Serpukhovian) may equally indicate Australia as the site of origin of *Oriocrassatella*. The genus appears to have originated during the Carboniferous in cold waters as seen by its distribution during this time in Argentina and Australia (see Mississippian occurrences in Fig. 1A). However, after this, during Pennsylvanian and Permian times the genus appears to have dispersed to warm waters of the Gondwana peripheral regions and Laurussia. However, exceptions to the pattern above are also known. For example, occurrences are recorded from Carboniferous and Permian of eastern Australia (Figs 1, 2[1, 9, 12,

16, 36]) within glacial intervals. This may indicate that species of this genus did not obligatorily avoided cold water settings. Indeed, González (2002) included *Oriocrassatella* as a genus commonly present in both cold and warm waters in faunas from the Pennsylvanian of western Argentina. In accordance to this interpretation, *Oriocrassatella piauiensis* n. sp. from the Piauí Formation is associated with a well diversified warm water fauna (Anelli *et al.* 2002, 2006, 2009). Moreover, Permian occurrences of *Oriocrassatella* in South America fit well with non glaciated areas during the upper part of Sakmarian stage in Argentina (Figs 1, 2[5]), Brazil (Figs 1, 2[15]) and Middle Permian in Venezuela (Northern South America; Figs 1, 2[32a, b]).

For that reason, the existence of the oldest known species of *Oriocrassatella* in Gondwana (*Oriocrassatella* sp., *O. compressa*, *O. piauiensis* n. sp. and *O. andina*) during the Carboniferous glacial period, in proximity to the glaciated areas (as *O. piauiensis* n. sp.), suggests that the origin of this genus was close to the cold seas of the southern hemisphere. This is not surprising as many lineages of bivalves appeared during the Carboniferous glaciations within the *Levipustula levis* fauna in central Patagonia, especially the endemic genera *Pyramus* Dana, 1847, *Megadesmus*? Sowerby, 1839, *Myonia* Dana, 1847, *Aphanaia* de Koninck, 1877, *Kolymia* Likharev, 1941, *Merismopteria* Etheridge, 1892 (see González 1972, 2002) and more rarely in Australia (Runnegar 1972). Also the precursors of the atomodesmids appear to have adapted to warm sea waters, during their migration to the northern hemisphere in the Permian (González & Waterhouse 2012). Similarly, *Oriocrassatella* younger species are everywhere associated with warm climate (Archbold *et al.* 1996; Figs 1, 4). We cannot rule out the possibility that the first species could be more tolerant to different temperatures, or perhaps *O. piauiensis* n. sp. was an early adaptation to warm waters, but certainly species of this genus adapted definitely to warm water environments during the Late Pennsylvanian interglacial. This warming eliminated the climatic barrier that interrupted the faunal interchange during the “Middle” Carboniferous glacial period and favored larval dispersion following a South to

North migration route through the eastern margins of Gondwana and the eastern Paleotethys (Shi & Grunt 2000; González & Waterhouse 2012). It should be noted, however, that the lack of fossil evidence in intermediate regions (as in western Antarctica, see Kelly *et al.* 2001) is not an evidence of absence. Through the western margin of Gondwana migratory sense may have varied in the course of the Late Paleozoic.

The Late Pennsylvanian time elapsed without records of *Oriocrassatella* (and the Gondwana fauna), except for occurrences in Argentina (see Fig. 1[6-8]), may be related to the generalized marine regressions in Gondwana (see Roberts *et al.* 1995; Dickins 1996). In addition, the pattern of occurrences described above for the Late Pennsylvanian may be a result of taphonomic bias and/or incomplete sampling. However, a preservational bias is difficult to conceive in some cases, as in the lack of *Oriocrassatella* records during the Serpukhovian-Moscovian stages in Australia, because the fossil record is accompanied by marine deposits yielding members of the *Levipustula levis* zone (see Fig. 1A). It is noteworthy that *Oriocrassatella* is also apparently absent in the same interval in Carboniferous deposits of the Alexander Island, Antarctica (Kelly *et al.* 2001).

The Carboniferous *Oriocrassatella* from Kazakhstan, Laurussia (Muromtzeva 1974 in Biakov 2005) may indicate a pre-Permian migration from Gondwana to northern Laurussia, but Early Carboniferous occurrences half way in central Pangea sequences are still missing. Hence, *Oriocrassatella* has a patchy distribution during Carboniferous. Finally, only during Permian times the genus spread-out through Pangea when the worldwide distribution was reached.

A NEW CARBONIFEROUS SPECIES OF *ORIOCRASSATELLA* FROM BRAZIL

GENERAL OBSERVATIONS

Specimens of *Oriocrassatellinae* described here were found in a dolostone bed informally named as “Mucambo”, which crops out near the village of José de Freitas (Fig. 3), Piauí. The specimens were collected and prepared according to standard paleontological techniques (Feldmann *et al.* 1989). For all




FIG. 3. — **A**, map showing the area of occurrence of the Upper Carboniferous Piauí Formation (in black) in the Parnaíba Basin, north-eastern Brazil; **B**, localization of the outcrop of Mucambo dolostone; **C**, stratigraphic section showing the Mucambo dolostone.

specimens, latex casts were prepared for assessment of muscle scars and especially for hinge features. Additionally, some latex casts of hinge features were photographed with scanning electronic microscope (SEM). The studied specimens are housed in the scientific collection of the Instituto de Geociências, Universidade de São Paulo, under the code GP/1T. In addition, some specimens belonging to the Departamento Nacional da Produção Mineral (DNPM), Rio de Janeiro, were also analyzed. The suprageneric systematics is based on Boyd & Newell (1968) and Amler (1999). The elongation and obesity indexes were calculated according to Stanley (1970).

SYSTEMATIC PALEONTOLOGY

Family CRASSATELLIDAE

Férussac, 1822

Subfamily ORIOCRASSATELLINAE

Boyd & Newell, 1968

Genus *Oriocrassatella* Etheridge Jr., 1907

Procrassatella Yakovlev, 1928: 114 (type species: *Schizodus planus* Golovkinsky, 1868, by monotypy).

TYPE SPECIES. — *Oriocrassatella stokesi* Etheridge Jr., 1907, by monotypy.

TABLE 1. — Dimensions (in mm) of *Oriocrassatella piauiensis* n. sp. Abbreviations: l, length; h, height; w, width; *, approximate for incomplete valve.

specimen	valve	length	height	width	elongation l/h	obesity h/w	umbonal angle (°)
GP/1T 2026	L	24	17.5	4	1.37	2.1	132
DNPM 219	L	24	17	4	1.41	2.1	124
DNPM 271	R	23	15.5	4	1.48	1.94	130
GP/1T 2022	R	21	13	4	1.61	1.62	132
DNPM 220	L	20	14.9	3	1.41	2.5	131
DNPM 148	–	20	13	3	1.54	2.2	131
GP/1T 2023	R	17	13	3	1.30	2.2	125
DNPM 234	R	17*	13	3	1.31*	2.2	129
GP/1T 2028	L	17	11	2.5	1.54	2.2	134
GP/1T 2024	R	14.5	10	2.5	1.45	2.0	129
DNPM 221a	L	14.2	9.5	2.5	1.49	1.90	129
GP/1T 2027	L	14	9	2	1.55	2.2	128

Oriocrassatella piauiensis n. sp.
(Fig. 4)

TYPE MATERIAL. — Holotype GP/1T 2022. — Paratypes GP/1T 2023, 2024, 2026–2028; DNPM 148, 219, 220, 221a, 234, 271.

ETYMOLOGY. — The specific epithet refers to Piauí State, northern Brazil, where the species was found.

TYPE HORIZON. — Mucambo dolostone, Piauí Formation, Parnaíba Basin.

AGE. — Bashkirian to Moscovian. The time-range assigned herein comprises the more reliable relative ages to the Piauí Formation, obtained from palynological data (Dino & Playford 2002; Souza *et al.* 2010) and comparisons with the palynostratigraphical scheme erected to the neighbor Amazonas Basin (Playford & Dino 2000).

MATERIAL EXAMINED. — Six internal molds of left valves and six of right valves; two external molds of left valves.

MEASUREMENTS. — See Table 1.

TYPE LOCALITY. — José de Freitas, (4°53'656"S, 42°35'871"W), Piauí, Brazil.

DIAGNOSIS. — *Oriocrassatella* with lateral teeth absent; non bifid cardinal tooth 2; ornamentation of regularly spaced lamellae with flattened surfaces showing rare bifurcation in anterior and posterior regions of shell.

DESCRIPTION

Internal and external molds up to 24 mm in length, and up to 17.5 mm in height. Shell thick based on internal molds, oval in shape, elongate, very compressed. Umbones depressed, beaks prosogy-

rate slightly elevated above hinge. Umbonal carina weak and rounded, running from beaks to posterior ventral angle, defining a large respiratory margin. As seen in casts of external molds, anterior dorsal margin straight; anterior extremity convex; ventral margin slightly convex; posterior extremity slightly truncated; posterior dorsal margin straight. Ornamentation of concentric lamellae, regularly spaced, about two per millimeter, some showing bifurcating pattern anteriorly and posteriorly. Pedal or siphonal gapes absent. Lunule and escutcheon absent.

Hinge formula as follows:

Posterior	RV	/nsr	–	3b	–	3a/
Anterior	LV	/nsr	4b	–	2	–/

Left anterior cardinal (2) strong, non arched or bifid; left posterior cardinal (4b) weak, laterally compressed, vertical, sometimes showing sigmoid shape. Right anterior cardinal (3a) well defined, separated from the anterior dorsal margin by shallow depression. Right posterior cardinal (3b) straight, vertical, slightly inflated in both extremities. Resilifer (r) in left valve placed behind posterior cardinal (4b), cover deck absent; septum (s) strong, well developed in both valves, originating immediately posterior to beak, bounding resilifer posteriorly; nymph (n) well-defined in posterior wall of septum, close to beak. Ligament furrow (lf) very narrow, located posterior to the septum. Lateral teeth absent.

Adductor muscles scars similar in size. Anterior scar positioned close to anterior dorsal angle, twice higher than long, with irregular shape at posterior




Fig. 4. — *Oriocrassatella piauiensis* n. sp., Mucambo limestone; **A**, holotype, internal mold of right valve, GP/1T 2022; **B**, internal mold of left valve, DNPM 076; **C**, latex cast, internal mold of right valve, DNPM 234; **D**, latex cast, magnified hinge area, internal mold of right valve, same specimen; **E**, latex cast, internal mold of left valve, magnified hinge area, DNPM 221b; **F**, latex cast, internal mold of left valve, same specimen; **G**, internal mold of right valve, GP/1T 2024; **H**, muscle scars based in specimen DNPM 076; **I**, latex cast, external mold of left valve, DNPM 220; **J**, internal mold of right valve, DNPM 234. Abbreviations: **aa**, anterior adductor; **apr**, anterior pedal retractor; **lf**, ligament furrow; **n**, nymphs; **pa**, posterior adductor; **pl**, pallial line; **r**, resilifer; **s**, septum; **ue**, umbonal elevator; **2**, left anterior cardinal; **3a**, right anterior cardinal; **3b**, right posterior cardinal; **4a**, left posterior cardinal. Scale bars: 0.5 cm.

margin. Anterior pedal retractor small, well marked, not fused but placed close to dorsal extremity of the adductor. Posterior adductor located above posterior dorsal angle, nearly circular in shape. Posterior pedal retractor apparently fused to dorsal end of posterior adductor. Pedal elevator scar well-marked, located at tip of beaks. Pallial line well defined, without sinus.

DISCUSSION

Boyd & Newell (1968: 28, 30) included among the diagnostic features for the genus *Oriocrassatella* the presence of a cover deck over the apical part of the resilifer and of a more or less bifid cardinal tooth 2 in left valve. Although both features may be easily removed from shell by taphonomic processes as abrasion or corrosion, our observations of several well preserved specimens from the Piauí Formation indicate that these features are absent in *Oriocrassatella piauiensis* n. sp. Furthermore, the cover deck feature was recorded only in the type species and *O. stokesi* (Newell 1958) from the Upper Permian of Eastern Australia. In the same way, the bifid cardinal tooth 2 is only observed in the type species and a few other Permian species of this genus (*O. itajaiensis* [Rocha-Campos 1970, Brazil], *O. stokesi* [Dickins 1963, Western], and *O. stokesi* [Newell 1958, Eastern]). Therefore, both the cover deck and bifid cardinal tooth 2 are known only in Permian species. If the presence of these characters are used to characterize species of *Oriocrassatella* then we must remove from this genus all species in which those characters are absent. Since both characters are present only in extremely well preserved Permian specimens we decide to assign our species to *Oriocrassatella* rather than erect a new genus.

Oriocrassatella species from Argentina comparable to *O. piauiensis* n. sp., include *O. sanjuanina* González, 1976, from Upper Carboniferous, which has very similar general profile, but differs by its more posteriorly developed resilifer in right valve, and by the presence of the anterior lateral tooth AII. *Oriocrassatella andina*, from the Late Mississippian-Early Pennsylvanian La Capilla Formation (González 2002) differs also by the presence of the lateral tooth AII and by its ornament of concentric striations.

Oriocrassatella sp. from Rio del Peñón Formation (González 1997) does not show hinge features for close comparison, but has a more elongated shell and more strongly truncated posterior extremity. The anterior pedal retractor is fused to the anterior adductor. *Oriocrassatella* sp. from Del Salto Formation (Manceñido *et al.* 1976) is more circular in shape and has all four lateral teeth in left valve.

Oriocrassatella itajaiensis described by Rocha-Campos (1970: 11, 12, fig. 3), has a more expanded septum in hinge, and although not completely visible, may have anterior and posterior lateral teeth in left valve. Finally, *O. elongate* Boyd & Newell, 1968 described from Permian rocks of Venezuela (Sanchez 1984) and North America (Boyd & Newell 1968) is more elongated in shape, and presents the complete set of lateral teeth in both valves.

Acknowledgements

This research was financially supported by the following scientific agencies: Conselho Nacional de Desenvolvimento Científico e Tecnológico (59969/92-3, 483947/2007-5), and Fundação de Amparo à Pesquisa do Estado de São Paulo (1999/10797-4). MGS and PAS are CNPq fellows. We also thank Suzana Damborenea, Claude Babin and Annemarie Ohler for their comments and suggestions that improved the early version of the manuscript, Tatiana Viktorovna Van Iten and Alexander Biakov for their help in translating parts of Russian papers.

REFERENCES

- AMLER M. R. W. 1999. — Synoptical classification of fossil and Recent Bivalvia. *Geologica et Palaeontologica* 33: 237-248.
- ANELLI L. E. 1999. — *Invertebrados neocarboníferos das formações Piauí (Bacia do Parnaíba) e Itaituba (Bacia do Amazonas): taxonomia; análise cladística das subfamílias Oriocrassatellinae (Crassatellacea, Bivalvia) e Neospiriferinae (Spiriferoidea, Brachiopoda)*. Programa de Pós-Graduação em Geologia Sedimentar, Universidade de São Paulo, Tese de Doutorado, 184 p. (unpublished).
- ANELLI L. E., ROCHA-CAMPOS A. C. & SIMÕES M. G. 2002. — Protobranch bivalves from the Piauí Forma-

- tion (Middle Pennsylvanian), Parnaíba Basin, Brazil. *Revista Española de Paleontología* 17: 165-176.
- ANELLI L. E., ROCHA-CAMPOS A. C. & SIMÕES M. G. 2006. — Pennsylvanian pteriomorphian bivalves from the Piauí Formation, Parnaíba Basin, Brazil. *Journal of Paleontology* 80 (6): 1125-1141.
- ANELLI L. E., ROCHA-CAMPOS A. C. & SIMÕES M. G., PECK R. L. 2009. — Pennsylvanian Heteroconchia (Mollusca, Bivalvia) from the Piauí Formation, Parnaíba Basin, Brazil. *Revista Brasileira de Paleontologia* 12 (2): 93-112.
- ANGIOLINI L., STEPHENSON M. H. & LEVEN Y. 2006. — Correlation of the Lower Permian surface Saiwan Formation and subsurface Haushi Limestone, central Oman. *GeoArabia* 11: 17-38.
- ANGIOLINI L., GAETANI M., GARZANTI E., MATTEI M., MUTTONI G., STEPHENSON M. H. & ZANCHI A. 2007. — Tethyan oceanic currents and climate gradients 300 m.y. ago. *Geology* 35: 1071-1074.
- ANGIOLINI L., JADOUL F., LENG M. J., STEPHENSON M. H., RUSHTON J., CHENE R. S. & CRIPPA G. 2009. — How cold were the Early Permian glacial tropics? Testing sea-surface temperature using the oxygen isotope composition of rigorously screened brachiopod shells. *Journal of the Geological Society* 166: 1-13.
- ARCHBOLD N. W., SWART R. H. & DICKINS J. M. 1996. — Permian fossil marine invertebrates from the Northern Territory described by Robert Etheridge Junior. *Proceedings of the Royal Society of Victoria* 102 (2): 63-71.
- AZCUY C. L., BERI A., OLIVEIRA M. E. C. M. B., CARRIZO H. A., DIPASQUO M., SARAVIA P. D., GONZÁLEZ C. R., IANNUZZI R., LEMOS V. B., MELO J. H. G., PAGANI A., ROHN R., AMENÁBAR C. R., SABATTINI N. & SOUZA P. A. 2007. — Bioestratigrafía del Paleozoico superior de América del Sur: primera etapa de trabajo hacia una nueva propuesta cronoestratigráfica. *Asociación Geológica Argentina, Publicación Especial* 11: 9-65.
- BIAKOV A. S. 2005. — New species of astartids and pholadomyids (Bivalvia) and beds with fauna from the Lower Permian of the Omolon massif, Northeastern Asia. *Paleontological Journal* 39 (2): 133-140.
- BOYD D. W. & NEWELL N. D. 1968. — Hinge grades in the evolution of Crassatellacean bivalves as revealed by Permian genera. *American Museum Novitates* 2328: 1-52.
- BOYD D. W. & NEWELL N. D. 1977. — An addition to the known geographic range of the Permian pelecypod *Oriocrassatella*. *Contributions to Geology* 16 (1): 55-57.
- CAMPANHA V. & ROCHA-CAMPOS A. C. 1979. — Alguns microfósseis da Formação Piauí (Neocarbonífero), Bacia do Parnaíba. *Boletim do Instituto de Geociências* 10: 57-67.
- CHARLTON T. R. 2001. — Permo-Triassic evolution of Gondwanan eastern Indonesia, and the final Mesozoic separation of SE Asia from Australia. *Journal of Asian Earth Sciences* 19 (5): 595-617.
- CIRIACKS K. W. 1963. — Permian and Eotriassic bivalves of the Middle Rockies. *Bulletin of the American Museum of Natural History* 125: 1-100.
- CLARKE M. J. 1987. — Late Permian (Late Lymingtonian = ?Kazanian) brachiopods from Tasmania. *Alcheringa* 11: 261-289.
- DICKINS J. M. 1961. — Permian pelecypods newly recorded from Eastern Australia. *Palaentology* 4 (1): 119-148.
- DICKINS J. M. 1963. — Permian pelecypods and gastropods from Western Australia. *Bulletin of the Bureau of Mineral Resources, Geology and Geophysics* 217: 59-65.
- DICKINS J. M. 1996. — Problems of a Late Palaeozoic glaciation in Australia and subsequent climate in the Permian. *Palaeoogeography, Palaeoclimatology, Palaeoecology* 125: 185-197.
- DICKINS J. M. & SHAH S. C. 1979. — Correlation of the Permian marine sequences of India and Western Australia, in LASKAR B. & RAJA RAO C. S. (eds), *Fourth International Gondwana Symposium: Papers*. Vol. 2. Hindustan Publishing Corporation, Delhi: 387-408.
- DICKINS J. M. & SKWARKO S. K. 1981. — Upper Paleozoic pelecypods and gastropods from Irian Jaya, Indonesia. *Geological Research and Development Centre, Bandung, Paleontology Series* 2: 43-52.
- DICKINS J. M., ARCHBOLD N. W., THOMAS G. A. & CAMPBELL H. J. 1989. — Mid-Permian correlation, in *XI^e Congrès international de stratigraphie et de géologie du carbonifère, Beijing, Compte Rendu*. Tome 2: 185-198.
- DINO R. & PLAYFORD G. 2002. — Stratigraphic and palaeoenvironmental significance of a Pennsylvanian (Upper Carboniferous) palynoflora from the Piauí Formation, Parnaíba Basin, northeastern Brazil. *Paleontological Research* 6 (1): 23-40.
- ETHERIDGE R. JR. 1907. — Official contributions to the paleontology of South Australia. *Adelaide, South Australia House of Assembly, Supplement to Parliamentary Paper* 55: 1-353.
- FELDMANN R. M., CHAPMAN R. E. & HANNIBAL J. T. 1989. — Paleotechniques. *Paleontological Society Special Publication* 4, 358 p.
- FIELDING C. R., FRANK T. D., BIRGENHEIER L. P., RYSEL M. C., JONES A. T. & ROBERTS J. 2008a. — Stratigraphic imprint of the Late Palaeozoic Ice Age in eastern Australia: a record of alternating glacial and nonglacial climate regime. *Journal of the Geological Society* 165: 129-140.
- FIELDING C. R., FRANK T. D., BIRGENHEIER L. P., RYSEL M. C., JONES A. T. & ROBERTS J. 2008b. — Stratigraphic record and facies associations of the Late Paleozoic Ice Age in eastern Australia (New South Wales and Queensland), in FIELDING C. R., FRANK T. D. & ISBELL J. L. (eds.), *Resolving the Late Paleozoic Ice Age in Time and Space. Geological Society of America, Special Paper* 441: 41-57.

- GOLOVINSKY N. A. 1868. — [The Permian Formation of the Central Kama-Volga Basin] O Permskoi format-sii v'tsenral'noi chasti Kamsko-Volshskago Basseina. *Materialy po Geologii Rossii* 1: 275-415 (in Russian).
- GONZÁLEZ C. R. 1972. — La Formación Las Salinas, Paleozoico superior de Chubut (República Argentina). Parte II. Taxonomía y Paleoecología. *Revista Asociación Geológica Argentina* 27 (2): 188-213.
- GONZÁLEZ C. R. 1976. — *Oriocrassatella* y *Stutchburia* (Bivalvia) en la ingression marina del Permico inferior de la Quebrada de Agua Negra, Depto. Iglesia, Provincia de San Juan (Argentina). *Ameghiniana* 13 (2): 127-140.
- GONZÁLEZ C. R. 1981. — El Paleozoico superior marino de la República Argentina. Bioestratigrafía y Paleoclimatología. *Ameghiniana* 18 (1/2): 51-65.
- GONZÁLEZ C. R. 1982. — Bivalvios de la Formación Agua del Jagüel, Permiano inferior de Mendoza (Argentina). *Ameghiniana* 19 (3-4): 367-376.
- GONZÁLEZ C. R. 1990. — Development of the Late Paleozoic glaciations of the South American Gondwana in western Argentina. *Palaeogeography, Palaeoclimatology, Palaeoecology* 79: 275-287.
- GONZÁLEZ C. R. 1997. — Late Carboniferous bivalvia from western Argentina. *Geologica et Paleontologica* 31: 193-214.
- GONZÁLEZ C. R. 2002. — Bivalves from Carboniferous glacial deposits of western Argentina. *Palaontologische Zeitschrift* 76 (1): 127-148.
- GONZÁLEZ C. R. & DÍAZ-SARAVIA P. 2007. — Faunas y Paleoclimatología del Paleozoico superior de Argentina: revisión crítica. *Acta Geológica Lilloana* 20 (1): 41-72.
- GONZÁLEZ C. R. & WATERHOUSE J. B. 2012. — Carboniferous atomodesmids (Mollusca) from central Patagonia, Argentina. *Geologica et Palaeontologica* 44: 27-36.
- HOLZ M., SOUZA P. A. & IANNUZZI R. 2008. — Chapter 8: Sequence stratigraphy and biostratigraphy of the Late Carboniferous to Early Permian glacial succession (Itararé subgroup) at the eastern-southeastern margin of the Paraná Basin, Brazil, in FIELDING C. R., FRANK T. D. & ISBELL J. L. (eds), Resolving the Late Paleozoic Ice Age in Time and Space. *Geological Society of America, Special Paper* 441: 115-129.
- HOLZ M., FRANÇA A. B., SOUZA P. A., IANNUZZI R. & ROHN R. 2010. — A stratigraphic chart of the Late Carboniferous/Permian succession of the eastern border of the Paraná Basin, Brazil, South America. *Journal of South American Earth Sciences* 29 (2): 381-399.
- HONEGGER K., DIETRICH V., FRANK W., GANSSER A., THÖNI M. & TROMMSDORFF V. 1982. — Magmatism and metamorphism in the Ladakh Himalayas (the Indus-Tsangpo suture zone): *Earth and Planetary Science Letters* 60: 253-292.
- JIN X. 2002. — Permo-Carboniferous sequences of Gondwana affinity in southwest China and their paleogeographic implications. *Journal of Asian Earth Sciences* 20 (6): 633-646.
- KANEV G. P. 1994. — [Permian Bivalves from the Chernyshev Range] *Permiskie dvustvorchatye Mollyuski gryady Chernysheva*. Rossiiskaya Akademya Nauk, Ural'skoe Otdelenie, Komi Nauchnyi Tsentr, Institut Geologii, Syktyvkar: 1-80 (in Russian).
- KELLY S. R. A., DOUBLEDAY P. A., BRUNTON C. H. C., DICKINS J. M., SEVASTOPULO G. D. & TAYLOR P. D. 2001. — First Carboniferous and ?Permian marine macrofaunas from Antarctica and their tectonic implications. *Journal of the Geological Society* 158 (2): 219-232.
- LARGHI C. 2005. — *Dickinsartella* fauna from the Saiwan Formation (Oman): a bivalve fauna testifying to the late Sakmarian (Early Permian) climatic amelioration along the north-eastern Gondwanan fringe. *Rivista Italiana di Paleontologia e Stratigrafia* 111: 353-375.
- LIMA FILHO F. P. 1991. — *Facies e ambientes deposicionais da Formação Piauí (Pensilvaniano), Bacia do Parnaíba, Brasil*. Programa de Pós-Graduação em Geologia Sedimentar, Universidade de São Paulo, Dissertação de Mestrado, 137 p. (unpublished).
- LIMA FILHO F. P. 1999. — *A seqüência Permo-Pensilvaniana da Bacia do Parnaíba*. Programa de Pós-Graduação em Geologia Sedimentar, Universidade de São Paulo, Tese de Doutorado, 155 p. (unpublished).
- LIU N. P. & CUI X. S. 1983. — Discovery of *Eurydesma*-fauna from Rutog, northwest Xizang (Tibet), and its biogeographic significance. *Earth Science, Journal of Wuhan College of Geology* 1: 79-92.
- LOBANOVA O. V. 1961. — [Permian bivalves from the Grodekovskii District (Western Prymorie)] *Permskie peletsipody Grodekovskogo raiona zapadnogo Prymor'ya. Sbornik Statei po Paleontologii i Biostratigrafii* 25: 52-80 (in Russian).
- MAEJIMA W., DAS R., PANDYA K. L. & HAYASHIL M. 2001. — Post-Glacial Sedimentation of Talchir Formation, Talchir Gondwana Basin, Orissa, India. *Gondwana Research* 4 (4): 694-695.
- MANCENIDO M. O., GONZÁLEZ C. R. & DAMBORENA S. E. 1976. — La fauna de la Formación del Salto (Paleozoico superior de la Provincia de San Juan). Parte III: Bivalvia 1. *Ameghiniana* 13 (1): 65-84.
- MAXWELL W. G. H. 1964. — The geology of the Yarrol Region. Part 1. Biostratigraphy. *Department of Geology University of Queensland, Papers* 5 (9): 1-79.
- MUROMTSEVA V. A. 1974. — Bivalves from the Carboniferous of Kazakhstan and Siberia, *Trudy Vnigri* 336: 1-150.
- MUROMTSEVA V. A. & GUSKOV V. A. 1984. — [Permian Marine Deposits and Bivalve Mollusks of the Soviet Arctic] *Permiskie morskije otlozheniya i dvustvorchatye mollyuski Sovetskoy Arktiki*. Trudy Vsesoyuznyi ordena Trudovogo Krasnogo Znameni Neftyanoi Nauchno-Issledovatel'skii Geologorazvedochnyi Institut, Nedra,

- Leningrad: 1-154, pls 1-53 (in Russian).
- NEWELL N. D. 1955. — Permian pelecypods of East Greenland. *Meddelelser om Gronland* 110 (4): 1-36.
- NEWELL N. D. 1958. — A note on Permian Crassatellid pelecypods. *American Museum Novitates* 1878: 1-6.
- PEOU S. & ENGEL B. A. 1979. — A Carboniferous fauna from Rawdon Vale, New South Wales. *Alcheringa* 1: 141-157.
- PLAYFORD G. & DINO R. 2000. — Palynostratigraphy of upper Palaeozoic strata (Tapajós Group), Amazonas Basin, Brazil. Part 2. *Palaeontographica Abteilung B* 255: 87-145.
- REED F. R. C. 1932. — New fossils from the agglomeratic slate of Kashmir. *Palaeontologia Indica* 20 (1): 1-106.
- ROBERTS J., HUNT J. W. & THOMPSON D. M. 1976. — Late Carboniferous marine invertebrate zones of eastern Australia. *Alcheringa* 1: 197-225.
- ROBERTS J., CLAQUE-LONG J., JONES P. J. & FOSTER C. B. 1995. — SHRIMP zircon age control of Gondwanan sequences in Late Carboniferous and Early Permian Australia, in DUNAY R. E., HAILWOOD E. A. (eds), Non-Biostratigraphical Methods of Dating and Correlation. *Geological Society, Special Publication* 89: 145-174.
- ROCHA-CAMPOS A. C. 1970. — Moluscos permianos da Formação Rio Bonito (Subgrupo Guatá), SC. *Boletim do Departamento Nacional de Produção Mineral, Divisão de Geologia e Mineralogia*, 251:1-89.
- RUNNEGAR B. 1972. — Anatomy of *Pholadomya candida* (Bivalvia) and the origin of the Pholadomyidae. *Proceedings Malacological Society of London* 40: 45-50.
- SABATTINI N., OTTONE E. & AZCUY C. 1990. — La zona de *Lissochonetes jachalensis-Streptorhynchus inaequiorinatus* (ex Fauna Intermedia) en la localidad de La Delfina, provincia de San Juan. *Ameghiniana* 27: 1, 2, 75-81.
- SANCHEZ M. T. 1984. — *Étude paléocologique du Paléozoïque supérieur dans le nord des Andes du Venezuela*. Université de Bretagne occidentale, Brest, collection *Biostratigraphie du Paléozoïque* 2: 1-165.
- SEN D. P. 1991. — Sedimentation patterns of the Talchir Group in the Giridih Gondwana basin, India: a case of multiple glacial advance and retreat. *Palaeogeography, Palaeoclimatology, Palaeoecology* 86: 339-352.
- SEN D. P. & BANERJI T. 1991. — Permo-Carboniferous proglacial-lake sedimentation in the Sahajuri Gondwana basin, India. *Sedimentary Geology* 71: 47-58.
- SHI G. R. & GRUNT T. A. 2000. — Permian Gondwana-Boreal antitropicality with special reference to brachiopod faunas. *Palaeogeography, Palaeoclimatology, Palaeoecology* 155: 239-263.
- SHI G. R., ARCHBOLD N. W., ANGIOLINI L., GRUNT T. A., KOTLYAR G. V., METCALFE I., SHEN S., WANG X., SRIVASTAVA A. K. & ZAKHAROV Y. D. 2002. — Working Group on "Using Transitional Biotas as Gateways for Global Correlation" and the "Permian Research Group of SE Asia", in Permophiles International Commission on Stratigraphy International Union of Geological Sciences Newsletter of the Subcommittee on Permian Stratigraphy Number 40, Joint Report of SPS.
- SOUZA P. A., MATZEMBACHER L. T., ABELHA M. & BORGHI L. 2010. — Palinologia da Formação Piauí, Pensilvaniano da Bacia do Parnaíba: biocronoestratigrafia de intervalo selecionado do poço 1-UN-09-PI (Caixas, MA, Brasil). *Revista Brasileira de Paleontologia* 13 (1): 57-66.
- STANLEY S. M. 1970. — Relation of shell form to life habits of the Bivalvia (Mollusca). *Memoir of the Geological Society of America* 125: 296 p.
- STEPHENSON M. H., ANGIOLINI L. & LENG M. J. 2007. — The Early Permian fossil record of Gondwana and its relationship to deglaciation: a review, in WILLIAMS M., HAYWOOD A. M., GREGORY F. J. & SCHMIDT D. N. (eds), *Deep-Time Perspectives on Climate Change: Marrying the Signal from Computer Models and Biological Proxies*. The Micropalaeontological Society, Special Publications, Geological Society, London, p. 103-122.
- STERREN A. F. 2005. — Bivalvos carboníferos de la Formación La Capilla en el área de Las Cambachas, provincia de San Juan, Argentina. *Ameghiniana* 42 (1): 209-219.
- SUVEIZDAS P., MIKLUKHO-MAKLAY K. V., UKHARSKAYA L. B., MOROZOVA I. P., GLUSHENKO N. V., IVANOV V. K. & PODOBA B. G. 1975. — [Permian deposits of the Baltic region (stratigraphy and fauna)]. Permskaya sistema Pribaltiki (fauna i stratigrafiya). *Lietuvos Geologijos Mokslinio Tyrimo Institutas, Trudy* 29, 305 p.
- TABOADA A. C. 1997. — Bioestratigrafía del Carbonífero marino del valle Calingasta-Uspallata, provincias de San Juan y Mendoza. *Ameghiniana* 34 (2): 215-246.
- TERMIER G., TERMIER H., LAPPARENT A. F. DE & MARIN P. 1974. — Monographie du Permo-Carbonifère de Wardak (Afghanistan Central). *Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon, Hors Série* 2: 1-167.
- VANDERHOR F., FLINT R. B., TYLER I. M. & HOCKING R. M. 2006. — 1:500 000 interpreted bedrock geology of Western Australia, 2006 update. Dataset within GeoVIEW.WA (interactive geological map) *Geological Survey of Western Australia map*.
- VEEVERS J. J. 1995. — Gondwana master basin of Peninsular India between Tethys and the interior of the Gondwanaland Province of Pangea, Boulder, Colorado. *Geological Society of America Memoir* 187, 72 p.
- VEEVERS J. J. & TEWARI R. C. 1995. — Permian-Carboniferous and Permian-Triassic magmatism in the rift zone bordering the Tethyan margin of southern Pangea. *Geology* 23 (5): 467-470.
- WANNER C. 1940. — *Neue Permische Lamellibranchiaten von Timor*, in BROUWER H. E. (ed.), Geological Expedition of the University of Amsterdam to the Lesser Sunda Islands in the Southeastern part of the

- Netherlands East Indies 1937 under leadership of H. A. Brouwer 2: 369-395.
- WATERHOUSE J. B. & BRIGGS D. J. C. 1986. — Late Palaeozoic Scyphozoa and Brachiopoda (Inarticulata, Strophomenida, Productida and Rhynchonellida) from the southeast Bowen Basin, Australia. *Palaeontographica Abteilung A* 193 (1-4): 1-76.
- YAKOVLEV N. N. 1902. — [Paleozoic representative of the Crassatellidae (*Schizodus planus* Golowk.)] Paleozoiskii predstavitel' Crassatellidae. (*Schizodus planus* Golowk.). *Izvestiya Geologicheskogo Komiteta* 21 (9): 755-760 (in Russian).
- YAKOVLEV N. N. 1928. — [Procrassatella, a new genus from Permian] *Procrassatella*, novyi rod iz Permi. *Ezhegodnik Russkogo Paleontologicheskogo Obshchestva* 7: 119-125 (in Russian).

*Submitted on 1st December 2010;
accepted on 19 April 2011.*