

























Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member C







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8



























































































































































































Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member E







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8

































































































































































Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member F







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8









































































Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member Lower G







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8



































































































































































































Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member H







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8







































Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member J







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8















































Photosimulations and false color elevation maps of scanned facets on molars of extinct *Kolpochoerus* from the Shungura Formation (Ethiopia), Member L







scanned at the PALEVOPRIM lab by A. Souron, University of Poitiers, France with "TRIDENT", a white light confocal microscope Leica DCM8














































































































140x100-Zinv-Kol-OMO377-101...stal-LUM3-lophX2-cb-fc-milieu_c



140x100-Zinv-Kol-OMO377-101...stal-LUM3-lophX2-cb-fc-milieu_c



140x100-Zinv-Kol-OMO377-101...stal-LUM3-lophX2-cb-fc-milieu_d



140x100-Zinv-Kol-OMO377-101...stal-LUM3-lophX2-cb-fc-milieu_d

